

Sensitivity of satellite NO₂ retrievals to aerosols

J. Leitão, A. Heckel, A. Richter, A. Kokhanovsky, and J.P. Burrows
Institute of Environmental Physics/Remote Sensing, University of Bremen, Germany
Email: jleitao@iup.physik.uni-bremen.de



Introduction

Satellite instruments such as GOME, GOME-2, SCIAMACHY and OMI perform measurements of the backscattered solar radiation from which trace gas distributions in the atmosphere can be retrieved. These data are of growing importance to investigate the global distribution of pollutants such as nitrogen dioxide (NO₂), sulphur dioxide (SO₂) and several others. With the satellite derived global fields one can identify emission sources and analyse long-term trends of pollutant concentrations.

The retrieval of tropospheric columns of NO₂ from satellite measurements is based on several assumptions that in one way or another contribute to the uncertainty in the final retrieval. The improvement of the a priori assumptions used for the computation of the airmass factor (AMF) is a main concern to obtain the correct values of NO₂ present in the troposphere.

A sensitivity study is shown in this poster as an example of the impact of a **different aerosol characteristics** such as optical properties and optical thickness (AOD) on the computation of the NO₂ AMF with the radiative transfer model (RTM) Sciatran (Rozanov *et al.*, 2005). Also the effect of **different boundary layer heights** values is studied.

Motivation...

It is known that the **presence of aerosols** (and its scattering effect) has an impact on the measurements of tropospheric NO₂. This effect is related for example to the vertical distribution of aerosol and its type.

Currently in IUP-Bremen, the NO₂ retrieval method uses data taken from climatological assumptions (Richter *et al.*, 2005). Therefore, the full spatial and time variability of aerosols and its characteristics is not well accounted. Hence, we attempt to include in the retrieval method data that can be continuously updated, i.e. using independent measurements when available and dynamical "climatology" when it is necessary to fill gaps. This can be done using, for example, AOD values calculated with the BAER algorithm (von Hoyningen-Huene *et al.*, 2006) from MERIS data.

Sensitivity Study - settings

Radiative transfer calculations

- RTM: Sciatran 2.2
- Surface albedo = 0.03
- Wavelength: 425, 437.5, 440, 450nm
- SZA: 20°, 35°, 60°, 70°
- BL heights: 0.6, 1, 2 and 3 km
- NO₂ and aerosol profiles – "box profile" of well mixed gas and aerosol along BL considered (see Fig.1)




Figure 1: Profiles of NO₂ for different BL heights. The aerosol profile is represented by the shaded area.

Aerosol Settings:

Phase function calculated with a FORTRAN program developed by Michael Mishchenko (de Rooij *et al.*, 1984; Mishchenko *et al.*, 1999) and adapted by Alexander Kokhanovsky.

• Optical properties of aerosols taken from Dubovik *et al.* (2002) – measurements from 12 worldwide AERONET stations

• "Typical" compositions of aerosol considered: urban/industrial, biomass burning, desert dust and oceanic (according to location)

• Legendre expansion coefficients determined for fine and coarse particles

• Wavelength: 440nm

• AOD: 0.1, 0.5, 0.9

• Aerosol absorption profile was determined according to the extinction coefficient profile – single scattering albedo of 0.933 (Dubovik *et al.*, 2002)

Sensitivity Study - results

Here only the results obtained with data measured in the AERONET station **Creteil-Paris, France** are shown. However, NO₂ AMF were calculated for several locations but the difference of the results is not so significant (see the example in Fig.2 on the right for stations classified as "desert" and "urban").




Figure 2: NO₂ AMF at 440nm, for BL=1km, with AOD=0.5. Aerosol optical properties taken from AERONET stations data in urban (in blue) and desert (in yellow/orange) background (Dubovik *et al.*, 2002).






Figure 3: NO₂ AMF at 437.5nm for AOD of 0.1 (top), 0.5 (middle) and 0.9 (bottom). AMF showed for different BL and for absorbing (Wabs) and non-absorbing aerosol (Nabs):

• Wabs BL600, Wabs BL1000, Wabs BL2000, Wabs BL3000

• Nabs BL600, Nabs BL1000, Nabs BL2000, Nabs BL3000

• Fine aerosol well mixed with NO₂ will derive higher AMF than those calculated when considering coarse particles (Fig.4). This difference is more pronounced for high aerosol loads (e.g. AOD=0.9).

• NO₂ signal is amplified for larger SZA and longer wavelengths.

• In general NO₂ AMF increases with the extension of BL to higher altitudes (Fig.3).

• The increase of AOD will lead to higher NO₂ AMF. Yet, in some cases, with AOD=0.9 it is possible to observe a decrease of AMF for SZA greater than 60° (Fig.4).

• When considering absorbing aerosol, the NO₂ AMF will be lower. The difference to the non-absorbing one is more relevant for high AOD values (Fig.3).




Figure 4: NO₂ AMF with profiles of BL=1km, at 425nm and 440nm and for several AOD values (0.1, 0.5 and 0.9). Coarse mode aerosol is represented by solid line and fine particles with hatched line.

Acknowledgements

- Part of this project is funded by the European Community through the GEMS project.
- Aerosol data from measurements performed at AERONET stations.

References

- Dubovik, O., B. Holben, T.F. Eck, A. Smirnov, Y. Kaufman, M.D. King, D. Tanré, I. Slutsker; *Variability of absorption and optical properties of key aerosol types observed in Worldwide locations*; J. Atmospheric Sciences; 59; 590-608; doi:10.1175/1520-0469(2002)059<0590:VOAOP>2.0.CO;2; 2002.
- Mishchenko, M. A., J. M. Dlugach, E. G. Yanovitskij, and N. T. Zakharova; *Bidirectional reflectance of flat, optically thick particulate layers: an efficient radiative transfer solution and applications to snow and soil surfaces*; J. Quant. Spectrosc. Radiat. Transfer; 63; 409-432; 1999.
- Richter, A., J. P. Burrows, H. Nüß, C. Granier, and U. Niemeier; *Increase in tropospheric nitrogen dioxide over China observed from space*; Nature; 437; 129-132; doi: 10.1038/nature04092, 2005.
- de Rooij, W. A., and C.C.A.H. van der Stap; *Expansion of Mie scattering matrices in generalized spherical functions*; Astronomy and Astrophysics; 131; 237-248; 1984.
- Rozanov, A., V. Rozanov, M. Buchwitz, A. Kokhanovsky, and J.P. Burrows; *SCIATRAN 2.0 - A new radiative transfer model for geophysical applications in the 175 - 2400 nm spectral region*; Adv.Space Res.; 36(5); 1015-1019; doi:10.1016/j.asr.2005.03.012; 2005.
- von Hoyningen-Huene, W., A. A. Kokhanovsky, J. P. Burrows, V. Bruniqnel-Pinel P. Regner, and F. Baret; *Simultaneous determination of aerosol- and surface characteristics from top-of-atmosphere reflectance using MERIS on board of ENVISAT*; Advances in Space Research; 37; 2172-2177; 2006.

Conclusions & Future work...

- From the sensitivity study performed it is clear that boundary layer height can have a large influence on the NO₂ AMF and such influence is intrinsically related to the NO₂ and aerosol profiles considered.
- Also, it was observed that changes of aerosol characteristics, such as load and type, are determinant variables in the calculations of AMF. Therefore, information on aerosol characteristics (e.g. if the aerosol is mainly composed by fine or coarse particles) is an important factor to obtain accurate AMF.
- It is concluded that the NO₂ AMF is dependent on many factors that need to be selected carefully. In order to improve the current retrieval method data from ECMWF (BL height) may be used to select and define the profiles and MERIS data (AOD measured) can be later incorporated in the retrieval method of tropospheric NO₂.