

**Seminar on Physics and Chemistry
of the Atmosphere
17.05.2024, SoSe 2024, IUP Bremen**

Improved tropospheric NO₂ columns retrieved from TROPOMI on board the Sentinel-5P satellite

**Thomas Visarius
(IUP)**

Abstract

NO₂ is a short-lived, highly reactive trace gas in the troposphere which is mainly co emitted with CO₂ during combustion processes. Therefore, NO₂ can be used as a proxy to estimate CO₂ emissions from large point sources such as coal fired power plants. Evaluating the potential of space-borne NO₂ observations for CO₂ emission estimates is one of the aims of the German ITMS (Integrated Tropospheric Monitoring System) project. To determine the vertical column density (VCD) of NO₂ the air mass factor (AMF) needs to be well known. It can be calculated using a priori information of the observation geometry and the state of the atmosphere. While the a priori data used in the operational TROPOMI retrieval is appropriate for global applications, it does not have sufficient spatial resolution for local retrievals for example over power plants.

In this work, spatially and temporally highly resolved AMF for the NO₂ retrieval window of the tropospheric monitoring instrument (TROPOMI) were calculated over Europe, using the bi-directional reflectance distribution function (BRDF) kernel coefficients obtained from the moderate imaging spectrometer (MODIS). The resulting AMFs are compared to AMFs obtained using the directional-dependent Lambertian equivalent reflectivity (DLER), which is used in the operational TROPOMI product.