



Product User Guide and Specification (PUGS) – ANNEX B for products CO2_GOS_SRFP & CH4_GOS_SRFP

C3S_312a_Lot6_IUP-UB – Greenhouse Gases

Issued by: R. G. Detmers, SRON, The Netherlands

Date: 20/10/2017

Ref: C3S_D312a_Lot6.3.1.5-v1_PUGS_ANNEX-B_v1.1

Official reference number service contract: 2016/C3S_312a_Lot6_IUP-UB/SC1



This document has been produced in the context of the Copernicus Climate Change Service (C3S). The activities leading to these results have been contracted by the European Centre for Medium-Range Weather Forecasts, operator of C3S on behalf of the European Union (Delegation Agreement signed on 11/11/2014). All information in this document is provided "as is" and no guarantee or warranty is given that the information is fit for any particular purpose. The user thereof uses the information at its sole risk and liability. For the avoidance of all doubts, the European Commission and the European Centre for Medium-Range Weather Forecasts has no liability in respect of this document, which is merely representing the authors view.



Contributors

**INSTITUTE OF ENVIRONMENTAL PHYSICS (IUP),
UNIVERSITY OF BREMEN, BREMEN, GERMANY
(IUP)**

M. Buchwitz

**SRON NETHERLANDS INSTITUTE FOR SPACE RESEARCH,
UTRECHT, THE NETHERLANDS
(SRON)**

I. Aben

R. G. Detmers

O. P. Hasekamp



Table of Contents

History of modifications	5
Related documents	6
Acronyms	7
General definitions	9
Scope of document	10
Executive summary	11
1. Data product description	12
2. Target requirements	14
3. Data usage information	14
3.1 Product Content and Format	14
3.2 Quality Flags and Metadata	17
3.3 Bias Correction	19
3.4 Recommended data usage	21
3.4 Tools for Reading the Data	21
3.5 Known Limitations and Issues	21
References	22



History of modifications

Version	Date	Description of modification	Chapters / Sections
1.0	21-August-2017	New document	All
1.0b	11-October-2017	Page header logo replaced	Page header
1.3	20-October-2017	Reference to main PUGS updated	Page 6



Related documents

Reference ID	Document
D1	Main PUGS: Buchwitz, M., et al., Product User Guide and Specification (PUGS) – Main document, C3S project C3S_312a_Lot6_IUP-UB – Greenhouse Gases, v1.3, 2017. <i>(this document is an ANNEX to the Main PUGS)</i>



Acronyms

Acronym	Definition
ATBD	Algorithm Theoretical Basis Document
CAR	Climate Assessment Report
C3S	Copernicus Climate Change Service
CCI	Climate Change Initiative
CDR	Climate Data Record
CDS	(Copernicus) Climate Data Store
CRG	Climate Research Group
D/B	Data base
EC	European Commission
ECMWF	European Centre for Medium Range Weather Forecasting
ECV	Essential Climate Variable
EO	Earth Observation
ESA	European Space Agency
EU	European Union
EUMETSAT	European Organisation for the Exploitation of Meteorological Satellites
FP	Full Physics retrieval method
FTIR	Fourier Transform InfraRed
FTS	Fourier Transform Spectrometer
GCOS	Global Climate Observing System
GEOSS	Global Earth Observation System of Systems
GHG	GreenHouse Gas
GOSAT	Greenhouse Gases Observing Satellite
IPCC	International Panel in Climate Change
IUP	Institute of Environmental Physics (IUP) of the University of Bremen, Germany
JAXA	Japan Aerospace Exploration Agency
KIT	Karlsruhe Institute of Technology
L1	Level 1
L2	Level 2
L3	Level 3
L4	Level 4
LMD	Laboratoire de Météorologie Dynamique
MACC	Monitoring Atmospheric Composition and Climate, EU GMES project
NA	Not applicable
NetCDF	Network Common Data Format
NIES	National Institute for Environmental Studies



NIR	Near Infra Red
NOAA	National Oceanic and Atmospheric Administration
Obs4MIPs	Observations for Climate Model Intercomparisons
ppb	Parts per billion
ppm	Parts per million
PR	(light path) PROxy retrieval method
PVIR	Product Validation and Intercomparison Report
QA	Quality Assurance
QC	Quality Control
REQ	Requirement
RMS	Root-Mean-Square
RTM	Radiative transfer model
SNR	Signal-to-Noise Ratio
SRON	SRON Netherlands Institute for Space Research
SWIR	Short Wave Infra Red
SZA	Solar Zenith Angle
TANSO	Thermal And Near infrared Sensor for carbon Observation
TANSO-FTS	Fourier Transform Spectrometer on GOSAT
TBC	To be confirmed
TBD	To be defined / to be determined
TCCON	Total Carbon Column Observing Network
TIR	Thermal Infra Red
TR	Target Requirements
TRD	Target Requirements Document
URD	User Requirements Document
WMO	World Meteorological Organization
Y2Y	Year-to-year (bias variability)



General definitions

Table 1 lists some general definitions relevant for this document.

Table 1: General definitions.

Item	Definition
XCO ₂	Column-averaged dry-air mixing ratios (mole fractions) of CO ₂
XCH ₄	Column-averaged dry-air mixing ratios (mole fractions) of CH ₄
L1	Level 1 satellite data product: geolocated radiance (spectra)
L2	Level 2 satellite-derived data product: Here: CO ₂ and CH ₄ information for each ground-pixel
L3	Level 3 satellite-derived data product: Here: Gridded CO ₂ and CH ₄ information, e.g., 5 deg times 5 deg, monthly
L4	Level 4 satellite-derived data product: Here: Surface fluxes (emission and/or uptake) of CO ₂ and CH ₄



Scope of document

This document is a Product User Guide and Specification (PUGS) for the Copernicus Climate Change Service (C3S, <https://climate.copernicus.eu/>) component as covered by project C3S_312a_Lot6 led by University of Bremen, Germany.

Within project C3S_312a_Lot6 satellite-derived atmospheric carbon dioxide (CO₂) and methane (CH₄) Essential Climate Variable (ECV) data products will be generated and delivered to ECMWF for inclusion into the Copernicus Climate Data Store (CDS) from which users can access these data products and the corresponding documentation.

The C3S_312a_Lot 6 satellite-derived data products are:

- Column-averaged dry-air mixing ratios (mole fractions) of CO₂ and CH₄, denoted XCO₂ (in parts per million, ppm) and XCH₄ (in parts per billion, ppb), respectively.
- Mid/upper tropospheric mixing ratios of CO₂ (in ppm) and CH₄ (in ppb).

This document describes the C3S products CO₂_GOS_SRFP and CH₄_GOS_SRFP.

These products are XCO₂ and XCH₄ Level 2 products as retrieved from GOSAT using algorithms developed at SRON, The Netherlands.



Executive summary

This document is the annex B of the Product User Guide (PUG), which is a deliverable of the C3S project. This document describes the RemoteC XCO₂ and XCH₄ Full Physics data products (CO₂_GOS_SRFP and CH₄_GOS_SRFP) so that it will be clear for the user how to use the product. The description includes quality flags and metadata, data format, product grid, known limitations, bias correction, and the product (column) averaging kernels and a description how to use them.



1. Data product description

The Japanese Greenhouse gases Observing SATellite (GOSAT) was launched on 23rd January 2009, *Yokota et al., 2009* by JAXA, the Japanese Space Agency. GOSAT provides the first dedicated global measurements of total column CO₂ and CH₄ from its SWIR bands, *Yoshida et al., 2010*. It is equipped with two instruments, the Thermal And Near Infrared Sensor for carbon Observations - Fourier Transform Spectrometer (TANSO-FTS) as well as a dedicated Cloud and Aerosol Imager (TANSO-CAI).

The TANSO-FTS instrument has four spectral bands with a high spectral resolution 0.3 cm⁻¹, three of which operate in the SWIR at around 0.76, 1.6 and 2.0 μm providing sensitivity to the near-surface absorbers with the fourth channel operating in the thermal infrared between 5.5 and 14.3 μm providing mid-tropospheric sensitivity.

The measurement strategy of TANSO-FTS is optimized for the characterization of continental-scale sources and sinks. TANSO-FTS utilizes a pointing mirror to perform off-nadir measurements at the same location on each 3-day repeat cycle. The pointing mirror allows TANSO-FTS to observe up to ±35° across track and ±20° along-track. These measurements nominally consist of 3 across track points spaced ~100km apart (although measurements are possible with 1, 3, 5, 7 or 9 across track points) with a ground footprint diameter of approximately 10.5 km and a 4 second exposure duration. Whilst the majority of data is limited to measurements over land where the surface reflectance is high, TANSO-FTS also observes in sunglint mode over the ocean within ±20° of the sub-solar latitude.

The CH₄_GOS_SRFP and CO₂_GOS_SRFP products are retrieved from GOSAT TANSO-FTS NIR and SWIR spectra using the RemoTeC algorithm that is being jointly developed at SRON and KIT. The algorithm retrieves simultaneously XCH₄ and XCO₂. For the retrieval, we analyse four spectral regions: the 0.77 μm oxygen band, two CO₂ bands at 1.61 and 2.06 μm, as well as a CH₄ band at 1.64 μm. Within the retrieval procedure the sub-columns of CO₂ and CH₄ in different altitude layers are being retrieved. To obtain the column averaged dry air mixing ratios XCO₂ and XCH₄ the sub-columns are summed up to get the total column which is divided by the dry-air columns obtained from ECMWF model data in combination with a surface elevation data base. The retrieved XCH₄ and XCO₂ have been extensively validated with ground based TCCON measurements. To further improve accuracy a bias correction has been developed based on TCCON comparisons. We use the GGG2014 TCCON dataset, *Wunch et al., 2015*.

More details on the technical aspects of the retrievals can be found in the ATBD Annex-B.



Figure 1: Global XCO₂ for the 2009-2016 period for the CO₂_GOS_SRFP product on a 1 by 1 degree resolution.





Figure 2: Global XCH₄ for the 2009-2016 period for the CH₄_GOS_SRFP product on a 1 by 1 degree resolution.





2. Target requirements

Table 1: Target requirements for XCH₄ and XCO₂

Random and systematic error requirements for XCO ₂ and XCH ₄					
Parameter	Req. type	Random error ("Precision")		Systematic error	Stability
		Single obs.	1000 ² km ² monthly		
XCO ₂	G	< 1 ppm	< 0.3 ppm	< 0.2 ppm (absolute)	As systematic error but per year
	B	< 3 ppm	< 1.0 ppm	< 0.3 ppm (relative ^s)	-"-
	T	< 8 ppm	< 1.3 ppm	< 0.5 ppm (relative [#])	-"-
XCH ₄	G	< 9 ppb	< 3 ppb	< 1 ppb (absolute)	< 1 ppb/year (absolute)
	B	< 17 ppb	< 5 ppb	< 5 ppb (relative ^s)	< 2 ppb/year (relative ^s)
	T	< 34 ppb	< 11 ppb	< 10 ppb (relative [#])	< 3 ppb/year (relative [#])

Table 1 shows the target requirements for XCH₄ and XCO₂ (Threshold, Breakthrough and Goal) as derived in the Target Requirements Document (TRD).

3. Data usage information

3.1 Product Content and Format

The CH₄_GOS_SRFP and CO₂_GOS_SRFP data products are stored per day in a single NetCDF file. Retrieval results are provided for the individual GOSAT spatial footprints, i.e. no averaging has been applied. The product file contains the key standard products, i.e. the retrieved column averaged dry air mixing ratios XCO₂ and XCH₄ with bias correction, averaging kernels and quality flags, as well as secondary products specific for the RemoTeC algorithm.



Table 2: Common variables for the CO2_GOS_SRFP product

Name	Type	Dim.	Units	Description
solar_zenith_angle	float	n	degrees	Angle between line of sight to the sun and local vertical
sensor_zenith_angle	float	n	degrees	Angle between the line of sight to the sensor and the local vertical
time	float	n	seconds	Seconds since 1970-01-01 00:00:00
longitude	float	n	degrees_ east	Center longitude
latitude	float	n	degrees_ north	Center latitude
pressure_levels	float	n, 13	hPa	Pressure levels
pressure_weight	float	n, 12		Layer dependent weights needed to apply the averaging kernels
xco2	float	n	1e-6	Retrieved column dry-air mole fraction of atmospheric carbon dioxide (XCO2) in ppm
xco2_uncertainty	float	n	1e-6	1-sigma uncertainty of the retrieved column-average dry-air mole fraction of atmospheric carbon dioxide
xco2_averaging_kernel	float	n, 12		Normalized column averaging kernel
co2_profile_apriori	float	n, 12	1e-6	A priori dry-air mole fraction profile of atmospheric carbon dioxide
xco2_quality_flag	int	n		Quality flag for XCO2 retrieval, 0 = good, 1 = bad

Table 2: Product specific (additional) variables for the CO2_GOS_SRFP product

Name	Type	Dim.	Units	Description
flag_landtype	int	n		0 = land, 1 = ocean
flag_sunlint	int	n		0 = no sunlint, 1 = sunlint
gain	char	n		gain setting of sensor, H = gain H, M = gain M
exposure_id	int	n		Exposure identification number of the sounding
l1b_name	char	n, 44		Name of the Level 1B file of the sounding
signal_to_noise_window	float	n, 4, 2		Signal to noise ratio per retrieval window and for both polarization directions
dry_airmass_layer	float	n, 12	m-2	Dry airmass per layer
altitude	float	n	m	Vertical distance above the surface
air_temperature	float	n, 13	K	The bulk temperature of the air at each level
surface_elevation_stdev	float	n	m	Standard deviation of the surface elevation within the sounding
x_wind	float	n, 13	m s-1	Eastward wind velocity
y_wind	float	n, 13	m s-1	Northward wind velocity
chi2	float	n		Chi-squared value of the sounding
optical_thickness_of_atmosphere_layer_due_to_ambient_aerosol	float	n, 4		Scattering optical thickness per retrieval window
raw_xco2	float	n	1e-6	Retrieved column dry-air mole fraction of atmospheric carbon dioxide (XCO2) in ppm before bias correction



raw_xco2_err	float	n	1e-6	1-sigma statistical uncertainty of the retrieved column-average dry-air mole fraction of atmospheric carbon dioxide
h2o_column	float	n	m-2	Retrieved total water column
surface_albedo_758	float	n		The retrieved albedo at 758 nm
surface_albedo_1593	float	n		The retrieved albedo at 1593 nm
surface_albedo_1629	float	n		The retrieved albedo at 1629 nm
surface_albedo_2042	float	n		The retrieved albedo at 2042 nm
intensity_offset_o2a	float	n	W cm-2	The retrieved intensity offset in the O2A band
aerosol_size	float	n		Retrieved size parameter of the aerosol distribution
aerosol_central_height	float	n	m	Peak height of the aerosol Gaussian height distribution
aerosol_total_column	float	n	m-2	Retrieved total aerosol column

Table 4: Common variables for the CH4_GOS_SRFP product

Name	Type	Dim.	Units	Description
solar_zenith_angle	float	n	degrees	Angle between line of sight to the sun and local vertical
sensor_zenith_angle	float	n	degrees	Angle between the line of sight to the sensor and the local vertical
time	float	n	seconds	Seconds since 1970-01-01 00:00:00
longitude	float	n	degrees_east	Center longitude
latitude	float	n	degrees_north	Center latitude
pressure_levels	float	n, 13	hPa	Pressure levels
pressure_weight	float	n, 12		Layer dependent weights needed to apply the averaging kernels
xch4	float	n	1e-9	Retrieved column dry-air mole fraction of atmospheric methane (XCH4) in ppb
xch4_uncertainty	float	n	1e-9	1-sigma uncertainty of the retrieved column-average dry-air mole fraction of atmospheric methane
xch4_averaging_kernel	float	n, 12		Normalized column averaging kernel
ch4_profile_apriori	float	n, 12	1e-9	A priori dry-air mole fraction profile of atmospheric methane
xch4_quality_flag	int	n		Quality flag for XCH4 retrieval, 0 = good, 1 = bad

Table 5: Product specific (additional) variables for the CH4_GOS_SRFP product

Name	Type	Dim.	Units	Description
flag_landtype	int	n		0 = land, 1 = ocean
flag_sunlint	int	n		0 = no sunlint, 1 = sunlint
gain	char	n		gain setting of sensor, H = gain H, M = gain M
exposure_id	int	n		Exposure identification number of the sounding
l1b_name	char	n, 44		Name of the Level 1B file of the sounding



signal_to_noise_window	float	n, 4, 2		Signal to noise ratio per retrieval window and for both polarization directions
dry_airmass_layer	float	n, 12	m-2	Dry airmass per layer
altitude	float	n	m	Vertical altitude above the surface
air_temperature	float	n, 13	K	The bulk temperature of the air at each level
surface_altitude_stdev	float	n	m	Standard deviation of the surface elevation within the sounding
x_wind	float	n, 13	m s-1	Eastward wind velocity
y_wind	float	n, 13	m s-1	Northward wind velocity
chi2	float	n		Chi-squared value of the sounding
optical_thickness_of_atmosphere_layer_due_to_ambient_aerosol	float	n, 4		Scattering optical thickness per retrieval window
raw_xch4	float	n	1e-9	Retrieved column dry-air mole fraction of atmospheric methane (XCH4) in ppb before bias correction
raw_xch4_err	float	n	1e-9	1-sigma statistical uncertainty of the retrieved column-average dry-air mole fraction of atmospheric methane
h2o_column	float	n	m-2	Retrieved total water column
surface_albedo_758	float	n		The retrieved albedo at 758 nm
surface_albedo_1593	float	n		The retrieved albedo at 1593 nm
surface_albedo_1629	float	n		The retrieved albedo at 1629 nm
surface_albedo_2042	float	n		The retrieved albedo at 2042 nm
intensity_offset_o2a	float	n	W cm-2	The retrieved intensity offset in the O2A band
aerosol_size	float	n		Retrieved size parameter of the aerosol distribution
aerosol_central_height	float	n	m	Peak height of the aerosol Gaussian height distribution
aerosol_total_column	float	n	m-2	Retrieved total aerosol column

3.2 Quality Flags and Metadata

There are quality flags "xco2_quality_flag" and "xch4_quality_flag" included in the data files. The quality flag can have 2 values:

- 0: retrievals for **H-gain**, **M-gain** or **sunlint** data, quality has been checked
- 1: data should not be used (e.g. bad fit to data, residual cloud contamination)

For a GOSAT ground pixel to be processed by the RemoTeC Full Physics algorithm it has to fulfill the following criteria: GOSAT nominal quality flags should be good and the standard deviation of the elevation in the pixel should be less than 1000 meters (to filter out the most extreme terrains). After the retrieval step the data that fulfill the following criteria are flagged as '0' for **gain H**:

- Error on retrieved XCO₂ < 1.2 ppm
- χ^2 of fit < 4.5
- χ^2 of fit in O2A-band < 4
- SNR > 50
- Standard deviation of surface elevation within GOSAT ground pixel should be < 80 m



- $0 < \text{aerosol_filter} < 300$
- Aerosol Optical Thickness < 0.3
- $3 < \text{Aero_size} < 5$
- $\text{SZA} < 70^\circ$
- $-1\text{e-}9 < \text{Intensity Offset O2A-band} < 3\text{e-}9$
- Blended Albedo < 0.9
- $0.99 < \text{CO2 (1.6 micron)} / \text{CO2 (2.0 micron)} < 1.015$
- $0.95 < \text{O2 (retrieved)} / \text{O2 (prior)} < 1.02$
- $0.95 < \text{H2O (1.6 micron)} / \text{H2O (2.0 micron)} < 1.08$

For **gain M**:

- Number of iteration steps in retrieval < 25
- Error on retrieved XCO₂ < 1.2 ppm
- χ^2 of fit < 5
- χ^2 of fit in O2A-band < 4
- χ^2 of fit in window 2 < 4
- SNR > 50
- Standard deviation of surface elevation within GOSAT ground pixel should be < 80 m
- $0 < \text{aerosol_filter} < 350$
- Aerosol Optical Thickness < 0.25
- $3 < \text{Aero_size} < 4$
- $\text{SZA} < 70^\circ$
- $0.99 < \text{CO2 (1.6 micron)} / \text{CO2 (2.0 micron)} < 1.015$
- $0.97 < \text{O2 (retrieved)} / \text{O2 (prior)} < 1.01$
- $0.95 < \text{H2O (1.6 micron)} / \text{H2O (2.0 micron)} < 1.05$

For **sunglint**:

- Number of iteration steps in retrieval < 31
- Cirrus Signal $< 8\text{e-}10$



- χ^2 of fit < 4
- χ^2 of fit in O2A-band < 4
- χ^2 of fit in window 4 < 10
- $-1e-5 < \text{Retrieved slope of albedo in window 4} < 5e-5$
- $-1e-9 < \text{Intensity Offset O2A-band} < 3e-9$
- $0.99 < \text{CO2 (1.6 micron)} / \text{CO2 (2.0 micron)} < 1.01$
- $0.965 < \text{O2 (retrieved)} / \text{O2 (prior)} < 1.00$
- $0.95 < \text{H2O (1.6 micron)} / \text{H2O (2.0 micron)} < 1.05$

3.3 Bias Correction

From comparison with TCCON it was found that the error in XCO₂ correlates with the solar zenith angle (sza) and the aerosol filter ϕ . Based on this correlation the following bias correction has been developed for XCO₂ for **gain H**:

$$XCO2_{corr} = XCO2 * (a + b * \phi + c * sza)$$

with $a = 0.999995$, $b = 2.8204e-5$, $c = 7.287e-5$.

For **gain M** we apply a correction based on just the aerosol filter ϕ :

$$XCO2_{corr} = XCO2 * (a + b * \phi)$$

and with $a = 1.004228$, $b = -3.00868e-6$.

For **sunglint** data we use a correction factor based on the O₂ ratio:

$$XCO2_{corr} = XCO2 * (a + b * RO2)$$

with $a = 1.283633$ and $b = -0.28368$.

The bias correction parameters are obtained from fits to the GOSAT-TCCON differences.

Similarly, for XCH₄ it was found that the error in XCH₄ correlates with the solar zenith angle (sza) and the aerosol filter ϕ . Based on this correlation the following bias correction has been developed for XCH₄ for **gain H**:

$$XCH4_{corr} = XCH4 * (a + b * \phi + c * sza)$$

with $a = 0.996226$, $b = 4.4482e-5$, $c = 6.0089e-5$.

For **gain M** we apply a correction based on just the aerosol filter ϕ :

$$XCH4_{corr} = XCH4 * (a + b * \phi)$$



with $a = 1.002728$, $b = 1.0053e-5$.

For **sunglint** data we use a correction factor based on the O2 ratio:

$$XCH4_{corr} = XCH4 * (a + b * RO2)$$

with $a = 1.18122$ and $b = -0.18569$.

The bias correction parameters are obtained from fits to the GOSAT-TCCON differences.

Figure 3: Co-located GOSAT-TCCON XCH4 measurements for gain H measurements





Figure 4: Co-located GOSAT-TCCON XCO2 measurements for gain H measurements





3.4 Recommended data usage

It is strongly recommended to only use the bias-corrected data in: "xco2" and "xch4" except if users explicitly correct for biases themselves (e.g. in an inverse modeling framework). Here, it should be noted that the bias correction has been developed independently for the different GOSAT-FTS instrument settings (**H-gain**, **M-gain**, **sunglint**).

Also, use only data over land (land type=0) except for sunglint cases.

If the data are to be compared with other XCO₂ and/or XCH₄ data for which vertical profile information is available (e.g. inverse modeling, comparison to models, comparison to measured profiles), the column averaging kernels should be used. Here it should be noted that **the column averaging kernels are to be applied to layer sub-columns (m-2)**, as these are the quantities directly retrieved in the RemoTeC algorithm.

For model comparisons the retrieved XCO₂ should be compared to $[VCO_2]'_{model}/[VAIR]_{model}$ where $[VAIR]_{model}$ is the total dry air column provided by the model and $[VCO_2]'_{model}$ is the model total CO₂ column after applying the column averaging kernel, viz.:

$$[VCO_2]'_{model} = [VCO_2]_{prior} + \mathbf{a}^T (\mathbf{x}_{model} - \mathbf{x}_{prior})$$

where $[VCO_2]_{prior}$ is the prior CO total column used in the retrieval, \mathbf{x}_{model} is the vertical CO₂ profile from the model (as sub-columns) and \mathbf{x}_{prior} is the prior vertical profile from the retrieval. For application of the column averaging kernel the model vertical profile should be re-calculated on the vertical grid of the retrieval (preferred) or the averaging kernel has to be interpolated to the vertical grid of the model. This procedure holds in the same way for the SRFP XCH₄ product, but then replacing all instances of CO₂ with CH₄.

3.4 Tools for Reading the Data

The data are stored in Netcdf format which can be read with standard tools in the common programming languages (IDL, Matlab, Python, Fortran90, C++, etc).

3.5 Known Limitations and Issues

- The data retrieved for the **H-gain** instrument settings are considered highest quality and are well validated. In the "raw" retrievals (i.e. before bias correction) there is a bias between **H-gain** and **sunglint** and **M-gain** retrievals, respectively. Although these biases have been corrected in the bias-corrected products, there may still be a small residual bias left, especially due to the limited number of validation sites for **sunglint** and **M-gain** retrievals.
- The 2nd half of December 2014 and the whole of January 2015 have no data due to the GOSAT satellite being in calibration mode after switching to the backup pointing system.



References

- Buchwitz et al., 2015:** Buchwitz, M., Reuter, M., Schneising, O., Boesch, H., Guerlet, S., Dils, B., Aben, I., Armante, R., Bergamaschi, P., Blumenstock, T., Bovensmann, H., Brunner, D., Buchmann, B., Burrows, J.P., Butz, A., Chédin, A., Chevallier, F., Crevoisier, C.D., Deutscher, N.M., Frankenberg, C., Hase, F., Hasekamp, O.P., Heymann, J., Kaminski, T., Laeng, A., Lichtenberg, G., De Mazière, M., Noël, S., Notholt, J., Orphal, J., Popp, C., Parker, R., Scholze, M., Sussmann, R., Stiller, G.P., Warneke, T., Zehner, C., Bril, A., Crisp, D., Griffith, D.W.T., Kuze, A., O'Dell, C., Oshchepkov, S., Sherlock, V., Suto, H., Wennberg, P., Wunch, D., Yokota, T., Yoshida, Y., The Greenhouse Gas Climate Change Initiative (GHG-CCI): comparison and quality assessment of near-surface-sensitive satellite-derived CO₂ and CH₄ global data sets. *Remote Sens. Environ.* 162:344–362, <http://dx.doi.org/10.1016/j.rse.2013.04.024>, 2015.
- Buchwitz et al., 2016:** Buchwitz, M., Reuter, M., Schneising, O., Hewson, W., Detmers, R. G., Boesch, H., Hasekamp, O. P., Aben, I., Bovensmann, H., Burrows, J. P., Butz, A., Chevallier, F., Dils, B., Frankenberg, C., Heymann, J., Lichtenberg, G., De Mazière, M., Notholt, J., Parker, R., Warneke, T., Zehner, C., Griffith, D. W. T., Deutscher, N. M., Kuze, A., Suto, H., and Wunch, D.: Global satellite observations of column-averaged carbon dioxide and methane: The GHG-CCI XCO₂ and XCH₄ CRDP3 data, *Remote Sensing of Environment (in press)*, Special Issue on Essential Climate Variables, DOI: 10.1016/j.rse.2016.12.027, (link: <http://dx.doi.org/10.1016/j.rse.2016.12.027>), 2016.
- Buchwitz et al., 2017:** ESA Climate Change Initiative (CCI) Product Validation and Intercomparison Report (PVIR) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG) for data set Climate Research Data Package No. 4 (CRDP#4), Version 5.0, 9. Feb. 2017, link: http://www.esa-ghg-cci.org/?q=webfm_send/352, 2017.
- Buchwitz et al., 2017a:** Buchwitz, M.; Reuter, M.; Aben, I.; Boesch, H.; Butz, A.; Detmers, R.G.; Frankenberg, C.; Hasekamp, O.P.; Parker, R.; Schneising, O.; Somkuti, P., ESA Greenhouse Gases Climate Change Initiative (GHG-CCI): Merged SCIAMACHY and GOSAT Level 3 gridded atmospheric column-average methane (XCH₄) product in Obs4MIPs format version 2 (CRDP#4), Technical Note, link: http://www.esa-ghg-cci.org/?q=webfm_send/349, pp. 11, 1 February 2017, 2017.
- Butz et al., 2011:** Butz, A., Guerlet, S., Hasekamp, O., et al., Toward accurate CO₂ and CH₄ observations from GOSAT, *Geophys. Res. Lett.*, doi:10.1029/2011GL047888, 2011.
- Butz et al., 2012:** Butz, A., Galli, A., Hasekamp, O., Landgraf, J., Tol, P., and Aben, I.: Remote Sensing of Environment, TROPOMI aboard Sentinel-5 Precursor : Prospective performance of CH₄ retrievals for aerosol and cirrus loaded atmospheres, 120, 267-276, doi:10.1016/j.rse.2011.05.030, 2012.
- Detmers et al., 2015:** Detmers, R. G., O. Hasekamp, I. Aben, S. Houweling, T. T. van Leeuwen, A. Butz, J. Landgraf, P. Koehler, L. Guanter, and B. Poulter, [Anomalous carbon uptake in Australia as seen by GOSAT](#), *Geophys. Res. Lett.*, 42, doi:10.1002/2015GL065161, 2015.
- ESA-CCI-GHG-URDv2.1:** Chevallier, F., et al., User Requirements Document (URD), ESA Climate Change Initiative (CCI) GHG-CCI project, Version 2.1, 19 Oct 2016, link: http://www.esa-ghg-cci.org/?q=webfm_send/344, 2016.



- Kuze et al., 2009:** Kuze, A., Suto, H., Nakajima, M., and Hamazaki, T. (2009), Thermal and near infrared sensor for carbon observation Fourier-transform spectrometer on the Greenhouse Gases Observing Satellite for greenhouse gases monitoring, *Appl. Opt.*, 48, 6716–6733, 2009.
- Kuze et al., 2016:** Kuze, A., Suto, H., Shiomi, K., Kawakami, S., Tanaka, M., Ueda, Y., Deguchi, A., Yoshida, J., Yamamoto, Y., Kataoka, F., Taylor, T. E., and Buijs, H. L.: Update on GOSAT TANSO-FTS performance, operations, and data products after more than 6 years in space, *Atmos. Meas. Tech.*, 9, 2445-2461, doi:10.5194/amt-9-2445-2016, 2016.
- Schepers et al., 2012:** Schepers, D., Guerlet, S., Butz, A., Landgraf, J., Frankenberg, C., Hasekamp, O., Blavier, J.-F., Deutscher, N. M., Griffith, D. W. T., Hase, F., Kyro, E., Morino, I., Sherlock, V., Sussmann, R., Aben, I. (2012), Methane retrievals from Greenhouse Gases Observing Satellite (GOSAT) shortwave infrared measurements: Performance comparison of proxy and physics retrieval algorithms, *J. Geophys. Res.*, 117, D10307, doi:10.1029/2012JD017549, 2012.
- TRD GHG, 2017:** Buchwitz, M., Aben, I., Anand, J., Armante, R., Boesch, H., Crevoisier, C., Detmers, R. G., Hasekamp, O. P., Reuter, M., Schneising-Weigel, O., Target Requirement Document, Copernicus Climate Change Service (C3S) project on satellite-derived Essential Climate Variable (ECV) Greenhouse Gases (CO₂ and CH₄) data products (project C3S_312a_Lot6), Version 1, 28-March-2017, pp. 52, 2017.
- Wunch et al. 2015:** Wunch, D., Toon, G.C., Sherlock, V., Deutscher, N.M., Liu, X., Feist, D.G., Wennberg, P.O., The Total Carbon Column Observing Network's GGG2014 Data Version. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA (available at: doi:10.14291/tccon.ggg2014.documentation.R0/1221662), 2015.
- Yokota et al., 2009:** Yokota, T., Y. Yoshida, N. Eguchi, Y. Ota, T. Tanaka, H. Watanabe, and S. Maksyutov (2009), Global concentrations of CO₂ and CH₄ retrieved from GOSAT: First preliminary results, *SOLA*, 5, 160-163.
- Yoshida et al., 2010:** Yoshida, Y., Y. Ota, N. Eguchi, N. Kikuchi, K. Nobuta, H. Tran, I. Morino, and T. Yokota (2010), Retrieval algorithm for CO₂ and CH₄ column abundances from short-wavelength infrared spectral observations by the greenhouse gases observing satellite, *Atmospheric Measurement Techniques Discussions*, 3(6), 4791-4833, doi:10.5194/amtd-3-4791-2010.

