



Copernicus Climate Change Service



Product Quality Assessment Report (PQAR) – ANNEX D for products XCO2_EMMA and XCH4_EMMA (v3.1, 2003-2017)

C3S_312a_Lot6_IUP-UB – Greenhouse Gases

Issued by: Maximilian Reuter, University of Bremen,
Institute of Environmental Physics (IUP)

Date: 04/10/2018

Ref: C3S_D312a_Lot6.3.1.7-v2_PQAR_ANNEX-D_v2.0

Official reference number service contract: 2016/C3S_312a_Lot6_IUP-UB/SC1



This document has been produced in the context of the Copernicus Climate Change Service (C3S).

The activities leading to these results have been contracted by the European Centre for Medium-Range Weather Forecasts, operator of C3S on behalf of the European Union (Delegation Agreement signed on 11/11/2014). All information in this document is provided "as is" and no guarantee or warranty is given that the information is fit for any particular purpose.

The user thereof uses the information at its sole risk and liability. For the avoidance of all doubts, the European Commission and the European Centre for Medium-Range Weather Forecasts has no liability in respect of this document, which is merely representing the authors view.



Contributors

**INSTITUTE OF ENVIRONMENTAL PHYSICS (IUP),
UNIVERSITY OF BREMEN, BREMEN, GERMANY
(IUP)**

M. Reuter

M. Buchwitz

O. Schneising-Weigel



Table of Contents

History of modifications	5
Related documents	6
Acronyms	7
General definitions	11
Scope of document	12
Executive summary	13
1. Product validation methodology	14
2. Validation Results	16
2.1 XCO₂_EMMA	16
2.1.1 Validation	16
2.1.2 Summary	24
2.2 XCH₄_EMMA	25
2.2.1 Validation	25
2.2.2 Summary	32
References	33



History of modifications

Version	Date	Description of modification	Chapters / Sections
1.1	20-October-2017	New document for data set CDR1 (2003-2016)	All
2.0	4-October-2018	Update for CDR2 (2003-2017)	All



Related documents

Reference ID	Document
D1	Main PQAR: Buchwitz, M., et al., Product Quality Assessment Report (PQAR) – Main document, C3S project C3S_312a_Lot6_IUP-UB – Greenhouse Gases, v2.0, 2018. <i>(this document is an ANNEX to the Main PQAR)</i>
D2	ATBD ANNEX D: Reuter, M., et al., Algorithm Theoretical Basis Document (ATBD) – ANNEX D for products XCO2_EMMA (v3.1, 2003-2017), XCH4_EMMA, C3S project C3S_312a_Lot6_IUP-UB – Greenhouse Gases, 2018.



Acronyms

Acronym	Definition
AIRS	Atmospheric Infrared Sounder
AMSU	Advanced Microwave Sounding Unit
ATBD	Algorithm Theoretical Basis Document
BESD	Bremen optimal ESTimation DOAS
CAR	Climate Assessment Report
C3S	Copernicus Climate Change Service
CCDAS	Carbon Cycle Data Assimilation System
CCI	Climate Change Initiative
CDR	Climate Data Record
CDS	(Copernicus) Climate Data Store
CMUG	Climate Modelling User Group (of ESA's CCI)
CRG	Climate Research Group
D/B	Data base
DOAS	Differential Optical Absorption Spectroscopy
EC	European Commission
ECMWF	European Centre for Medium Range Weather Forecasting
ECV	Essential Climate Variable
EMMA	Ensemble Median Algorithm
ENVISAT	Environmental Satellite (of ESA)
EO	Earth Observation
ESA	European Space Agency
EU	European Union



EUMETSAT	European Organisation for the Exploitation of Meteorological Satellites
FCDR	Fundamental Climate Data Record
FoM	Figure of Merit
FP	Full Physics retrieval method
FTIR	Fourier Transform InfraRed
FTS	Fourier Transform Spectrometer
GCOS	Global Climate Observing System
GEO	Group on Earth Observation
GEOSS	Global Earth Observation System of Systems
GHG	GreenHouse Gas
GOME	Global Ozone Monitoring Experiment
GMES	Global Monitoring for Environment and Security
GOSAT	Greenhouse Gases Observing Satellite
IASI	Infrared Atmospheric Sounding Interferometer
IMAP-DOAS (or IMAP)	Iterative Maximum A posteriori DOAS
IPCC	International Panel in Climate Change
IUP	Institute of Environmental Physics (IUP) of the University of Bremen, Germany
JAXA	Japan Aerospace Exploration Agency
JCGM	Joint Committee for Guides in Metrology
L1	Level 1
L2	Level 2
L3	Level 3
L4	Level 4
LMD	Laboratoire de Météorologie Dynamique
MACC	Monitoring Atmospheric Composition and Climate, EU GMES project
NA	Not applicable
NASA	National Aeronautics and Space Administration



NetCDF	Network Common Data Format
NDACC	Network for the Detection of Atmospheric Composition Change
NIES	National Institute for Environmental Studies
NIR	Near Infra Red
NLIS	LMD/CNRS <i>neuronal</i> network mid/upper tropospheric CO ₂ and CH ₄ retrieval algorithm
NOAA	National Oceanic and Atmospheric Administration
Obs4MIPs	Observations for Climate Model Intercomparisons
OCO	Orbiting Carbon Observatory
OE	Optimal Estimation
PBL	Planetary Boundary Layer
ppb	Parts per billion
ppm	Parts per million
PR	(light path) PROxy retrieval method
PVIR	Product Validation and Intercomparison Report
QA	Quality Assurance
QC	Quality Control
REQ	Requirement
RMS	Root-Mean-Square
RTM	Radiative transfer model
SCIAMACHY	SCanning Imaging Absorption spectroMeter for Atmospheric ChartographY
SCIATRAN	SCIAMACHY radiative transfer model
SRON	SRON Netherlands Institute for Space Research
SWIR	Short Wava Infra Red
TANSO	Thermal And Near infrared Sensor for carbon Observation
TANSO-FTS	Fourier Transform Spectrometer on GOSAT
TBC	To be confirmed
TBD	To be defined / to be determined



TCCON	Total Carbon Column Observing Network
TIR	Thermal Infra Red
TR	Target Requirements
TRD	Target Requirements Document
WFM-DOAS (or WFMD)	Weighting Function Modified DOAS
UoL	University of Leicester, United Kingdom
URD	User Requirements Document
WMO	World Meteorological Organization
Y2Y	Year-to-year (bias variability)



General definitions

Table 1 lists some general definitions relevant for this document.

Table 1: General definitions.

Item	Definition
XCO ₂	Column-averaged dry-air mixing ratios (mole fractions) of CO ₂
XCH ₄	Column-averaged dry-air mixing ratios (mole fractions) of CH ₄
L1	Level 1 satellite data product: geolocated radiance (spectra)
L2	Level 2 satellite-derived data product: Here: XCO ₂ and XCH ₄ information for each ground-pixel
L3	Level 3 satellite-derived data product: Here: Gridded XCO ₂ and XCH ₄ information, e.g., 5°x5°, monthly
L4	Level 4 satellite-derived data product: Here: Surface fluxes (emission and/or uptake) of CO ₂ and CH ₄



Scope of document

This document is a Product Quality Assessment Report (PQAR) for the Copernicus Climate Change Service (C3S, <https://climate.copernicus.eu/>) component as covered by project C3S_312a_Lot6 led by University of Bremen, Germany.

Within project C3S_312a_Lot6 satellite-derived atmospheric carbon dioxide (CO₂) and methane (CH₄) Essential Climate Variable (ECV) data products will be generated and delivered to ECMWF for inclusion into the Copernicus Climate Data Store (CDS) from which users can access these data products and the corresponding documentation.

The C3S_312a_Lot 6 satellite-derived data products are:

- Column-averaged dry-air mixing ratios (mole fractions) of CO₂ and CH₄, denoted XCO₂ (in parts per million, ppm) and XCH₄ (in parts per billion, ppb), respectively.
- Mid/upper tropospheric mixing ratios of CO₂ (in ppm) and CH₄ (in ppb).

This document describes the validation / quality assessment of the C3S products XCO₂_EMMA and XCH₄_EMMA.

These products are merged multi-sensor XCO₂ and XCH₄ Level 2 products generated using algorithms developed at University of Bremen, Germany.



Executive summary

This Product Quality Assessment Report (PQAR) describes the validation of the EMMA v3.1 CO₂ and EMMA v3.1 CH₄ products (in the following also referred to as XCO₂_EMMA and XCH₄_EMMA) with TCCON ground based measurements. Originally, the EMMA algorithm (v1.3) was described and validated in the publication of *Reuter et al. (2013)* and their validation is the blueprint for this PQAR. Several of the shown figures are updated versions of figures shown in the publication of *Reuter et al. (2013)*. EMMA is composed of an ensemble of individual SCIAMACHY and GOSAT L2 algorithms and this document also contributes to the inter-comparison of the contributing algorithms.

For XCO₂ we find that the individual algorithms have a single measurement precision in the range of 1.74ppm (ACOS) to 2.30ppm (NIES v02). EMMA has a single measurement precision of 1.92ppm. EMMA's combined regional and seasonal biases (0.58ppm) are on the lower end of the range of the individual algorithms (0.43ppm for BESD to 1.00ppm for PPDF-S). The found linear drifts are small and predominantly not significant, i.e., the trend is usually smaller than twice its uncertainty. The linear drift found for EMMA's XCO₂ is 0.00 ± 0.09 ppm/a. The year-to-year stability is in the range of 0.27ppm/a (ACOS) and 0.50ppm/a (NIES v02bc). EMMA's year-to-year stability of 0.29ppm/a is on the lower end of this range.

For XCH₄ we find that the individual algorithms have a single measurement precision in the range of 13.4ppb (UoL-FP) to 14.4ppb (PPDF-S) except for WFMD which has a single measurement precision of 100.5ppb. EMMA has a single measurement precision of 13.7ppb (excluding the period till 04/2010). EMMA's combined regional and seasonal biases (4.3ppb) are at the lower end of the range of the individual GOSAT algorithms (4.3ppb for UoL-FP to 14.6ppb for WFMD). The found linear drifts are small and not significant, i.e., the trend is smaller than twice its uncertainty. The linear drift found for EMMA's XCH₄ is -0.05 ± 0.67 ppb/a. The year-to-year stability is in the range of 1.7ppb/a (UoL-PR, NIES v02) and 7.0ppb/a (WFMD). EMMA's year-to-year stability of 1.8ppb/a is at the lower end of the range.



1. Product validation methodology

As described in D2, EMMA v3.1 CO₂ and CH₄ make use of the following satellite data products: BESD v02.01.02 (*Reuter et al., 2016*), RemoTeC v2.3.8 (*Detmers et al., 2017a*), ACOS v7.3.10a (*O'Dell et al., 2012*), UoL-FP v7.2 (*Boesch and Anand, 2017*), and NIES v02 (*Yoshida et al., 2013*) without and with (v02bc) bias correction, and PPDF-S v02 (*Bril et al., 2012*) for XCO₂ and WFMD v4.0 (*Schneising et al., 2016*), RemoTeC-FP v2.3.8 (*Detmers et al., 2017a*), RemoTeC-PR v2.3.9 (*Detmers et al., 2017b*), UoL-FP v7.2 (*Boesch and Anand, 2017*), UoL-PR v7.2 (*Boesch and Anand, 2017*), and NIES v02 (*Yoshida et al., 2011*) without and with (v02bc) bias correction, and PPDF-S v02 (*Bril et al., 2012*) for XCH₄.

The EMMA CO₂ and CH₄ data products are validated with TCCON (using version GGG2014 R0/1) measurements in a similar way as done by *Reuter et al. 2011*. The co-location criteria are defined by a maximal time difference of two hours, a maximal spatial distance of 500km, and a maximal surface elevation difference of 250m.

For each TCCON site with more than 250 co-locations and covering a time period of at least one year, the performance statistics number of co-locations, station bias, seasonal bias, linear drift, and single measurement precision are calculated. The validation period ranges from 01/2003 to 12/2017.

The station bias is defined as average difference to TCCON and the single measurement precision as standard deviation of the difference to TCCON. Seasonal bias and linear drift are derived by fitting the following trend model:

$$\Delta X = a_0 + a_1 t + a_2 \sin(2\pi t + a_3) + \varepsilon$$

Here, ΔX represents the difference satellite minus TCCON, ε the residual, and a_{0-3} the free fit parameters. Specifically, a_1 represents the linear drift and a_2 the amplitude of the seasonal bias.

Based on the per station statistics, the following summarizing statistics are calculated: Total number of co-locations used for validation, (quadratic) average single measurement precision, station-to-station bias (standard deviation of the station biases), average seasonal bias (standard deviation of the seasonal bias term), and average linear drift. As the linear drift can be assumed to be globally constant, the station-to-station standard deviation of the linear drift is a measure for its uncertainty.

Additionally, a measure for the year-to-year stability is computed: For each TCCON site, the residual difference (satellite - TCCON) which is not explained by station bias, seasonal bias, and/or linear drift is derived by subtracting the fit of the trend model ΔX from the satellite minus TCCON difference. These time series are smoothed by a running average of 365 days. Only days with more than 10 co-locations contributing to the running average of at least 5 TCCON sites are further considered. At these days, the station-to-station average is calculated. The corresponding expected uncertainty is computed from the standard error of the mean (derived from the station-to-station standard deviation and the number of stations) and by error propagation of the reported single sounding uncertainties.



Due to the relatively large uncertainty, we do not compute the maximum minus minimum as a measure for the year-to-year stability because this quantity can be expected to increase with length of the time series simply due to statistics. Therefore, we estimate the year-to-year stability by randomly selecting pairs of dates with a time difference of at least 365 days. For each selection we compute the difference modified by a random component corresponding to the estimated uncertainty. From 1000 of such pairs we compute the standard deviation as estimate for the year-to-year stability. We repeat this experiment 1000 times and compute the average and standard deviation.

As EMMA is constructed from an ensemble of individual L2 algorithms, the purpose of this document is not only product validation but also inter-comparison. Therefore, all individual algorithms contributing to EMMA are validated in the exact same way as EMMA.

Additionally, this document shows assessments of temporal and spatial bias patterns based on $10^{\circ} \times 10^{\circ}$ monthly gridded level 3 data sets.

We calculate the fraction of potential outliers according to unrealistically large spatial gradients ($>3\text{ppm}/10^{\circ}$ for XCO₂ and $>20\text{ppb}/10^{\circ}$ for XCH₄), unrealistically large deviations from CarbonTracker CT2017 ($>3\text{ppm}$ for XCO₂ and $>20\text{ppb}/10^{\circ}$ for XCH₄), and larger deviations from EMMA ($>2.5\text{ppm}$ for XCO₂ and $>10\text{ppb}/10^{\circ}$ for XCH₄).

We also compare the north/south (N/S) gradient of each month with CT2017 and TCCON by averaging all northern and southern hemispheric grid boxes (using the same sampling). However, it shall be noted that the statistics in comparing to TCCON are less robust because only few grid boxes include TCCON stations.

Additionally, we compare the seasonal (peak-to-peak) amplitude of each grid box with CT2017 and TCCON by calculating the difference between annual maximum and minimum. We consider only those grid boxes with at least six valid months and use the same sampling. Again, the TCCON statistics are probably not very robust because they rely on few grid boxes with seasonal cycles.



2. Validation Results

2.1 XCO₂_EMMA

2.1.1 Validation

Figure 1 shows all co-located EMMA and TCCON retrievals used for validation. Additionally, it includes all co-locations of the individual algorithms contributing to EMMA. The overall statistics per contributing algorithm are summarized in **Table 2**. **Table 3** shows the validation summary specifically for EMMA v3.1 CO₂, i.e., the XCO₂_EMMA product. The results are valid for the time periods covered by the individual algorithms but starting in 2009, respectively.

The individual algorithms have a single measurement precision in the range of 1.74ppm (ACOS) – 2.30ppm (NIES v02). EMMA has a single measurement precision of 1.92ppm. EMMA's combined regional and seasonal biases (0.58ppm) are on the lower end of the range of the individual algorithms (0.43ppm for BESD to 1.00ppm for PPDF-S).

Figure 2 (left) shows the anomaly of station biases of the used algorithms. One can see that most satellite retrievals have a high bias of about 0.3ppm – 1.0ppm at the Sodankylä, Garmisch-Partenkirchen, Bremen, and Karlsruhe TCCON sites and low bias of similar magnitude at Lamont and the southern hemispheric sites Darwin and Wollongong. This feature considerably contributes to the algorithms station-to-station bias statistics. Currently, it is unclear whether this discrepancy comes from the satellite retrievals or the TCCON.

The drift analysis in **Figure 2** (right) shows more or less small negative trends (typically below -0.1ppm/a) for most algorithms at the sites Lamont and Wollongong and positive trends of the same magnitude for most algorithms at Orleans and Bremen. This is a bit surprising because some of the sites are located in similar latitude bands.

Figure 3 shows the smoothed average residual difference (satellite - TCCON) which is not explained by station bias, seasonal bias, and/or linear drift. The year-to-year stability computed from the variability of the average is in the range of 0.27ppm/a (ACOS) and 0.50ppm/a (NIES v02bc). EMMA's year-to-year stability of 0.29ppm/a lies at the lower end of this range.

Analyses of gridded L3 data show that all algorithms reproduce large scale features well, however, there are still differences of a few ppm when looking into the details (**Figure 4**).

The satellite retrieved seasonal amplitudes are in similarly good agreement with TCCON and CarbonTracker (**Figure 5**, top left).

Comparison of the north/south gradients show also similar performances when comparing against CarbonTracker and TCCON. However, this should not be over interpreted because TCCON contributes only to few grid boxes especially on the southern hemisphere (**Figure 5**, top right). Most satellite retrievals have north/south gradients which are a few tenth of a ppm larger than in CarbonTracker and TCCON.




Figure 1: Validation of individual XCO₂ algorithms and EMMA v3.1 CO₂ with TCCON GGG2014.

C3S_312a_Lot6_IUP-UB_2016SC1 – Product Quality Assessment Report ANNEX-D v2



Table 2: Summarizing XCO₂ validation statistics for all TCCON sites that have been used for the validation. Listed are the number of co-locations (#), average single measurement precision, regional and seasonal accuracy, linear trend, year-to-year stability, and the probability that the accuracy and stability TR are met.

Algorithm	#	Precision [ppm]	Accuracy [ppm]		Stability [ppm/a]		Probability that TR is met [%]	
			Regional	Seasonal	Trend	Year2Year	Accuracy	Stability
BESD v02.01.02	18585	1.84	0.29	0.32	0.13±0.28	0.29	59	83
RemoTeC v2.3.8	12587	2.13	0.60	0.32	0.06±0.08	0.31	28	97
ACOS v7.3.10a	11985	1.74	0.58	0.32	-0.08±0.07	0.27	-	-
UoL-FP v7.2	11974	1.80	0.43	0.37	-0.10±0.09	0.29	42	96
NIES v02bc	8952	2.26	0.63	0.42	0.00±0.14	0.50	-	-
PPDF-S v02	8469	1.95	0.94	0.33	-0.15±0.12	0.37	-	-
NIES v02	14654	2.30	0.67	0.41	0.03±0.14	0.36	-	-
EMMA v3.1 CO2	15743	1.92	0.50	0.29	0.00±0.09	0.29	40	98




Figure 2: Anomaly of station biases (left) and station drift (right).




Figure 3: Stability analyses for EMMA and the contributing individual algorithms. The black curve shows the average station bias and the red curves its uncertainty represented by the station-to-station standard deviation and error propagation from single sounding measurement noise.

In terms of the frequency of potential outliers and standard deviation of the difference to CarbonTracker and TCCON, EMMA is usually under the top three performing individual algorithms (**Figure 5**, bottom).

Once the GOSAT algorithms kick in in 2009, ACOS usually provides the largest part of the integrated data weight in EMMA (**Figure 6**).

The average inter-algorithm spread has values most times between 0.4ppm and 1.6 ppm and is typically below 1.0ppm (**Figure 7**, left). The largest inter-algorithm spreads are observed in the tropics, Asia, and in high latitudes. Only a small fraction of the inter-algorithm spread can be explained with differences expected due to measurement noise so that most of the differences can be considered systematic. Only in high latitudes and at some coast-lines measurement noise is expected to explain a significant part of the inter-algorithm spread (**Figure 7**, right).

It is interesting to note that the average inter-algorithm spread usually reduced with every new EMMA version (always including the most recent algorithm versions, **Figure 8**). This means that EMMA observed kind of a convergence among the individual algorithms. It is not entirely clear



where this convergence is coming from and many effects may contribute to the explanation: algorithms are improved and bugs are removed but algorithms may also become more similar by using the same input data (e.g., spectroscopy, elevation model). However, the most recent version of EMMA does not follow this trend because the average inter-algorithm spread increased slightly. Most probably, this is caused by adding the not bias corrected operational NIES product and NIES' PPDF-S product. Additionally, the EMMA time period enhanced so that small drifts in the data sets, which are not corrected by EMMA's overall offset correction, can contribute to a larger extend.




Figure 4: Monthly gridded XCO₂ averages and inter-algorithm spread at the example of September 2009 for EMMA v3.1 CO₂.




Figure 5: Top left: Difference of seasonal cycle amplitude of all individual algorithms as well as EMMA compared with CarbonTracker v2017 and TCCON GGG2014. **Top right:** Difference of north/south gradient of all individual algorithms as well as EMMA compared with CarbonTracker v2017 and TCCON. **Bottom left:** Frequency of potential outliers estimated by large gradients, large differences to CarbonTracker, and large differences to EMMA. **Bottom right:** Standard deviation of the difference of all algorithms and EMMA to CarbonTracker and TCCON.




Figure 6: EMMA's normalized relative data weight proportional to $\sum 1/\sigma_i^2$ (**top**) and number of soundings (**bottom**) per algorithm and month.




Figure 7: Average inter-algorithm scatter of monthly $10^\circ \times 10^\circ$ averages from January 2003 till December 2017 (left) and corresponding expected contribution of measurement noise (right).




Figure 8: Average inter-algorithm spread of all EMMA versions compared with the approximately expected contribution of retrieval noise and a rough estimate of the representation error.



2.1.2 Summary

The validation results are summarized in **Table 3**.

Table 3: Product Quality Summary Table for product XCO2_EMMA.

Product Quality Summary Table for Product: XCO2_EMMA Level: 2, Version: 3.0, Time period covered: 1.2009 – 12.2017				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppm]	1.92	< 8 (T) < 3 (B) < 1 (G)	-	-
Uncertainty ratio in [-]: Ratio reported uncertainty to standard deviation of satellite-TCCON difference	1.00	-	-	No requirement but value close to unity expected for a high quality data product.
Mean bias [ppm]	0.48	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppm]	Spatial – spatiotemporal: 0.50 – 0.58	< 0.5	Probability that accuracy TR is met: 40%	-
Stability: Drift [ppm/year]	0.00 +/- 0.09 (1-sigma)	< 0.5	Probability that stability TR is met: 98%	-
Stability: Year-to-year bias variability [ppm/year]	0.29 (1-sigma)	< 0.5	-	-



2.2 XCH4_EMMA

2.2.1 Validation

Figure 9 shows all co-located EMMA and TCCON retrievals used for validation. Additionally, it includes all co-locations of the individual algorithms contributing to EMMA. The overall statistics per contributing algorithm are summarized in **Table 4**. **Table 5** shows the validation summary specifically for EMMA v3.1 CH4 i.e., the XCH4_EMMA product. The results are valid for the time periods covered by the individual algorithms but starting in 04/2010 (i.e., after the WFMD contribution to EMMA ended), respectively. Note that this significantly limits the validation period for WFMD which ends at the beginning of 2012. Therefore, the validation results for WFMD are less robust and do not cover the years 2003-2005 when the SCIAMACHY instrument performed best in respect to XCH4.

The individual algorithms have a single measurement precision in the range of 13.38ppb (UoL-FP) – 14.40ppb (PPDF-S) except for WFMD which has a single measurement precision of 100.54ppb. EMMA has a single measurement precision of 13.68ppb which is similar as for the GOSAT algorithms. EMMA's combined regional and seasonal biases (4.27ppb) are smaller than those of the individual algorithms ranging from (4.32ppb for UoL-FP to 14.59 for WFMD).

Figure 10 (left) shows the anomaly of station biases of the used algorithms. One can see that most satellite retrievals have a high bias of typically 4ppb at the Sodankylä and Park Falls TCCON sites and low bias of similar magnitude at the southern hemispheric sites Darwin and Wollongong. This feature is somewhat similar to the feature observed for XCO₂ and considerably contributes to the algorithms station-to-station bias statistics. Currently, it is unclear whether this discrepancy comes from the satellite retrievals or the TCCON.

The drift analysis in **Figure 10** (right) shows more or less small but consistent negative trends (typically below -0.5ppb/a) at the sites Lamont and Park Falls and positive trends of the same magnitude at Bremen and Sodankylä. This is a bit surprising because some of the sites are located in similar latitude bands. This feature is similar to the feature observed for the drifts of XCO₂.

Figure 11 shows the smoothed average residual difference (satellite - TCCON) which is not explained by station bias, seasonal bias, and/or linear drift. The year-to-year stability computed from the variability of the average is in the range of 1.7ppb/a (UoL-PR and NIES v02) to 2.4ppb/a (PPDF-S) for the GOSAT algorithms and 7.0ppb/a for WFMD. EMMA's year-to-year stability of 1.8ppb/a is at the lower end of the range of GOSAT algorithms.

Analyses of gridded L3 data show that all algorithms reproduce large scale features well, however, there are still differences of a some 10ppb when looking into the details (**Figure 12**).

All algorithms see a consistently larger north/south gradient than TCCON (**Figure 13**, left). The same is true for the seasonal amplitude (**Figure 13**, left). In terms of the standard deviation of the difference to TCCON and the frequency of potential outliers, EMMA performs similarly as the best individual algorithms (**Figure 13**).




Figure 9: Validation of individual XCH₄ algorithms and EMMA v3.1 CH₄ with TCCON GGG2014.



Table 4: Summarizing XCH₄ validation statistics for all TCCON sites that have been used for the validation. Listed are the number of co-locations (#), average single measurement precision, regional and seasonal accuracy, linear trend, year-to-year stability, and the probability that the accuracy and stability TR are met. Note, due to the short validation period of WFMD and the instrumental issues of SCIAMACHY during this period, the WFMD results are less robust and trend and probabilities for meeting the TRs have not been assessed.

Algorithm	#	Precision [ppb]	Accuracy [ppb]		Stability [ppb/a]		Probability that TR is met [%]	
			Regional	Seasonal	Trend	Year2Year	Accuracy	Stability
WFMD v4.0	13358	100.54	8.38	11.94	-	7.0	-	-
RemoTeC-FP v2.3.8	11288	13.98	4.58	3.24	0.02±0.53	1.8	100	99
RemoTeC-PR v2.3.9	35293	14.26	3.29	3.73	0.62±0.39	2.0	100	99
UoL-FP v7.2	10942	13.38	3.36	2.71	-0.56±0.98	1.9	100	95
UoL-PR v7.2	30592	13.68	4.42	2.01	0.05±0.53	1.7	100	99
NIES v02bc	8046	14.04	3.90	3.45	-0.64±1.42	1.9	-	-
PPDF-S v02	7512	14.40	6.66	3.41	-1.44±1.60	2.4	-	-
NIES v02	13802	13.95	4.08	3.19	-0.32±1.05	1.7	-	-
EMMA v3.1 CO2	20881	13.68	3.70	2.13	-0.05±0.67	1.8	100	99




Figure 10: Anomaly of station biases (left) and station drift (right).




Figure 11: Stability analyses for EMMA and the contributing individual algorithms. The black curve shows the average station bias and the red curves its uncertainty represented by the station-to-station standard deviation and error propagation from single sounding measurement noise.

Once the GOSAT algorithms kick in in 2009, the proxy algorithms RemoTeC-PR and UoL-PR clearly provide the largest part of the integrated data weight in EMMA (**Figure 14**). Note also the drop in the integrated data weight of WFMD end of 2005 due to instrument degradation.

The average inter-algorithm spread has values mostly between 2ppb and 12ppb and is typically below 9ppb (**Figure 15**, left). The largest inter-algorithm spreads are observed in the tropics, Asia, and in high latitudes. Only a small fraction of the inter-algorithm spread can be explained with differences expected due to measurement noise so that most of the differences can be considered systematic errors. Only in high latitudes and at some coast-lines measurement noise is expected to explain a significant part of the inter-algorithm spread (**Figure 15**, right).

The average inter-algorithm spread enhanced with the last EMMA versions (always including the most recent algorithm versions, **Figure 16**). The most recent enhancement is most probably caused by adding the not bias corrected operational NIES product and NIES' PPDF-S product. Additionally, the EMMA time period enhanced so that small drifts in the data sets, which are not corrected by EMMA's overall offset correction, can contribute to a larger extend.




Figure 12: Monthly gridded XCH₄ averages and inter-algorithm spread at the example of September 2010 for EMMA v3.1 CH₄.




Figure 13: Left: Difference of the north/south gradient, difference of the seasonal cycle amplitude, and standard deviation of the difference in respect to TCCON for all individual algorithms as well as EMMA. **Right:** Frequency of potential outliers estimated by large gradients, large differences to TCCON, and large differences to EMMA.




Figure 14: EMMA's normalized relative data weight proportional to $\sum 1/\sigma_i^2$ (**top**) and number of soundings (**bottom**) per algorithm and month.




Figure 15: Average inter-algorithm scatter of monthly 10°x10° averages from January 2003 till December 2017 (left) and corresponding expected contribution of measurement noise (right).




Figure 16: Average inter-algorithm spread of all EMMA versions compared with the approximately expected contribution of retrieval noise and a rough estimate of the representation error.



2.2.2 Summary

The validation results are summarized in **Table 5**.

Table 5: Product Quality Summary Table for product XCH4_EMMA. Note that the achieved performance corresponds to the time period after 03/2010 when only GOSAT algorithms are part of EMMA.

Product Quality Summary Table for Product: XCH4_EMMA Level: 2, Version: 3.0, Time period covered: 4.2010 – 12.2017				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppb]	13.68	< 34 (T) < 17 (B) < 9 (G)	-	
Uncertainty ratio) in [-]: Ratio reported uncertainty to standard deviation of satellite-TCCON difference	0.97	-	-	No requirement but value close to unity expected for a high quality data product.
Mean bias [ppb]	-6.80	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppb]	Spatial – spatiotemporal: 3.70 – 4.27	< 10	Probability that accuracy TR is met: 100%	-
Stability: Linear bias trend [ppb/year]	-0.05 +/- 0.67 (1-sigma)	< 3	Probability that stability TR is met: 99%	-
Stability: Year-to-year bias variability [ppb/year]	1.8 (1-sigma)	< 3	-	



References

- Boesch and Anand, 2017:** H. Boesch and J. Anand, Algorithm Theoretical Basis Document (ATBD) – ANNEX A for products CO₂_GOS_OCFP, CH₄_GOS_OCFP & CH₄_GOS_OCFR, Copernicus Climate Change Service (C3S) project on satellite-derived Essential Climate Variable (ECV) Greenhouse Gases (CO₂ and CH₄) data products (project C3S_312a_Lot6), Version 1 (21/08/2017), 2017
- Bril et al., 2012:** Bril, A.; Oshchepkov, S.; Yokota, T. Application of a probability density function-based atmospheric light-scattering correction to carbon dioxide retrievals from GOSAT over-sea observations. *Remote Sensing of the Environment* 2012, 117, 301–306
- Detmers, 2017a:** R. Detmers, Algorithm Theoretical Basis Document (ATBD) – ANNEX B for products CO₂_GOS_SRFP & CH₄_GOS_SRFP, Copernicus Climate Change Service (C3S) project on satellite-derived Essential Climate Variable (ECV) Greenhouse Gases (CO₂ and CH₄) data products (project C3S_312a_Lot6), Version 1 (21/08/2017), 2017
- Detmers, 2017b:** R. Detmers, Algorithm Theoretical Basis Document (ATBD) – ANNEX C for product CH₄_GOS_SRPR, Copernicus Climate Change Service (C3S) project on satellite-derived Essential Climate Variable (ECV) Greenhouse Gases (CO₂ and CH₄) data products (project C3S_312a_Lot6), Version 1 (21/08/2017), 2017
- O'Dell et al., 2012:** O'Dell, C. W., Connor, B., Bösch, H., O'Brien, D., Frankenberg, C., Castano, R., Christi, M., Eldering, D., Fisher, B., Gunson, M., McDuffie, J., Miller, C. E., Natraj, V., Oyafuso, F., Polonsky, I., Smyth, M., Taylor, T., Toon, G. C., Wennberg, P. O., and Wunch, D.: The ACOS CO₂ retrieval algorithm – Part 1: Description and validation against synthetic observations, *Atmos. Meas. Tech.*, 5, 99–121, doi:10.5194/amt-5-99-2012, 2012
- Reuter et al., 2011:** M. Reuter, H. Bovensmann, M. Buchwitz, J. P. Burrows, B. J. Connor, N. M. Deutscher, D. W. T. Griffith, J. Heymann, G. Keppel-Aleks, J. Messerschmidt, J. Notholt, C. Petri, J. Robinson, O. Schneising, V. Sherlock, V. Velasco, T. Warneke, P. O. Wennberg, D. Wunch: Retrieval of atmospheric CO₂ with enhanced accuracy and precision from SCIAMACHY: Validation with FTS measurements and comparison with model results. *Journal of Geophysical Research - Atmospheres*, 116, D04301, doi: 10.1029/2010JD015047, 2011
- Reuter et al., 2013:** M. Reuter, H. Bösch, H. Bovensmann, A. Bril, M. Buchwitz, A. Butz, J. P. Burrows, C. W. O'Dell, S. Guerlet, O. Hasekamp, J. Heymann, N. Kikuchi, S. Oshchepkov, R. Parker, S. Pfeifer, O. Schneising, T. Yokota, and Y. Yoshida: A joint effort to deliver satellite retrieved atmospheric CO₂ concentrations for surface flux inversions: the ensemble median algorithm EMMA. *Atmospheric Chemistry and Physics*, doi:10.5194/acp-13-1771-2013, 13, 1771-1780, 2013
- Yoshida et al., 2011:** Yoshida, Y., Ota, Y., Eguchi, N., Kikuchi, N., Nobuta, K., Tran, H., Morino, I., and Yokota, T.: Retrieval algorithm for CO₂ and CH₄ column abundances from short-wavelength infrared spectral observations by the Greenhouse gases observing satellite, *Atmos. Meas. Tech.*, 4, 717–734, doi:10.5194/amt-4-717-2011, 2011



ECMWF - Shinfield Park, Reading RG2 9AX, UK

Contact: info@copernicus-climate.eu