



Product Quality Assessment Report (PQAR) – ANNEX E for IASI CO₂ and CH₄ and AIRS CO₂ mid-tropospheric products

C3S_312a_Lot6_IUP-UB – Greenhouse Gases

Issued by: C. Crevoisier, LMD/CNRS, France

Date: 04/10/2018

Ref: C3S_D312a_Lot6.3.1.7-v2_PQAR_ANNEX-E_v2.0

Official reference number service contract: 2016/C3S_312a_Lot6_IUP-UB/SC1



This document has been produced in the context of the Copernicus Climate Change Service (C3S). The activities leading to these results have been contracted by the European Centre for Medium-Range Weather Forecasts, operator of C3S on behalf of the European Union (Delegation Agreement signed on 11/11/2014). All information in this document is provided "as is" and no guarantee or warranty is given that the information is fit for any particular purpose. The user thereof uses the information at its sole risk and liability. For the avoidance of all doubts, the European Commission and the European Centre for Medium-Range Weather Forecasts has no liability in respect of this document, which is merely representing the authors view.



Contributors

**INSTITUTE OF ENVIRONMENTAL PHYSICS (IUP),
UNIVERSITY OF BREMEN, BREMEN, GERMANY
(IUP)**

M. Buchwitz

**CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE (CNRS),
LABORATOIRE DE METEOROLOGIE DYNAMIQUE (LMD),
PALAISEAU, FRANCE
(LMD/CNRS)**

C. Crevoisier

R. Armante



Table of Contents

History of modifications	5
Related documents	6
Acronyms	7
General definitions	10
Scope of document	11
Executive summary	12
1. Product validation methodology	13
1.1 CH ₄ and CO ₂ mid-tropospheric column averaged mole fractions	13
1.2 Validation data and method	14
2. Validation Results	14
2.1 Products CO ₂ _IASA_NLIS and CO ₂ _IASB_NLIS	14
2.1.1 Validation	14
2.1.2 Validation summary	17
2.2 Products CH ₄ _IASA_NLIS and CH ₄ _IASB_NLIS	18
2.2.1 Validation	18
2.2.2 Validation summary	22
2.3 Product CO ₂ _AIRS_NLIS	23
2.3.1 Validation	23
2.3.2 Validation summary	24
References	25



History of modifications

Version	Date	Description of modification	Chapters / Sections
1.1	20-October-2017	New document for data set CDR1 (until 2016)	All
2.0	4-October-2018	Update for CDR2 (until 2017)	All



Related documents

Reference ID	Document
D1	<p>Main PQAR:</p> <p>Buchwitz, M., et al., Product Quality Assessment Report (PQAR) – Main document, C3S project C3S_312a_Lot6_IUP-UB – Greenhouse Gases, v2.0, 2018.</p> <p><i>(this document is an ANNEX to the Main PQAR)</i></p>



Acronyms

Acronym	Definition
AIRS	Atmospheric Infrared Sounder
AMSU	Advanced Microwave Sounding Unit
ATBD	Algorithm Theoretical Basis Document
BESD	Bremen optimal ESTimation DOAS
CAR	Climate Assessment Report
C3S	Copernicus Climate Change Service
CCDAS	Carbon Cycle Data Assimilation System
CCI	Climate Change Initiative
CDR	Climate Data Record
CDS	(Copernicus) Climate Data Store
CMUG	Climate Modelling User Group (of ESA's CCI)
CRG	Climate Research Group
D/B	Data base
DOAS	Differential Optical Absorption Spectroscopy
EC	European Commission
ECMWF	European Centre for Medium Range Weather Forecasting
ECV	Essential Climate Variable
EMMA	Ensemble Median Algorithm
ENVISAT	Environmental Satellite (of ESA)
EO	Earth Observation
ESA	European Space Agency
EU	European Union
EUMETSAT	European Organisation for the Exploitation of Meteorological Satellites
FCDR	Fundamental Climate Data Record
FoM	Figure of Merit
FP	Full Physics retrieval method
FTIR	Fourier Transform InfraRed
FTS	Fourier Transform Spectrometer
GCOS	Global Climate Observing System
GEO	Group on Earth Observation
GEOSS	Global Earth Observation System of Systems
GHG	GreenHouse Gas
GOME	Global Ozone Monitoring Experiment
GMES	Global Monitoring for Environment and Security
GOSAT	Greenhouse Gases Observing Satellite
IASI	Infrared Atmospheric Sounding Interferometer



IMAP-DOAS (or IMAP)	Iterative Maximum A posteriori DOAS
IPCC	International Panel in Climate Change
IUP	Institute of Environmental Physics (IUP) of the University of Bremen, Germany
JAXA	Japan Aerospace Exploration Agency
JCGM	Joint Committee for Guides in Metrology
L1	Level 1
L2	Level 2
L3	Level 3
L4	Level 4
LMD	Laboratoire de Météorologie Dynamique
MACC	Monitoring Atmospheric Composition and Climate, EU GMES project
NA	Not applicable
NASA	National Aeronautics and Space Administration
NetCDF	Network Common Data Format
NDACC	Network for the Detection of Atmospheric Composition Change
NIES	National Institute for Environmental Studies
NIR	Near Infra Red
NLIS	LMD/CNRS <i>neuronal</i> network mid/upper tropospheric CO ₂ and CH ₄ retrieval algorithm
NOAA	National Oceanic and Atmospheric Administration
Obs4MIPs	Observations for Climate Model Intercomparisons
OCO	Orbiting Carbon Observatory
OE	Optimal Estimation
PBL	Planetary Boundary Layer
ppb	Parts per billion
ppm	Parts per million
PR	(light path) PROxy retrieval method
PVIR	Product Validation and Intercomparison Report
QA	Quality Assurance
QC	Quality Control
REQ	Requirement
RMS	Root-Mean-Square
RTM	Radiative transfer model
SCIAMACHY	SCanning Imaging Absorption spectroMeter for Atmospheric Chartography
SCIATRAN	SCIAMACHY radiative transfer model
SRON	SRON Netherlands Institute for Space Research
SWIR	Short Wave Infra Red
TANSO	Thermal And Near infrared Sensor for carbon Observation
TANSO-FTS	Fourier Transform Spectrometer on GOSAT
TBC	To be confirmed



TBD	To be defined / to be determined
TCCON	Total Carbon Column Observing Network
TIR	Thermal Infra Red
TR	Target Requirements
TRD	Target Requirements Document
WFM-DOAS (or WFMD)	Weighting Function Modified DOAS
UoL	University of Leicester, United Kingdom
URD	User Requirements Document
WMO	World Meteorological Organization
Y2Y	Year-to-year (bias variability)



General definitions

Table 1 lists some general definitions relevant for this document.

Table 1: General definitions.

Item	Definition
XCO ₂	Column-averaged dry-air mixing ratios (mole fractions) of CO ₂
XCH ₄	Column-averaged dry-air mixing ratios (mole fractions) of CH ₄
L1	Level 1 satellite data product: geolocated radiance (spectra)
L2	Level 2 satellite-derived data product: Here: CO ₂ and CH ₄ information for each ground-pixel
L3	Level 3 satellite-derived data product: Here: Gridded CO ₂ and CH ₄ information, e.g., 5 deg times 5 deg, monthly
L4	Level 4 satellite-derived data product: Here: Surface fluxes (emission and/or uptake) of CO ₂ and CH ₄



Scope of document

This document is a Product Quality Assessment Report (PQAR) for the Copernicus Climate Change Service (C3S, <https://climate.copernicus.eu/>) component as covered by project C3S_312a_Lot6 led by University of Bremen, Germany.

Within project C3S_312a_Lot6 satellite-derived atmospheric carbon dioxide (CO₂) and methane (CH₄) Essential Climate Variable (ECV) data products will be generated and delivered to ECMWF for inclusion into the Copernicus Climate Data Store (CDS) from which users can access these data products and the corresponding documentation.

The C3S_312a_Lot 6 satellite-derived data products are:

- Column-average dry-air mixing ratios (mole fractions) of CO₂ and CH₄, denoted XCO₂ (in parts per million, ppm) and XCH₄ (in parts per billion, ppb), respectively.
- Mid/upper tropospheric mixing ratios of CO₂ (in ppm) and CH₄ (in ppb).

This document describes the validation / quality assessment of the C3S products CO2_IASA_NLIS, CH4_IASA_NLIS, CO2_IASB_NLIS, CH4_IASB_NLIS, CO2_AIRS_NLIS.

These products are mid-tropospheric CO₂ and CH₄ Level 2 products as retrieved from the IASI sensors on Metop-A and Metop-B and mid-tropospheric CO₂ from AIRS using algorithms developed at CNRS-LMD, France.



Executive summary

This document describes the performance for the Level 2 CO₂ and CH₄ data products retrieved from IASI observations at CNRS-LMD and delivered to the Copernicus Climate Change Service (C3S) . These products are mid-tropospheric-averaged dry-air mixing ratios (mole fractions) of CH₄ and CO₂, retrieved at 9:30 am/pm (local time) from observations made by the IASI and AMSU instruments onboard the European Metop-A (since July 2006) and Metop-B (since February 2013) platforms.

IASI and AIRS observations were spatially and temporally collocated with observations made from aircraft measurements from the CONTRAIL and HIPPO programs, as well as with observations made from balloons using AirCores. When enough in-situ data were available, a number of statistics, including accuracy and stability, have been computed from the difference between in-situ measurements and retrievals from space observation. Overall, the CNRS-LMD products are found to be highly stable and meet the Target Requirement (TR) requirements for accuracy and stability. It has to be noted that, due to too sparse a validation data for CH₄, the TR for stability could not be computed. This calls for continuous effort in performing and developing continuous airborne observations of greenhouse gases.



1. Product validation methodology

1.1 CH₄ and CO₂ mid-tropospheric column averaged mole fractions

The validation is performed for five Level 2 products:

- CO₂_IASA_NLIS: mid-tropospheric column averaged mole fractions of CO₂ retrieved from IASI onboard Metop-A.
- CO₂_IASB_NLIS: mid-tropospheric column averaged mole fractions of CO₂ retrieved from IASI onboard Metop-B.
- CH₄_IASA_NLIS: mid-tropospheric column averaged mole fractions of CH₄ retrieved from IASI onboard Metop-A.
- CH₄_IASB_NLIS: mid-tropospheric column averaged mole fractions of CH₄ retrieved from IASI onboard Metop-B.
- CO₂_AIRS_NLIS: mid-tropospheric column averaged mole fractions of CO₂ retrieved from AIRS onboard Aqua.

The four first products have been retrieved from simultaneous observations of the IASI and AMSU instruments flying together onboard the Metop satellites using a non-linear inference scheme using Multi-Layer Perceptrons with 2 hidden layers. IASI hyperspectral observations in the thermal infrared at 7.7 μm (resp. 15 μm), which are sensitive to both temperature and gas concentrations of CH₄ (resp. CO₂) are used in conjunction with microwave observations from the AMSU instruments, only sensitive to temperature, to decorrelate both signals (*Crevoisier et al., 2009a, 2009b, 2013*). The fifth product has been similarly obtained with AIRS and AMSU observations.

Potential radiative systematic biases existing between simulations used in the inference scheme and observations are computed for each channel by averaging, over the instruments full years of operation, the differences between simulations and collocated (in time and space) satellite observations. The simulations are performed using the 4A/OP-2009 forward model (*Scott and Chédin, 1981*; <http://www.noveltis.net/4AOP/>), which is based on the updated 2011 version of the GEISA spectroscopic database (available at <http://ether.ipsl.jussieu.fr/>) (*Jacquinet-Husson et al., 2011*), and radiosonde measurements from the Analyzed RadioSoundings Archive database (available at <http://ara.lmd.polytechnique.fr>). IASI calibrated radiance spectra (level1c) are received through the EUMETCast near real time data distribution service via the French Ether center (<http://ether.ipsl.jussieu.fr>).

The retrieved CO₂ and CH₄ integrated columns are weighted to the tropical mid-troposphere with peak sensitivity at about 230 hPa (~11 km), half the peak sensitivity at 100 and 500 hPa (~6 and 16 km), and no sensitivity to the surface. Retrievals are performed over land and sea, by night and day (9:30 am/pm local time) for clear-sky only (no clouds, no aerosols). The CO₂ retrievals are limited to the tropical region (30N:30S) because of the greater stability of the temperature atmospheric



profile, which helps decorrelating temperature from gas in the observed radiances, yielding a much better precision compared to the extratropics.

1.2 Validation data and method

Validation against high precision / low systematic errors reference observations is required for the mid/upper troposphere CO₂ and CH₄ data products. Unfortunately, measurements of both gases in the free troposphere and stratosphere are very sparse. Validation thus mostly relies on existing aircraft and airborne measurements.

A promising way consists in using 0-30 km profiles measured by balloon-borne AirCores (*Karion et al., 2010; Membrive et al., 2017*) to which averaging kernels can be applied to derive columns that can then be compared to those derived from space. So far, only a few profiles have been acquired, all in the northern hemisphere. In this validation exercise, use will be made of CH₄ profiles measured at Timmins (Ontario, Canada) and Sodankylä (Finland). Spanning 2014-2016, they will be used to validate both Metop-A and Metop-B retrievals (CH₄_IASA_NLIS and CH₄_IASB_NLIS).

2. Validation Results

2.1 Products CO₂_IASA_NLIS and CO₂_IASB_NLIS

2.1.1 Validation

2.1.1.1 Validation with aircraft measurements

Figure 1 shows comparison of IASI CO₂ mid-tropospheric columns with commercial aircraft measurements made as part of the CONTRAIL project (*Matsueda et al. 2008, Machida et al., 2008, Sawa et al., 2008*) as monthly means in 12 latitudinal bands of 5° each. Figure 2 shows the scatter plot of IASI CO₂ vs. CONTRAIL CO₂ for the whole period. The R correlation coefficient is 0.96, the bias and the standard deviation of the difference between both being 0.57 ± 0.99 ppmv.

To compute the various parameters summarized in the following tables, the time series in each latitudinal bands displayed in Figure 1 has been used separately.



Table 1. Mean and standard deviation of CO₂ (ppm): difference between CONTRAIL and IASI over 12 latitudinal bands of 5° each. Statistics over July 2007-December 2014.

Latitudinal band	30S: 25S	25S: 20S	20S: 15S	15S: 10S	10S: 5S	5S :EQ	EQ: 5N	5N: 10N	10N: 15N	15N: 20N	20N: 25N	25N: 30N
IASI- CONTRAIL	1.09 ± 1.04	0.92 ± 0.79	0.64 ± 0.68	0.34 ± 0.59	0.11 ± 0.78	0.26 ± 0.88	0.21 ± 0.80	0.08 ± 0.69	0.26 ± 0.71	0.56 ± 1.00	0.96 ± 1.17	1.51 ± 1.40

Table 1 shows the mean CONTRAIL-IASI CO₂ difference together with the associated standard deviation recorded in each latitudinal band. The mean single measurement is 0.99 ppm, while the mean CONTRAIL-IASI bias over all latitudinal band is 0.57 ppm.

Figure 1: Monthly variation of IASI mid-tropospheric CO₂ V4.0 (full line) and of CONTRAIL CO₂







Figure 2: Scatter plot of IASI mid-tropospheric CO₂ vs. CONTRAIL CO₂ measured at 10 km over the whole period depicted in Fig. 3.1 (July 2007 – December 2014) measured by aircraft at 10 km (dashed line) in 12 latitudinal bands of 5° each.



The relative systematic error is computed as the standard deviation of the CONTRAIL – IASI bias obtained in each latitudinal band. It is computed as two values:

- the “relative spatial bias”, which is the standard deviation of the mean per-latitudinal band bias computed over the entire time series. It comes to 0.46 ppm.
- The “relative spatio-temporal bias”, which is the standard deviation of the seasonal mean bias in each latitudinal band (i.e. JFM, AMJ, JAS, OND). It comes to 0.49 ppm.

For each latitudinal band, the linear drift was computed as the slope of the linear regression of the mean CONTRAIL –IASI bias against time. Table 2 shows the resulting drift and error. The main drift over the whole bands is: -0.01 ± 0.01 ppm/year.

Table 2. Mean and standard deviation of CO₂ (ppm): difference between CONTRAIL and IASI over 12 latitudinal bands of 5° each. Statistics over July 2007-December 2014.

Latitudinal band	30S: 25S	25S: 20S	20S: 15S	15S: 10S	10S: 5S	5S :EQ	EQ: 5N	5N: 10N	10N: 15N	15N: 20N	20N: 25N	25N: 30N
Linear drift [ppm/year]	-0.01	0.08	0.06	-0.02	-0.04	-0.09	-0.12	-0.07	-0.03	0.01	0.04	0.01

Finally, the year-to-year stability in each latitudinal band was computed as the difference between the maximum and the minimum values of the monthly differences within each year. This stability was found to be 2.64 ± 0.79 ppm/year.


2.1.1.2 Consistency between Metop-A and Metop-B

A direct comparison between mid-tropospheric CO₂ fields retrieved from Metop-A and Metop-B (version V4.2) yields a global bias close to zero and a standard deviation less than 0.1 ppm. The bias that was observed between the 2 products in version V4.1 has been corrected, as detailed in



Section 3.1.6 of the ATBD_ANNEX-E. Figure 3 shows the full time series of mid-tropospheric CO₂ retrieved from Metop-A and Metop-B. For most of the common period, the two products, which are recorded at the same local time but with a 180° shift in the orbit, are on top of each other. Only towards the end of Metop-A series (first half of 2015), a bias is seen between the 2. This is due to the slight degradation of AMSU channel 7 which started degrading beginning of 2015 and exceeded specifications in 2015.

Figure 3: Mid-tropospheric CO₂ retrieved from IASI/AMSU onboard Metop-A (blue) between July 2007 and June 2015, and from IASI/AMSU onboard Metop-B (green) between February 2013 and December 2017.



2.1.2 Validation summary

The validation results are summarized in the table below.

Table 2 - Product Quality Summary Table for product CO2_IASA_NLIS.

Product Quality Summary Table for Product: CO2_IASA_NLIS Level: 2, Version: 8.0, Time period covered: 7.2007 – 05.2015				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppm]	0.99	< 8 (T) < 3 (B) < 1 (G)	-	-
Mean bias [ppm]	0.57	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppm]	Spatial – spatiotemporal: 0.46 / 0.49	< 0.5	Probability that accuracy TR is met: 100%	-
Stability: Drift [ppm/year]	-0.01 ± 0.01 (1-sigma)	< 0.5	Probability that stability TR is met: 100%	-
Stability: Year-to-year bias variability [ppm/year]	2.64 ± 0.79 (1-sigma)	< 0.5	-	-

For CO₂_IASB_NLIS, it has not been possible to perform such an analysis yet, due to lack of coverage of the aircraft data for the corresponding years. As detailed in Section 2.1.1.2, a direct comparison between CO₂_IASA_NLIS and CO₂_IASB_NLIS has revealed an average bias of 0.0 ± 0.1 ppm between the two products for version V4.2. This comparison between the 2 products, measured at the same local time but with a 180° shift in the orbit, implies that the conclusion detailed in Table 3 for CO₂_IASA_NLIS apply for CO₂_IASB_NLIS as well.

2.2 Products CH₄_IASA_NLIS and CH₄_IASB_NLIS

2.2.1 Validation

For CH₄ products, only two quantities have been evaluated so far: single measurement precision, and mean bias with both aircraft and AirCore measurements. Due to limited time series of both aircraft (2 full years of CONTRAIL data for CH₄, only 17 AirCore profiles spanning 2014-2016), it has not yet been possible to evaluate the stability criteria.

2.2.1.1 Validation with aircraft measurements

Retrievals are compared with measurements made in the framework of the CONTRAIL project (*Machida et al., 2007, 2008; Matsueda et al., 2008; Sawa et al., 2008*). All IASI retrievals falling in a 5°x5° grid cell centered on each CONTRAIL measurement are averaged. Figure 4 shows the scatter plot of each pair of CONTRAIL / IASI CH₄. Over the whole dataset (311 pairs), the difference between CONTRAIL and IASI CH₄ is -0.89 ± 16.13 ppb, with a correlation R factor of 0.81.

Figure 4: CONTRAIL CH₄ vs. IASI CH₄ for all CONTRAIL measurements over July 2007-December 2014. The 1x1 line is shown as black. Difference between CONTRAIL and IASI CH₄ is -0.89 ± 16.13 ppb, with a correlation R factor of 0.81.




Figure 5 shows the monthly evolution of CH₄ as measured by CONTRAIL (dashed lines) and retrieved by IASI (full line) for 6 latitudinal bands of 10° each. The monthly evolution observed on both




datasets is consistent whatever the latitude is, both in terms of seasonality and amplitude. Table 3 summarizes the statistics (mean and standard deviation) obtained within each 7 latitudinal bands for IASI, CONTRAIL and the difference between both. Both datasets are statistically in agreement. Biases of less than 5 ppb, except for 20S:10S is observed. The standard deviations of IASI and CONTRAIL inside a given latitudinal band are noticeably close to each other, with again the exception of the 20S:10S band for which very few observations are available.

Table 4. Mean and standard deviation of CH₄ (ppb): as measured by CONTRAIL aircrafts, as retrieved by IASI, and difference between the two over 7 latitudinal bands of 10° each. Statistics over July 2007-December 2014.

Latitudinal band	30S:20S	20S:10S	10S:EQ	EQ:10N	10N:20N	20N:30S	30N:40N
CONTRAIL	1773.6 ± 12.1	1769.3 ± 9.3	1772.7 ± 10.0	1785.1 ± 13.3	1803.1 ± 18.5	1811.1 ± 14.9	1810.4 ± 22.7
IASI	1763.7 ± 13.3	1766.3 ± 19.8	1764.4 ± 11.0	1781.5 ± 22.5	1809.1 ± 21.0	1809.8 ± 16.1	1809.6 ± 19.8
IASI- CONTRAIL	9.9 ± 15.7	2.9 ± 14.0	8.2 ± 10.5	3.6 ± 19.8	-5.9 ± 18.0	1.31 ± 20.8	0.79 ± 27.3

Figure 5: Comparison between CONTRAIL and IASI CH₄ over July 2007-December 2014. Monthly evolution of CONTRAIL CH₄ (dashed line) and IASI CH₄ (full line) for 6 latitudinal bands of 10° each.






From Table 4, it is straightforward to compute the “relative spatial bias” of the “relative systematic error”, which is the standard deviation of the mean per-latitudinal band bias computed over the whole time series. The Accuracy is found to be 5.2 ppb. Due to several gaps in the time series, as well seen in Figure 5, it is not possible to compute the “relative spatio-temporal bias” which is the standard deviation of the seasonal mean bias in each latitudinal band.

2.2.1.2 Validation with AirCore 0-30 km profiles

Here, IASI CH₄ retrievals are compared to several AirCore profiles from measurements made at Timmins (Ontario, Canada) and Sodankylä (Finland) (*Membrive et al., 2017*). All IASI retrievals falling in a 5°x5° grid cell centered on each AirCore profile for the same day are averaged. Figure 6 shows the scatter plot of each pair of AirCore/IASI CH₄. Over the whole dataset (17 pairs for Metop-A and for Metop-B), the difference between AirCore and IASI CH₄ is 2.8 ± 8.5 ppb for Metop-A and 0.4 ± 9.9 ppb for Metop-B.

Figure 6: Comparison between IASI CH₄ (red for Metop-A and blue for Metop-B) and AirCore CH₄.




2.2.1.3 Comparison between CH₄_IASA_NLIS (Metop-A) and CH₄_IASB_NLIS (Metop-B)

A direct comparison between mid-tropospheric CH₄ fields retrieved from Metop-A and Metop-B yields a global bias close to zero and a standard deviation less than 5 ppbv. Figure 7 shows the seasonal maps (3 month average) of mid-tropospheric CH₄ for the year 2014 retrieved from Metop-



A and Metop-B, as well as the difference between the two and the associated standard deviation. The Metop-A and -B derived fields are close to each other. However, a small but negative bias over tropical oceans and a small but positive bias over Africa and to a lesser extent Australia can be observed on the map. These biases appear to be constant throughout the year. Such constant biases might be due to the fact that the systematic radiances for Metop-B used in the retrieval process have been computed over a reduced period, with very few radiosoundings over sea and



without taking into account any potential scan dependency. The availability of more years of observations to compute the radiative biases will help clarifying this point. The consistency of cloud detections between both satellites should also be assessed.

Figure 7: Seasonal maps (3 month average) of mid-tropospheric CH₄ as retrieved from Metop-A (1st column), Metop-B (2nd column), and mean (3rd column) and standard deviation (4th column) of the difference between Metop-A and Metop-B for the year 2014.



2.2.2 Validation summary

The validation results are summarized in the table below for CH4_IASA_NLIS. Please refer to Section 2.2.1.3 for comparison between CH4_IASA_NLIS and CH4_IASB_NLIS.

Table 3: Product Quality Summary Table for products CH4_IASA_NLIS (NC stands for Not computed due to lack of available data).

Product Quality Summary Table for Product: CH4_IASA_NLIS Level: 2, Version: 8.0, Time period covered: 7.2007 – 5.2015				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppb]	11.9	< 34 (T) < 17 (B) < 9 (G)	-	-
Mean bias [ppb]	-1.3	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: relative systematic error [ppb]	5.2	< 10	Probability that accuracy TR is met: 100%	-
Stability: Linear bias trend [ppb/year]	NC	< 3	NC	Time series of available aircraft/AirCore obs are not long enough to compute these 2 parameters
Stability: Year-to-year bias variability [ppb/year]	NC	< 3	-	



2.3 Product CO₂_AIRS_NLIS


2.3.1 Validation

Use is made of in situ observations made by commercial airliners from April 1993 to March 2007 between Japan and Australia (data available at <http://gaw.kishou.go.jp/wdcgg.html>). These observations, partly analyzed by *Matsueda et al. (2002)*, are available on a monthly basis. They cover the altitude range 9–13 km. Several gaps have affected the measurements throughout the period, which prevents making robust statistics from them.

The seasonal cycles of CO₂ measured by JAL are plotted in Fig. 8 for the period 2005-2006 (only period with full coverage). AIRS retrieved CO₂ cycle is also plotted in Fig. 8. There is a good agreement between both datasets in terms of the phase and amplitude of the seasonal cycle. Overall, the JAL – AIRS CO₂ difference is -0.43 ± 1.32 ppm.

Beginning of mid-2006, the bias between aircraft and AIRS CO₂ increases (negative sign), up to July 2007 when AIRS channels used to perform the retrievals were lost. This might be due to a non-corrected trend, which has affected either the AMSU observations in late 2006 and 2007, or some of the AIRS channels, which started exceeding radiometric specifications.

Figure 8: CO₂ seasonal cycle in the northern tropics as measured by JAL/CONTRAIL aircrafts (black) and as retrieved from AIRS (orange).





2.3.2 Validation summary

Table 6: Product Quality Summary Table for products CO2_AIRS_NLIS.

Product Quality Summary Table for Product: CO2_AIRS_NLIS Level: 2, Version: 3.0, Time period covered: 4.2003 – 7.2007				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppb]	1.32	< 8 (T) < 3 (B) < 1 (G)	-	-
Mean bias [ppb]	-0.43	-	-	No requirement but value close to zero expected for a high quality data product.



References

- Buchwitz et al., 2017:** ESA Climate Change Initiative (CCI) Product Validation and Intercomparison Report (PVIR) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG) for data set Climate Research Data Package No. 4 (CRDP#4), Version 5.0, 9. Feb. 2017, link: http://www.esa-ghg-cci.org/?q=webfm_send/352, 2017.
- Chédin et al. 2003:** Chédin, A., Saunders, R., Hollingsworth, A., Scott, N. A., Matricardi, M., Etcheto, J., Clerbaux, C., Armante, R. and Crevoisier, C.: The feasibility of monitoring CO₂ from high resolution infrared sounders. *J. Geophys. Res.*, 108, ACH 6-1–6-19, doi: 10.1029/2001JD001443, 2003.
- Chevallier et al., 2005:** Chevallier, F., R. J. Engelen, and P. Peylin, The contribution of AIRS data to the estimation of CO₂ sources and sinks. *Geophys. Res. Lett.*, 32, L23801, doi:10.1029/2005GL024229, 2005.
- CMUG-RBD, 2010:** Climate Modelling User Group Requirements Baseline Document, Deliverable 1.2, Number D1.2, Version 1.3, 2 Nov 2010.
- Cressot et al., 2014:** Cressot, C., F. Chevallier, P. Bousquet, et al., On the consistency between global and regional methane emissions inferred from SCIAMACHY, TANSO-FTS, IASI and surface measurements, *Atmos. Chem. Phys.*, 14, 577-592, 2014.
- Crevoisier et al., 2004:** Crevoisier, C., S. Heilliette, A. Chédin, S. Serrar, R. Armante, and N. A. Scott, Midtropospheric CO₂ concentration retrieval from AIRS observations in the tropics, *Geophys. Res. Lett.*, 31, L17106, doi:10.1029/2004GL020141, 2004.
- Crevoisier et al., 2009:** Crevoisier, C., Chédin, A., Matsueda, H., et al., First year of upper tropospheric integrated content of CO₂ from IASI hyperspectral infrared observations, *Atmos. Chem. Phys.*, 9, 4797-4810, 2009.
- Crevoisier et al. 2009b:** Crevoisier, C., Nobileau, D., Fiore, A., Armante, R., Chédin, A., and Scott, N. A.: Tropospheric methane in the tropics – first year from IASI hyperspectral infrared observations, *Atmos. Chem. Phys.*, 9, 6337–6350, doi:10.5194/acp-9-6337-2009, 2009b.
- Crevoisier et al., 2013:** Crevoisier, C., Nobileau, D., Armante, R., et al., The 2007–2011 evolution of tropical methane in the mid-troposphere as seen from space by MetOp-A/IASI, *Atmos. Chem. Phys.*, 13, 4279-4289, 2013.
- GCOS-154:** Global Climate Observing System (GCOS), SYSTEMATIC OBSERVATION REQUIREMENTS FOR SATELLITE-BASED PRODUCTS FOR CLIMATE, Supplemental details to the satellite-based component of the “Implementation Plan for the Global Observing System for Climate in Support of the UNFCCC (2010 update)”, Prepared by World Meteorological Organization (WMO), Intergovernmental Oceanographic Commission, United Nations Environment Programme (UNEP), International Council for Science, Doc.: GCOS 154, ink: <https://www.wmo.int/pages/prog/gcos/Publications/gcos-154.pdf>, 2010.
- GCOS-200:** The Global Observing System for Climate: Implementation Needs, World Meteorological Organization (WMO), GCOS-200 (GOOS-214), pp. 325, link: http://unfccc.int/files/science/workstreams/systematic_observation/application/pdf/gcos_ip_10oct2016.pdf, 2016.



- Jacquinet-Husson et al. 2011:** Jacquinet-Husson N., L. Crépeau, R. Armante, C. Boutammine, A. Chédin, et al.. The 2009 edition of the GEISA spectroscopic database, *J. Quant. Spectrosc. Radiat. Transfer*, 112, 2395–2445 doi:10.1016/j.jqsrt.2011.06.004, 2011.
- Machida et al. 2008:** Machida, T., Matsueda, H., Sawa, Y., Nakagawa, Y., Hirokuni, K., Kondo, N., Goto, K., Nakazawa, T., Ishikawa, K., and Ogawa, T.: Worldwide measurements of atmospheric CO₂ and other trace gas species using commercial airlines, *J. Atmos. Ocean. Tech.*, 25(10), 1744–1754, doi:10.1175/2008JTECHA1082.1, 2008.
- Matsueda et al. 2008:** Matsueda, H., Machida, T., Sawa, Y., Nakagawa, Y., Hirokuni, K., Ikeda, H., Kondo, N., and Goto, K.: Evaluation of atmospheric CO₂ measurements from new flask air sampling of JAL airliner observation, *Pap. Meteorol. Geophys.*, 59, 1–17, 2008.
- Membrive et al. 2017:** Membrive, O., Crevoisier, C., Sweeney, C., Danis, F., Hertzog, A., Engel, A., Bönisch, H., and Picon, L.: AirCore-HR: a high-resolution column sampling to enhance the vertical description of CH₄ and CO₂, *Atmos. Meas. Tech.*, 10, 2163–2181, <https://doi.org/10.5194/amt-10-2163-2017>, 2017.
- Scott and Chédin 1981:** Scott N.A. and Chédin A., A fast line-by-line method for atmospheric absorption computations: The Automatized Atmospheric Absorption Atlas, *J. Appl. Meteor.*, 20, n° 7, pp. 802–812, 1981.
- TRD GHG, 2017:** Buchwitz, M., Aben, I., Anand, J., Armante, R., Boesch, H., Crevoisier, C., Detmers, R. G., Hasekamp, O. P., Reuter, M., Schneising-Weigel, O., Target Requirement Document, Copernicus Climate Change Service (C3S) project on satellite-derived Essential Climate Variable (ECV) Greenhouse Gases (CO₂ and CH₄) data products (project C3S_312a_Lot6), Version 1, 28-March-2017, pp. 52, 2017.
- Wofsy et al. 2012:** Wofsy, S. C., Daube, B. C., Jimenez, R., et al.: HIPPO Merged 10-second Meteorology, Atmospheric Chemistry, Aerosol Data (R 20121129), Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA, http://dx.doi.org/10.3334/CDIAC/hippo_010 (Release 29 November 2012), 2012.
- Yoshida et al. 2013:** Yoshida, Y., Kikuchi, N., Morino, I., Uchino, O., Oshchepkov, S., Bril, A., Saeki, T., Schutgens, N., Toon, G. C., Wunch, D., Roehl, C. M., Wennberg, P. O., Griffith, D. W. T., Deutscher, N. M., Warneke, T., Notholt, J., Robinson, J., Sherlock, V., Connor, B., Rettinger, M., Sussmann, R., Ahonen, P., Heikkinen, P., Kyrö, E., Mendonca, J., Strong, K., Hase, F., Dohe, S., and Yokota, T.: Improvement of the retrieval algorithm for GOSAT SWIR XCO₂ and XCH₄ and their validation using TCCON data, *Atmos. Meas. Tech.*, 6, 1533–1547, doi:10.5194/amt-6-1533-2013, 2013.

