



Copernicus Climate Change Service



# **Product User Guide and Specification (PUGS) – ANNEX D for products XCO2\_EMMA and XCH4\_EMMA (v3.1, 2003-2017)**

## **C3S\_312a\_Lot6\_IUP-UB – Greenhouse Gases**

Issued by: Maximilian Reuter, University of Bremen,  
Institute of Environmental Physics (IUP)

Date: 04/10/2018

Ref: C3S\_D312a\_Lot6.3.1.5-v2\_PUGS\_ANNEX-D\_v2.0

Official reference number service contract: 2016/C3S\_312a\_Lot6\_IUP-UB/SC1



*This document has been produced in the context of the Copernicus Climate Change Service (C3S).*

*The activities leading to these results have been contracted by the European Centre for Medium-Range Weather Forecasts, operator of C3S on behalf of the European Union (Delegation Agreement signed on 11/11/2014). All information in this document is provided "as is" and no guarantee or warranty is given that the information is fit for any particular purpose.*

*The user thereof uses the information at its sole risk and liability. For the avoidance of all doubts, the European Commission and the European Centre for Medium-Range Weather Forecasts has no liability in respect of this document, which is merely representing the authors view.*



## Contributors

**INSTITUTE OF ENVIRONMENTAL PHYSICS (IUP),  
UNIVERSITY OF BREMEN, BREMEN, GERMANY  
(IUP)**

M. Reuter

M. Buchwitz

O. Schneising-Weigel



## Table of Contents

<b>History of modifications</b>	<b>5</b>
<b>Related documents</b>	<b>6</b>
<b>Acronyms</b>	<b>7</b>
<b>General definitions</b>	<b>11</b>
<b>Scope of document</b>	<b>12</b>
<b>Executive summary</b>	<b>13</b>
<b>1. Product description</b>	<b>14</b>
<b>2. Target requirements</b>	<b>16</b>
<b>3. Data usage information</b>	<b>17</b>



## History of modifications

Version	Date	Description of modification	Chapters / Sections
1.3	20-October-2017	New document for data set CDR1 (2003-2016)	All
2.0	4-October-2018	Update for CDR2 (2003-2017)	All



## Related documents

Reference ID	Document
D1	Main PUGS: Buchwitz, M., et al., Product User Guide and Specification (PUGS) – Main document, C3S project C3S_312a_Lot6_IUP-UB – Greenhouse Gases, v2.0, 2018. <i>(this document is an ANNEX to the Main ATBD)</i>
D2	Corresponding ATBD: <b>ATBD ANNEX D:</b> Reuter, M., et al., Algorithm Theoretical Basis Document (ATBD) – ANNEX D for products XCO <sub>2</sub> _EMMA and XCH <sub>4</sub> _EMMA (v3.1, 2003-2017), C3S project C3S_312a_Lot6_IUP-UB – Greenhouse Gases, v2.0, 2018.
D3	<b>TRD GHG, 2017:</b> Buchwitz, M., Aben, I., Anand, J., Armante, R., Boesch, H., Crevoisier, C., Detmers, R. G., Hasekamp, O. P., Reuter, M., Schneising-Weigel, O., Target Requirement Document, Copernicus Climate Change Service (C3S) project on satellite-derived Essential Climate Variable (ECV) Greenhouse Gases (CO <sub>2</sub> and CH <sub>4</sub> ) data products (project C3S_312a_Lot6), Version 1.3, 20-October-2017, pp. 53, 2017.



## Acronyms

Acronym	Definition
AIRS	Atmospheric Infrared Sounder
AMSU	Advanced Microwave Sounding Unit
ATBD	Algorithm Theoretical Basis Document
BESD	Bremen optimal ESTimation DOAS
CAR	Climate Assessment Report
C3S	Copernicus Climate Change Service
CCDAS	Carbon Cycle Data Assimilation System
CCI	Climate Change Initiative
CDR	Climate Data Record
CDS	(Copernicus) Climate Data Store
CMUG	Climate Modelling User Group (of ESA's CCI)
CRG	Climate Research Group
D/B	Data base
DOAS	Differential Optical Absorption Spectroscopy
EC	European Commission
ECMWF	European Centre for Medium Range Weather Forecasting
ECV	Essential Climate Variable
EMMA	Ensemble Median Algorithm
ENVISAT	Environmental Satellite (of ESA)
EO	Earth Observation
ESA	European Space Agency
EU	European Union



EUMETSAT	European Organisation for the Exploitation of Meteorological Satellites
FCDR	Fundamental Climate Data Record
FoM	Figure of Merit
FP	Full Physics retrieval method
FTIR	Fourier Transform InfraRed
FTS	Fourier Transform Spectrometer
GCOS	Global Climate Observing System
GEO	Group on Earth Observation
GEOSS	Global Earth Observation System of Systems
GHG	GreenHouse Gas
GOME	Global Ozone Monitoring Experiment
GMES	Global Monitoring for Environment and Security
GOSAT	Greenhouse Gases Observing Satellite
IASI	Infrared Atmospheric Sounding Interferometer
IMAP-DOAS (or IMAP)	Iterative Maximum A posteriori DOAS
IPCC	International Panel in Climate Change
IUP	Institute of Environmental Physics (IUP) of the University of Bremen, Germany
JAXA	Japan Aerospace Exploration Agency
JCGM	Joint Committee for Guides in Metrology
L1	Level 1
L2	Level 2
L3	Level 3
L4	Level 4
LMD	Laboratoire de Météorologie Dynamique
MACC	Monitoring Atmospheric Composition and Climate, EU GMES project
NA	Not applicable
NASA	National Aeronautics and Space Administration





NetCDF	Network Common Data Format
NDACC	Network for the Detection of Atmospheric Composition Change
NIES	National Institute for Environmental Studies
NIR	Near Infra Red
NLIS	LMD/CNRS <i>neural</i> network mid/upper tropospheric CO <sub>2</sub> and CH <sub>4</sub> retrieval algorithm
NOAA	National Oceanic and Atmospheric Administration
Obs4MIPs	Observations for Climate Model Intercomparisons
OCO	Orbiting Carbon Observatory
OE	Optimal Estimation
PBL	Planetary Boundary Layer
ppb	Parts per billion
ppm	Parts per million
PR	(light path) PROxy retrieval method
PVIR	Product Validation and Intercomparison Report
QA	Quality Assurance
QC	Quality Control
REQ	Requirement
RMS	Root-Mean-Square
RTM	Radiative transfer model
SCIAMACHY	SCanning Imaging Absorption spectroMeter for Atmospheric ChartographY
SCIATRAN	SCIAMACHY radiative transfer model
SRON	SRON Netherlands Institute for Space Research
SWIR	Short Wava Infra Red
TANSO	Thermal And Near infrared Sensor for carbon Observation
TANSO-FTS	Fourier Transform Spectrometer on GOSAT
TBC	To be confirmed
TBD	To be defined / to be determined



TCCON	Total Carbon Column Observing Network
TIR	Thermal Infra Red
TR	Target Requirements
TRD	Target Requirements Document
WFM-DOAS (or WFMD)	Weighting Function Modified DOAS
UoL	University of Leicester, United Kingdom
URD	User Requirements Document
WMO	World Meteorological Organization
Y2Y	Year-to-year (bias variability)



## General definitions

**Table 1** lists some general definitions relevant for this document.

**Table 1:** General definitions.

Item	Definition
XCO <sub>2</sub>	Column-averaged dry-air mixing ratios (mole fractions) of CO <sub>2</sub>
XCH <sub>4</sub>	Column-averaged dry-air mixing ratios (mole fractions) of CH <sub>4</sub>
L1	Level 1 satellite data product: geolocated radiance (spectra)
L2	Level 2 satellite-derived data product: Here: XCO <sub>2</sub> and XCH <sub>4</sub> information for each ground-pixel
L3	Level 3 satellite-derived data product: Here: Gridded XCO <sub>2</sub> and XCH <sub>4</sub> information, e.g., 5°x5°, monthly
L4	Level 4 satellite-derived data product: Here: Surface fluxes (emission and/or uptake) of CO <sub>2</sub> and CH <sub>4</sub>



## Scope of document

This document is an ANNEX to a Product User Guide and Specification (PUGS, see *D1*) for the Copernicus Climate Change Service (C3S, <https://climate.copernicus.eu/>) component as covered by project C3S\_312a\_Lot6 led by University of Bremen, Germany.

Within project C3S\_312a\_Lot6 satellite-derived atmospheric carbon dioxide (CO<sub>2</sub>) and methane (CH<sub>4</sub>) Essential Climate Variable (ECV) data products will be generated and delivered to ECMWF for inclusion into the Copernicus Climate Data Store (CDS) from which users can access these data products and the corresponding documentation.

The C3S\_312a\_Lot 6 satellite-derived data products are:

- Column-averaged dry-air mixing ratios (mole fractions) of CO<sub>2</sub> and CH<sub>4</sub>, denoted XCO<sub>2</sub> (in parts per million, ppm) and XCH<sub>4</sub> (in parts per billion, ppb), respectively.
- Mid/upper tropospheric mixing ratios of CO<sub>2</sub> (in ppm) and CH<sub>4</sub> (in ppb).

This document describes the C3S products XCO<sub>2</sub>\_EMMA and XCH<sub>4</sub>\_EMMA.

These products are merged multi-sensor XCO<sub>2</sub> and XCH<sub>4</sub> Level 2 products generated using algorithms developed at University of Bremen, Germany (see *D2*).



## Executive summary

The EMMA database consists of individual level 2 soundings retrieved by algorithms which can change from grid box to grid box and month to month. Therefore, it can be used in the same manner as any other XCO<sub>2</sub> or XCH<sub>4</sub> satellite retrieval, i.e., the EMMA database includes all information needed for inverse modeling (geo-location, time, XCO<sub>2</sub> or XCH<sub>4</sub>, averaging kernels, etc.).

The data fields and guidance on their use are provided in the main PUGS document (*D1*) describing, e.g., the common variables of all XCO<sub>2</sub> and XCH<sub>4</sub> L2 data sets provided by the Copernicus C3S\_312a\_Lot6 sub-project.

Additionally, to the common variables, EMMA includes information on, e.g., the inter-algorithm spread which informs about potential regional uncertainties and on the source-algorithm of each individual sounding within the EMMA data base. Such variables are subject to this ANNEX describing only the EMMA v3.1 CO<sub>2</sub> and EMMA v3.1 CH<sub>4</sub> specific aspects of the EMMA L2 data base.



## 1. Product description

Additionally to the common variables described in the main PUGS document (*D1*) the EMMA L2 data base includes the variables listed in **Table 2** and described in the following.

**Table 2:** EMMA v3.1 CO<sub>2</sub> and EMMA v3.1 CH<sub>4</sub> specific variables.

Name	Type	Dimension	Units	Short Description
median_processor_id	Integer	n	[-]	A unique ID for each L2 algorithm contributing to EMMA
median_uncertainty	Float	n	For XCO2: ppm, i.e., 10 <sup>-6</sup> For XCH4: ppb, i.e., 10 <sup>-9</sup>	Inter algorithm spread defined as standard deviation of the L3 products in the corresponding grid box (see <i>D2</i> )
median_uncertainty_se	Float	n	For XCO2: ppm, i.e., 10 <sup>-6</sup> For XCH4: ppb, i.e., 10 <sup>-9</sup>	Standard error of the median uncertainty (see <i>D2</i> )
median_uncertainty_ex	Float	n	For XCO2: ppm, i.e., 10 <sup>-6</sup> For XCH4: ppb, i.e., 10 <sup>-9</sup>	Inter-algorithm spread as expected from measurement noise (see <i>D2</i> )
<xco2 xch4>_accuracy	Float	n	For XCO2: ppm, i.e., 10 <sup>-6</sup> For XCH4: ppb, i.e., 10 <sup>-9</sup>	Potential spatio/temporal XCO2 or XCH4 bias (1-sigma) estimated from TCCON co-locations (see <i>D2</i> )
contributing_algorithms	Byte	n	[-]	Number of L2 algorithms contributing to median calculation in a specific grid box

**Table 3:** Unique L2 algorithm IDs used in EMMA v3.1 CO<sub>2</sub> and CH<sub>4</sub>.

ID	Quantity	Name	Version	Institution
1	XCO2	BESD	v02.01.02	IUP
2	XCO2	ACOS	v7.3.10a	NASA
3	XCO2	RemoTeC	v2.3.8	SRON
4	XCO2	UoL-FP	v7.2	UoL
5	XCO2	PPDF-S	v02	NIES
6	XCO2	NIES	v02bc (bias corrected)	NIES
-6	XCO2	NIES	v02	NIES
30	XCH4	RemoTeC-PR	v2.3.9	SRON
31	XCH4	RemoTeC-FP	v2.3.8	SRON
40	XCH4	UoL-PR	v7.2	UoL
41	XCH4	UoL-FP	v7.2	UoL



50	XCH4	PPDF-S	v02	NIES
60	XCH4	NIES	v02bc (bias corrected)	NIES
-60	XCH4	NIES	v02	NIES
70	XCH4	WFMD	v4.0	IUP

### Description of each parameter:

#### ***median\_processor\_id***

A unique ID for each L2 algorithm contributing to EMMA v3.1 CO<sub>2</sub> and CH<sub>4</sub>. See listing in **Table 3**.

#### ***median\_uncertainty***

Inter algorithm spread defined as standard deviation of the L3 products in the corresponding grid box (see D2).

#### ***median\_uncertainty\_se***

Standard error of the median uncertainty (see D2).

#### ***median\_uncertainty\_ex***

Inter-algorithm spread as expected from measurement noise (see D2).

#### ***<xco2/xch4>\_accuracy***

Potential spatio/temporal XCO<sub>2</sub> or XCH<sub>4</sub> bias (1-sigma) estimated from TCCON co-locations (see D2).

#### ***contributing\_algorithms***

Number of L2 algorithms contributing to median calculation in a specific grid box.



## 2. Target requirements

The target requirements for these products are described in the Target Requirement Document (TRD) (*TRD GHG, 2017*), see *D3*.







### 3. Data usage information

The EMMA database consists of individual level 2 soundings which can be used in the same manner as any other XCO<sub>2</sub> or XCH<sub>4</sub> satellite retrieval, i.e., the EMMA database includes all information needed for inverse modelling (geo-location, time, XCO<sub>2</sub> or XCH<sub>4</sub>, averaging kernels, etc.). The main PUGS document (*D1*) provides guidance on how to use the information.

**Figure 1** shows for an example month (September 2009) the EMMA XCO<sub>2</sub> and XCH<sub>4</sub> and the corresponding algorithm selected by the median (see *D2*). **Figure 2** shows the average inter-algorithm spread (01/2003 – 12/2017) and the expected average inter-algorithm spread due to measurement noise for XCO<sub>2</sub> and XCH<sub>4</sub>.



**Figure 1:** EMMA L2 XCO<sub>2</sub> and XCH<sub>4</sub> (**left**) and corresponding selected algorithm (**right**) for EMMA v3.1 CO<sub>2</sub> (**top**) and EMMA v3.1 CH<sub>4</sub> (**bottom**) at the example of September 2009.



**Figure 2:** Average inter-algorithm spread (01/2003 – 12/2017) (**left**) and expected average inter-algorithm spread due to measurement noise (**right**) for EMMA v3.1 CO<sub>2</sub> (**top**) and EMMA v3.1 CH<sub>4</sub> (**bottom**).



ECMWF - Shinfield Park, Reading RG2 9AX, UK

Contact: [info@copernicus-climate.eu](mailto:info@copernicus-climate.eu)