



Copernicus Climate Change Service



Product Quality Assessment Report (PQAR) – Main document for Greenhouse Gas (GHG: CO₂ & CH₄) data set CDR 3 (2003-2018)

C3S_312b_Lot2_DLR – Atmosphere

Issued by: Michael Buchwitz, University of Bremen,
Institute of Environmental Physics (IUP)

Date: 03/11/2019

Ref: C3S_D312b_Lot2.2.2.3-v1.0_PQAR-GHG_MAIN_v3.1

Official reference number service contract: 2018/C3S_312b_Lot2_DLR/SC1



This document has been produced in the context of the Copernicus Climate Change Service (C3S). The activities leading to these results have been contracted by the European Centre for Medium-Range Weather Forecasts, operator of C3S on behalf of the European Union (Delegation Agreement signed on 11/11/2014). All information in this document is provided "as is" and no guarantee or warranty is given that the information is fit for any particular purpose. The user thereof uses the information at its sole risk and liability. For the avoidance of all doubts, the European Commission and the European Centre for Medium-Range Weather Forecasts has no liability in respect of this document, which is merely representing the authors view.



Contributors

**INSTITUTE OF ENVIRONMENTAL PHYSICS (IUP),
UNIVERSITY OF BREMEN, BREMEN, GERMANY
(IUP)**

M. Buchwitz

M. Reuter

O. Schneising-Weigel

**SRON NETHERLANDS INSTITUTE FOR SPACE RESEARCH,
UTRECHT, THE NETHERLANDS
(SRON)**

I. Aben

L. Wu

O. P. Hasekamp

**UNIVERSITY OF LEICESTER, LEICESTER, UK
(UoL)**

H. Boesch

A. Di Noia

J. Anand

**CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE (CNRS),
LABORATOIRE DE METEOROLOGIE DYNAMIQUE (LMD),
PALAISEAU, FRANCE
(LMD/CNRS)**

C. Crevoisier

R. Armante



Table of Contents

History of modifications	6
Related documents	7
Acronyms	8
General definitions	11
Scope of document	13
Executive summary	16
1. Overview data products and instruments	22
1.1 Column-average mixing ratios of CO₂ and CH₄ (XCO₂ and XCH₄)	23
1.1.1 Overview	23
1.1.2 Instruments	23
1.1.3 XCO ₂	26
1.1.4 XCH ₄	27
1.1.5 List of XCO ₂ and XCH ₄ data products	28
1.2 Mid-tropospheric mixing ratios of CO₂ and CH₄	30
1.2.1 Overview	30
1.2.2 Instruments	30
1.2.3 CO ₂	31
1.2.4 CH ₄	31
1.2.5 List of mid-tropospheric CO ₂ and CH ₄ data products	32
2. Product validation methodology	33
2.1 Description of reference data used for validation	33
2.1.1 Reference data for validation of the XCO ₂ and XCH ₄ Level 2 products	33
2.1.2 Reference data for validation of the mid/upper tropospheric CO ₂ and CH ₄ products	34
2.2 Description of product validation methodology	35
2.2.1 Methods for validation of XCO ₂ and XCH ₄ Level 2 products	35
2.2.2 Methods for validation of XCO ₂ and XCH ₄ Level 3 Obs4MIPs products	45
2.2.3 Methods for validation of CO ₂ and CH ₄ Level 2 mid/upper troposphere products	46
3. Validation results	48
3.1 Validation results for Level 2 XCO₂ products	48
3.1.1 Validation results for product CO2_SCI_BESD	49
3.1.2 Validation results for product CO2_SCI_WFMD	53



3.1.3 Validation results for product CO2_GOS_OCFP	55
3.1.4 Validation results for product CO2_GOS_SRFp	57
3.1.5 Validation results for product XCO2_EMMA	59
3.2 Validation results of Level 2 XCH₄ products	61
3.2.1 Validation results for product CH4_SCI_WFMD	62
3.2.2 Validation results for product CH4_SCI_IMAP	65
3.2.3 Validation results for product CH4_GOS_OCPR	67
3.2.4 Validation results for product CH4_GOS_SRPR	69
3.2.5 Validation results for product CH4_GOS_OCFP	71
3.2.6 Validation results for product CH4_GOS_SRFp	73
3.2.7 Validation results for product XCH4_EMMA	75
3.3 Validation results for Level 3 XCO₂ product	77
3.4 Validation results for Level 3 XCH₄ products	79
3.5 Validation results for Level 2 mid-tropospheric products	81
4. Application(s) specific assessments	84
5. Compliance with user requirements	87
5.1 Level 2 XCO ₂ and XCH ₄ products	87
5.2 Level 3 XCO ₂ and XCH ₄ products	88
5.3 Level 2 mid-tropospheric products	89
References	90
6. Acknowledgement	101
7. List of ANNEXes	102
7.1 ANNEX A: PQAR for products CO2_GOS_OCFP, CH4_GOS_OCFP, CH4_OCPR	102
7.2 ANNEX B: PQAR for products CO2_GOS_SRFp, CH4_GOS_SRFp	102
7.3 ANNEX C: PQAR for product CH4_GOS_SRPR	102
7.4 ANNEX D: PQAR for products XCO2_EMMA, XCH4_EMMA, XCO2_OBS4MIPS, XCH4_OBS4MIPS	102
7.5 ANNEX E: PQAR for IASI CO ₂ and CH ₄ and AIRS CO ₂ products	102



History of modifications

Version	Date	Description of modification	Chapters / Sections
1.3	20-October-2017	New document for data set CDR1 (temporal coverage: 2003-2016)	All
2.0	16-October-2018	Update for data set CDR2 (temporal coverage: 2003-2017)	All
3.0	12-August-2019	Update for data set CDR3 (temporal coverage: 2003-2018)	All
3.1	03-November-2019	<p>Update after review by Assimila: Correction of typos and broken links.</p> <p>Other changes: Figures 35 and 36 updated and reference to Reuter et al., 2019, added. Updates of figures and tables in Sects. 3.1 and 3.2 to correct for a bug in the computation of “Uncertainty ratio”.</p>	First 7 pages (date, version number, etc.), update of Figs. 35 and 36, updates of figures and tables in Sects. 3.1 and 3.2.



Related documents

Reference ID	Document
D1	GCOS-154: Global Climate Observing System (GCOS): SYSTEMATIC OBSERVATION REQUIREMENTS FOR SATELLITE-BASED DATA PRODUCTS FOR CLIMATE - 2011 Update - Supplemental details to the satellite-based component of the “Implementation Plan for the Global Observing System for Climate in Support of the UNFCCC (2010 Update)”, December 2011, prepared by World Meteorological Organization (WMO), Intergovernmental Oceanographic Commission, United Nations Environment Programme (UNEP), International Council for Science, Doc.: GCOS 154, link: http://cci.esa.int/sites/default/files/gcos-154.pdf , 2011.
D2	GCOS-200: The Global Observing System for Climate: Implementation Needs, World Meteorological Organization (WMO), GCOS-200 (GOOS-214), pp. 325, link: http://unfccc.int/files/science/workstreams/systematic_observation/application/pdf/gcos_ip_10oct2016.pdf , 2016.
D3	ESA-CCI-GHG-URDv2.1: Chevallier, F., et al., User Requirements Document (URD), ESA Climate Change Initiative (CCI) GHG-CCI project, Version 2.1, 19 Oct 2016, link: http://www.esa-ghg-cci.org/?q=webfm_send/344 , 2016.
D4	TRD GHG, 2017: Buchwitz, M., Aben, I., Anand, J., Armante, R., Boesch, H., Crevoisier, C., Detmers, R. G., Hasekamp, O. P., Reuter, M., Schneising-Weigel, O., Target Requirement Document, Copernicus Climate Change Service (C3S) project on satellite-derived Essential Climate Variable (ECV) Greenhouse Gases (CO ₂ and CH ₄) data products (project C3S_312a_Lot6), Version 1.3, 20-October-2017, pp. 53, 2017.
D5	ATBD GHG, 2019: Buchwitz, M., Aben, I., J., Armante, R., Boesch, H., Crevoisier, C., Di Noia, A., Hasekamp, O. P., Reuter, M., Schneising-Weigel, O., Wu, L., Algorithm Theoretical Basis Document (ATBD) – Main document for Greenhouse Gas (GHG: CO ₂ & CH ₄) data set CDR 3 (2003-2018), C3S project C3S_312b_Lot2_DLR, v3.1, 2019.
D6	PUGS GHG, 2019: Buchwitz, M., Aben, I., J., Armante, R., Boesch, H., Crevoisier, C., Di Noia, A., Hasekamp, O. P., Reuter, M., Schneising-Weigel, O., Wu, L., Product User Guide and Specification (PUGS) – Main document for Greenhouse Gas (GHG: CO ₂ & CH ₄) data set CDR 3 (2003-2018), C3S project C3S_312b_Lot2_DLR, v3.1, 2019.



Acronyms

Acronym	Definition
AIRS	Atmospheric Infrared Sounder
AMSU	Advanced Microwave Sounding Unit
ATBD	Algorithm Theoretical Basis Document
BESD	Bremen optimal ESTimation DOAS
CAR	Climate Assessment Report
C3S	Copernicus Climate Change Service
CCDAS	Carbon Cycle Data Assimilation System
CCI	Climate Change Initiative
CDR	Climate Data Record
CDS	(Copernicus) Climate Data Store
CMUG	Climate Modelling User Group (of ESA's CCI)
CRG	Climate Research Group
D/B	Data base
DOAS	Differential Optical Absorption Spectroscopy
EC	European Commission
ECMWF	European Centre for Medium Range Weather Forecasting
ECV	Essential Climate Variable
EMMA	Ensemble Median Algorithm
ENVISAT	Environmental Satellite (of ESA)
EO	Earth Observation
ESA	European Space Agency
EU	European Union
EUMETSAT	European Organisation for the Exploitation of Meteorological Satellites
FCDR	Fundamental Climate Data Record
FoM	Figure of Merit
FP	Full Physics retrieval method
FTIR	Fourier Transform InfraRed
FTS	Fourier Transform Spectrometer
GCOS	Global Climate Observing System
GEO	Group on Earth Observation
GEOSS	Global Earth Observation System of Systems
GHG	GreenHouse Gas
GHG-CCI	GHG project of ESA's CCI
GOME	Global Ozone Monitoring Experiment
GMES	Global Monitoring for Environment and Security
GOSAT	Greenhouse Gases Observing Satellite
IASI	Infrared Atmospheric Sounding Interferometer



IMAP-DOAS (or IMAP)	Iterative Maximum A posteriori DOAS
IPCC	International Panel in Climate Change
IUP	Institute of Environmental Physics (IUP) of the University of Bremen, Germany
JAXA	Japan Aerospace Exploration Agency
JCGM	Joint Committee for Guides in Metrology
L1	Level 1
L2	Level 2
L3	Level 3
L4	Level 4
LMD	Laboratoire de Météorologie Dynamique
MACC	Monitoring Atmospheric Composition and Climate, EU GMES project
NA	Not applicable
NASA	National Aeronautics and Space Administration
NetCDF	Network Common Data Format
NDACC	Network for the Detection of Atmospheric Composition Change
NIES	National Institute for Environmental Studies
NIR	Near Infra Red
NLIS	LMD/CNRS <i>neuronal</i> network mid/upper tropospheric CO ₂ and CH ₄ retrieval algorithm
NOAA	National Oceanic and Atmospheric Administration
Obs4MIPs	Observations for Climate Model Intercomparisons
OCO	Orbiting Carbon Observatory
OE	Optimal Estimation
PBL	Planetary Boundary Layer
ppb	Parts per billion
ppm	Parts per million
PR	(light path) PROxy retrieval method
PUGS	Product User Guide and Specification
PVIR	Product Validation and Intercomparison Report
QA	Quality Assurance
QC	Quality Control
REQ	Requirement
RMS	Root-Mean-Square
RTM	Radiative transfer model
SCIAMACHY	SCanning Imaging Absorption spectroMeter for Atmospheric ChartographY
SCIATRAN	SCIAMACHY radiative transfer model
SRON	SRON Netherlands Institute for Space Research
SWIR	Short Wava Infra Red
TANSO	Thermal And Near infrared Sensor for carbon Observation
TANSO-FTS	Fourier Transform Spectrometer on GOSAT



TBC	To be confirmed
TBD	To be defined / to be determined
TCCON	Total Carbon Column Observing Network
TIR	Thermal Infra Red
TR	Target Requirements
TRD	Target Requirements Document
WFM-DOAS (or WFMD)	Weighting Function Modified DOAS
UoL	University of Leicester, United Kingdom
URD	User Requirements Document
WMO	World Meteorological Organization
Y2Y	Year-to-year (bias variability)



General definitions

Table 1 lists some general definitions relevant for this document. Other definitions, which require more detailed explanations, are given below.

Table 1 - General definitions.

Item	Definition
XCO ₂	Column-average dry-air mixing ratio (mole fraction) of CO ₂
XCH ₄	Column-average dry-air mixing ratio (mole fraction) of CH ₄
L1	Level 1 satellite data product: geolocated radiance (spectra)
L2	Level 2 satellite-derived data product: Here: CO ₂ and CH ₄ information for each ground-pixel
L3	Level 3 satellite-derived data product: Here: Gridded CO ₂ and CH ₄ information, e.g., 5 deg times 5 deg, monthly
L4	Level 4 satellite-derived data product: Here: Surface fluxes (emission and/or uptake) of CO ₂ and CH ₄

In the following some relevant Target Requirement (TR) related definitions are given. For details please see *TRD GHG, 2017*, *ESA-CCI-GHG-URDv2.1* and *CMUG-RBD, 2010*:

Systematic error: component of measurement error that in replicate measurements remains constant or varies in a predictable manner

Note: “Systematic error” = “Absolute systematic error” (in contrast to “Relative systematic error” defined below).

For satellite GHG ECV products especially the “Relative systematic error” is important. The definition as used here is as follows:

Relative systematic error: Identical with “Systematic error” but after bias correction and without considering a possible “global offset” (overall mean bias). Reflects the importance of spatially and temporally correlated errors (“spatio-temporal biases”). Computed from standard deviations of spatial and temporal biases.

Bias: estimate of a systematic measurement error (*JCGM, 2008*).

Precision is the measure of reproducibility or repeatability of the measurement without reference to an international standard so that precision is a measure of the random and not the systematic error. Suitable averaging of the random error can improve the precision of the measurement but does not establish the systematic error of the observation (*CMUG-RBD, 2010*).



Note: Precision (as explained in *TRD GHG, 2017*) is quantified with the standard deviation (1-sigma) of the error distribution.

Stability is a term often invoked with respect to long-term records when no absolute standard is available to quantitatively establish the systematic error - the bias defining the time-dependent (or instrument-dependent) difference between the observed quantity and the true value (*CMUG-RBD, 2010*).

Note: Stability requirements cover inter-annual error changes. If the change in the average bias from one year to another is larger than the defined values, the corresponding product does not meet the stability requirement.

Representativity is important when comparing with or assimilating in models. Measurements are typically averaged over different horizontal and vertical scales compared to model fields. If the measurements are smaller scale than the model it is important. The sampling strategy can also affect this term (*CMUG-RBD, 2010*).

Threshold requirement: The threshold is the limit at which the observation becomes ineffectual and is not of use for climate-related applications (*CMUG-RBD, 2010*).

Goal requirement: The goal is an ideal requirement above which further improvements are not necessary (*CMUG-RBD, 2010*).

Breakthrough requirement: The breakthrough is an intermediate level between the “threshold” and “goal” requirements, which - if achieved - would result in a significant improvement for the targeted application. The breakthrough level may be considered as an optimum, from a cost-benefit point of view when planning or designing observing systems (*CMUG-RBD, 2010*).

Horizontal resolution is the area over which one value of the variable is representative of (*CMUG-RBD, 2010*).

Vertical resolution is the height over which one value of the variable is representative of. Only used for profile data (*CMUG-RBD, 2010*).

Observing Cycle is the temporal frequency at which the measurements are required (*CMUG-RBD, 2010*).

Note: In this document also the term “Revisit time” may be used. The definition is identical with the definition of “Observing cycle”. Both terms refer to the (average) temporal frequency at a given location.



Scope of document

This document is the Product Quality Assurance Report (PQAR) for the Copernicus Climate Change Service (C3S, <https://climate.copernicus.eu/>) component as covered by the greenhouse gas (GHG) sub-project of project C3S_312b_Lot2 led by DLR, Germany (a follow-on activity of project C3S_312a_Lot6 led by University of Bremen, Germany), in the following referred to as C3S/GHG project.

Within this project satellite-derived atmospheric carbon dioxide (CO₂) and methane (CH₄) Essential Climate Variable (ECV) data products have been generated and provided to ECMWF for inclusion into the Copernicus Climate Data Store (CDS) from which users can access these data products and the corresponding documentation.

The satellite-derived data products described and quality assessed in this document are:

- Column-average dry-air mixing ratios (mole fractions) of CO₂ and CH₄, denoted XCO₂ (in parts per million, ppm) and XCH₄ (in parts per billion, ppb), respectively.
- Mid/upper tropospheric mixing ratios of CO₂ (in ppm) and CH₄ (in ppb).

An overview about the products is given in Table 2 for the CO₂ products and in Table 3 for the CH₄ products.

For an overview of the merged Level 2 data products XCO₂_EMMA and XCH₄_EMMA and of the merged Level 3 data products XCO₂_OBS4MIPS and XCH₄_OBS4MIPS see also *Reuter et al., 2019*.

Requirements on data quality are formulated in the corresponding Target Requirement Document (TRD) (*TRD GHG, 2017*).

The main purpose of this document is to describe the validation / quality assessment of the satellite-derived CO₂ and CH₄ greenhouse gas (GHG) ECV data products.



Table 2: Overview CO₂ products. “CRD#” indicates the Climate Data Record Number. CRD1 has been released in 2017, CDR2 in 2018 and CDR3 will be released in 2019. Level 2 (L2) products contains information for each individual satellite footprint (ground pixel) whereas Level 3 (L3) products are gridded /averaged spatially and temporally. If CDR# is 2-3 then this means that the CDR3 product is the same as the CDR2 product (no update).

Product ID (Level)	Version	CDR#	Temporal coverage	Comments
CO2_SCI_BESD (L2)	02.01.02	1-3	01.2003 – 03.2012	XCO ₂ from SCIAMACHY as retrieved with Univ. Bremen’s BESD algorithm. Brokered from GHG-CCI.
CO2_SCI_WFMD (L2)	4.0	1-3	10.2002 – 04.2012	XCO ₂ from SCIAMACHY as retrieved with Univ. Bremen’s WFMD algorithm. Brokered from GHG-CCI.
CO2_GOS_OCFP (L2)	7.1 7.2	1 2-3	04.2009 – 12.2016 04.2009 – 12.2018	XCO ₂ from GOSAT as retrieved with Univ. Leicester’s OCFP algorithm.
CO2_GOS_SRF (L2)	2.3.8	1-3	04.2009 – 12.2018	XCO ₂ from GOSAT as retrieved with SRON’s SRF (RemoTeC) algorithm.
XCO2_EMMA (L2)	3.0 3.1 4.1	1 2 3	01.2003 – 12.2016 01.2003 – 12.2017 01.2003 – 12.2018	Merged L2 XCO ₂ product using Univ. Bremen’s EMMA algorithm.
XCO2_OBS4MIPS (L3)	3 3.1 4.1	1 2 3	01.2003 – 12.2016 01.2003 – 12.2017 01.2003 – 12.2018	Merged L3 XCO ₂ product in OBS4MIPS format.
CO2_AIRS_NLIS (L2)	3.0	1-3	04.2003 – 07.2007	Mid-tropospheric CO ₂ mixing ratios as retrieved from AIRS using LMD’s NLIS algorithm. Brokered from GHG-CCI.
CO2_IASA_NLIS (L2)	8.0	1-3	7.2007 – 05.2015	Mid-tropospheric CO ₂ mixing ratios as retrieved from IASI/Metop-A using LMD’s NLIS algorithm.
CO2_IASB_NLIS (L2)	4.0 4.2	1 2-3	2.2013 – 12.2016 2.2013 – 12.2018	Mid-tropospheric CO ₂ mixing ratios as retrieved from IASI/Metop-B using LMD’s NLIS algorithm.



Table 3: Overview CH₄ products. “CRD#” indicates the Climate Data Record Number. CRD1 has been released in 2017, CDR2 in 2018 and CDR3 will be released in 2019. Level 2 (L2) products contains information for each individual satellite footprint (ground pixel) whereas Level 3 (L3) products are gridded /averaged spatially and temporally. If CDR# is 2-3 then this means that the CDR3 product is the same as the CDR2 product (no update).

Product ID (Level)	Version	CDR#	Temporal coverage	Comments
CH4_SCI_WFMD (L2)	4.0	1-3	10.2002 – 12.2011	XCH ₄ from SCIAMACHY as retrieved with Univ. Bremen’s WFMD algorithm. Brokered from GHG-CCI.
CH4_SCI_IMAP (L2)	7.2	1-3	01.2003 – 04.2012	XCH ₄ from SCIAMACHY as retrieved with SRON/JPL’s IMAP algorithm. Brokered from GHG-CCI.
CH4_GOS_OCPR (L2)	7.0 7.2	1 2-3	04.2009 – 12.2016 04.2009 – 12.2018	XCH ₄ from GOSAT as retrieved with Univ. Leicester’s OCPR algorithm.
CH4_GOS_SRPR (L2)	2.3.8 2.3.9	1 2-3	04.2009 – 12.2016 04.2009 – 12.2018	XCH ₄ from GOSAT as retrieved with SRON’s SRPR (RemoTeC) algorithm.
CH4_GOS_OCFP (L2)	7.1 7.2	1 2-3	04.2009 – 12.2016 04.2009 – 12.2018	XCH ₄ from GOSAT as retrieved with Univ. Leicester’s OCFP algorithm.
CH4_GOS_SRFP (L2)	2.3.8 2.3.8	1 2-3	04.2009 – 12.2016 04.2009 – 12.2018	XCH ₄ from GOSAT as retrieved with SRON’s SRFP (RemoTeC) algorithm.
XCH4_EMMA (L2)	3.0 3.1 4.1	1 2 3	01.2003 – 12.2016 01.2003 – 12.2017 01.2003 – 12.2018	Merged L2 XCH ₄ product using Univ. Bremen’s EMMA algorithm.
XCH4_OBS4MIPS (L3)	3 3.1 4.1	1 2 3	01.2003 – 12.2016 01.2003 – 12.2017 01.2003 – 12.2018	Merged L3 XCH ₄ product in OBS4MIPS format.
CH4_IASA_NLIS (L2)	8.4	1-3	7.2007 – 05.2015	Mid-tropospheric CH ₄ mixing ratios as retrieved from IASI/Metop-A using LMD’s NLIS algorithm.
CH4_IASB_NLIS (L2)	8.1 8.1	1 2-3	2.2013 – 12.2016 2.2013 – 12.2018	Mid-tropospheric CH ₄ mixing ratios as retrieved from IASI/Metop-B using LMD’s NLIS algorithm.



Executive summary

In this document the validation / quality assessment of satellite-derived atmospheric carbon dioxide (CO₂) and methane (CH₄) Climate Data Record (CDR) data products as generated via the C3S_312b_Lot2 project of the Copernicus Climate Change Service (C3S, <https://climate.copernicus.eu/>) is described.

The C3S_312b_Lot2 satellite-derived greenhouse gas (GHG) data products are:

- Column-average dry-air mixing ratios (mole fractions) of CO₂ and CH₄, denoted XCO₂ (in parts per million, ppm) and XCH₄ (in parts per billion, ppb), respectively.
- Mid/upper tropospheric mixing ratios of CO₂ (in ppm) and CH₄ (in ppb).

These data products are generated from the satellite instruments SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT (XCO₂ and XCH₄ products) and AIRS and IASI (mid/upper troposphere products). All data products are available as Level 2 (individual ground pixels) products in NetCDF format. The XCO₂ and XCH₄ Level 2 products correspond to individual satellite sensors but are also available as merged multi-sensor products. In addition, also merged Level 3 (i.e., gridded) products in Obs4MIPs format are available for the XCO₂ and XCH₄ products. For details on data format etc. please see the Product User Guide and Specification (PUGS) document (*PUGS GHG, 2018*).

CO₂ and CH₄ are important climate-relevant atmospheric gases, so-called greenhouse gases (GHG). Because of their important role for climate they are classified as Essential Climate Variables (ECVs). The ECV GHG as formulated by GCOS (Global Climate Observing System) is defined as: “Retrievals of greenhouse gases, such as CO₂ and CH₄, of sufficient quality to estimate regional sources and sinks” (GCOS-154). This definition contains already the main application of these atmospheric data products, namely to use them (in combination with appropriate (inverse) modelling) to obtain (improved) information on their (primarily surface) sources and sinks.

Both gases, CO₂ and CH₄, have a long lifetime in the atmosphere. As a consequence of this fact and related human emissions the atmospheric concentrations of these gases are relatively high (currently about 400 ppm for CO₂ and 1.8 ppm (1800 ppb) for CH₄) compared to other atmospheric trace gases. As a result of this, even a moderate to strong (surface) source or sink typically only results in a relatively small local or regional change (enhancement or depletion relative to the surrounding region) in their vertical columns or their mid/upper tropospheric concentration. The observational requirements are therefore very demanding in particular with respect to random and systematic errors and stability.

Because of their long lifetime and atmospheric transport, elevated (or depleted) atmospheric CO₂ and CH₄ concentrations can be higher (or lower) relative to the background far away from the surface source (or sink), which has emitted (or taken up) these atmospheric gases. In order to obtain source/sink information from the atmospheric observations it is therefore required to take atmospheric transport (and in particular for methane also atmospheric chemistry) into account and to consider the exact time and location of the atmospheric observations. As a consequence, the most relevant data products are the Level 2 (L2) products, which contain detailed information (time,



location, etc.) for each individual satellite ground pixel. The requirements as formulated in the Target Requirement Document (*TRD GHG, 2017*) are, therefore, mostly L2 requirements. However, for XCO₂ and XCH₄ also (gridded) Level 3 (L3) products have been generated (in Obs4MIPs format) and also their validation is described in this document.

The C3S_312a_Lot6 project is essentially the (pre-)operational continuation of the research and development (R&D) pre-cursor project GHG-CCI (<http://www.esa-ghg-cci.org/>) of ESA's Climate Change Initiative (CCI). The main goal of the C3S_312a_Lot6 project is to extend (in time) the data base of GHG-CCI pre-cursor data products.

The first C3S_312a_Lot6 GHG data set - Climate Data Record 1 (CDR1) - covered the time period 2003-2016 and had been delivered to ECMWF in 2017. CDR2 covers the time period 2003-2017.

The latest data set - Climate Data Record 2 (CDR3) - covers the time period 2003-2018. This document is an update for data set CDR3.

This document is the MAIN PQAR document. It provides an overview of all products including validation / quality assessment results (including the latest versions of SCIAMACHY XCO₂ and XCH₄ products as generated in the framework of the GHG-CCI project). Additional detailed validation results for each product are provided in separate ANNEXes:

- ANNEX A: PQAR for products CO₂_GOS_OCFP, CH₄_GOS_OCFP, CH₄_OCPR (University of Leicester's GOSAT products)
- ANNEX B: PQAR for products CO₂_GOS_SRFp, CH₄_GOS_SRFp (SRON's "full physics" GOSAT products)
- ANNEX C: PQAR for product CH₄_GOS_SRPR (SRON's "proxy" GOSAT XCH₄ product)
- ANNEX D: PQAR for products XCO₂_EMMA, XCH₄_EMMA, XCO₂_OBS4MIPS, XCH₄_OBS4MIPS (University of Bremen's merged Level 2 products)
- ANNEX E: PQAR for IASI CO₂ and CH₄ products and AIRS CO₂ product (LMD/CNRS's IASI and AIRS products)

Table 4 provides an overview about all products and their estimated data quality in terms of Target Requirement (TR) assessments.

Figure 1 presents an overview of the achieved data quality for all Level 2 XCO₂ data products and Figure 2 presents this overview for the Level 2 XCH₄ data products.



Table 4 - Overview quality assessment results of products in terms of Target Requirements (TRs). For additional quality assessment results see the following two figures.

Product ID	Level	Description	Probability that TR is met		Details see Sect.
			Accuracy	Stability	
XCO ₂ products			Required: < 0.5 ppm	Required: < 0.5 ppm/year	
CO2_SCI_BESD	2	XCO ₂ from SCIAMACHY retrieved using Univ. Bremen’s BESD algorithm	64%	87%	3.1.1
CO2_SCI_WFMD	2	XCO ₂ from SCIAMACHY retrieved using Univ. Bremen’s WFMD algorithm	41%	95%	3.1.2
CO2_GOS_OCFP	2	XCO ₂ from GOSAT retrieved using Univ. Leicester’s OCFP algorithm	51%	96%	3.1.3
CO2_GOS_SRFP	2	XCO ₂ from GOSAT retrieved using SRON’s SRFP (RemoTeC) algorithm	36%	95%	3.1.4
XCO2_EMMA	2	Merged multi-satellite XCO ₂ via Univ. Bremen’s EMMA algorithm	49%	97%	3.1.7
XCO2_OBS4MIPS	3	Merged multi-satellite XCO ₂ via Univ. Bremen’s OBS4MIPS algorithm	47%	98%	3.3
XCH ₄ products			Required: < 10 ppb	Required: < 3 ppb/year	
CH4_SCI_WFMD	2	XCH ₄ from SCIAMACHY retrieved using Univ. Bremen’s WFMD algorithm	46%	85%	3.2.1
CH4_SCI_IMAP	2	XCH ₄ from SCIAMACHY retrieved using the IMAP algorithm of SRON/JPL	61%	98%	3.2.2
CH4_GOS_OCPR	2	XCH ₄ from GOSAT retrieved using Univ. Leicester’s OCPR algorithm	89%	97%	3.2.3



CH4_GOS_SRPR	2	XCH ₄ from GOSAT retrieved using SRON's SRPR (RemoTeC) algorithm	89%	97%	3.2.4
CH4_GOS_OCFP	2	XCH ₄ from GOSAT retrieved using Univ. Leicester's OCFP algorithm	87%	95%	3.2.5
CH4_GOS_SRFP	2	XCH ₄ from GOSAT retrieved using SRON's SRFP (RemoTeC) algorithm	87%	99%	3.2.6
XCH4_EMMA	2	Merged multi-satellite XCH ₄ via Univ. Bremen's EMMA algorithm	88%	98%	3.2.7
XCH4_OBS4MIPS	3	Merged multi-satellite XCH ₄ via Univ. Bremen's OBS4MIPS algorithm	89%	100%	3.4
Mid/upper troposphere CO ₂ products			Required: < 0.5 ppm	Required: < 0.5 ppm/year	
CO2_AIR_NLIS	2	LMD's product from AIRS	-	-	3.5
COS_IASA_NLIS	2	LMD's product from IASI/Metop-A	70%	100%	3.5
CO2_IASB_NLIS	2	LMD's product from IASI/Metop-B	-	-	3.5
Mid/upper troposphere CH ₄ products			Required: < 10 ppb	Required: < 3 ppb/year	
CH4_IASA_NLIS	2	LMD's product from IASI/Metop-A	90%	-	3.5
CH4_IASB_NLIS	2	LMD's product from IASI/Metop-B	-	-	3.5




Figure 1 - Overview data quality assessment results for Level 2 XCO₂ data products. The green bars refer to the “Quality Assessment / Quality control” (QA/QC) results as described in this document. The red bars refer to results obtained by the data providers (DPs), as described in separate Annexes (see Sect. 7). The blue bars result from an assessment using the EMMA method (see Sect. 3.1.5). For “Accuracy” and “Stability” also the numerical values for the “Probability that TR is met” are given (computed as mean value if more than one value (bar) exists). Also listed (in grey on the right hand side) is the uncertainty of the reference data as used for the Target Requirements (TR) assessments. The DP values for the “GHG-CCI brokered products” CO₂_SCI_BESD and CO₂_SCI_WFM are from *Buchwitz et al., 2017*.

C3S Level 2 products: XCO₂

QA/QC method (applied to all products)

DP methods (applied by data provider)

EMMA method



Michael.Buchwitz@iup.physik.uni-bremen.de, 12-Aug-2019




Figure 2 - Overview data quality assessment results for Level 2 XCH₄ data products. The green bars refer to the “Quality Assessment / Quality control” (QA/QC) results as described in this document. The red bars refer to results obtained by the data providers (DPs), as described in separate Annexes (see Sect. 7). The blue bars result from an assessment using the EMMA method (see Sect. 3.2.7). For “Accuracy” and “Stability” also the numerical values for the “Probability that TR is met” are given (computed as mean value if more than one value (bar) exists). Also listed (in grey on the right hand side) is the uncertainty of the reference data as used for the Target Requirements (TR) assessments. The DP values for the “GHG-CCI brokered products” CH4_SCI_WFMD and CH4_SCI_IMAP are from *Buchwitz et al., 2017*.

C3S Level 2 products: XCH₄

QA/QC method (applied to all products)

DP methods (applied by data provider)

EMMA method



Michael.Buchwitz@iup.physik.uni-bremen.de, 12-Aug-2019



1. Overview data products and instruments

In this section an overview of the data products - specified in terms of variable, its property, processing level(s) and instrument(s) - is given.

The data products are (see also *Buchwitz et al., 2013b, 2016, 2017*):

- Column-average dry-air mixing ratios (mole fractions) of CO₂ and CH₄, denoted XCO₂ (in parts per million, ppm) and XCH₄ (in parts per billion, ppb).
- Mid/upper tropospheric mixing ratios of CO₂ and CH₄.

Carbon dioxide and methane are important atmospheric greenhouse gases (e.g., *IPCC 2013*) but despite their importance our knowledge on their various and variable natural and anthropogenic sources and sinks has significant gaps (e.g., *IPCC 2013; Ciais et al., 2014; 2015; Kirschke et al., 2013; Nisbet et al., 2014*, and references given therein). A purpose of the satellite data products described in this document is to contribute to enhancing our knowledge on the CO₂ and CH₄ sources and sinks (via appropriate (inverse) modelling).

Carbon dioxide and methane are so-called Essential Climate Variables (ECVs) and the need to monitor them has been clearly identified along with the definition of key observational requirements (e.g., *GCOS-154, GCOS-200*). In recent years several satellite-derived ECV data products have been generated in particular in the framework of the Climate Change Initiative (CCI) of ESA (e.g., *Hollmann et al., 2013*) including CO₂ and CH₄ (e.g., *Buchwitz et al., 2013a, 2016, 2017*).

Previous version of these satellite-derived CO₂ and CH₄ data products have been used for a number of (primarily scientific) applications, e.g.,

- to improve our knowledge on the various natural and anthropogenic (surface) sources and sinks of these important greenhouse gases (GHG) (see, e.g., *Alexe et al., 2015; Bergamaschi et al., 2015; Chevallier et al., 2014, 2016a, 2016b; Cressot et al., 2014; Detmers et al., 2015; Guerlet et al., 2013; Houweling et al., 2015; McNorton et al., 2016; Pandey et al., 2016; Reuter et al., 2014b, 2017; Schneising et al., 2014b; Turner et al., 2015, 2016*, and references given therein)
- to monitor the global distribution of CO₂ and CH₄ (e.g., *Buchwitz et al., 2007, 2016b; Schneising et al., 2011; Frankenberg et al., 2011; Massart et al., 2016*)
- to improve our knowledge on emission ratios, e.g., for biomass burning (e.g., *Ross et al., 2013; Parker et al., 2016*)
- for comparisons with (chemistry) climate models (e.g., *Shindell et al., 2013; Hayman et al., 2014; Lauer et al., 2017*) and other models (e.g., *Schneising et al., 2014a; Parker et al., 2016*)

In the following sub-sections an overview about the satellite-derived CO₂ and CH₄ data products is given.



1.1 Column-average mixing ratios of CO₂ and CH₄ (XCO₂ and XCH₄)

1.1.1 Overview

Satellite radiance observations in the Near Infrared / Short Wave Infrared (NIR/SWIR) spectral region in nadir (downlooking) observation viewing mode are sensitive to atmospheric CO₂ and CH₄ concentration changes with good sensitivity down to the Earth's surface (because solar radiation reflected at the Earth's surface is observed). These measurements permit to obtain "total column information" but do not permit to obtain (detailed) information on the vertical profiles of CO₂ and CH₄. The CO₂ and CH₄ products derived from these satellites are column-averaged dry-air mixing ratios (more precisely: mole fractions) of CO₂ and CH₄ denoted XCO₂ (e.g., in ppm) and XCH₄ (e.g., in ppb).

In the following, several satellite instruments are shortly described which have been used (or will be used in the future) to generate XCO₂ and/or XCH₄ data products.

1.1.2 Instruments

In this section a short overview about relevant satellite instruments is given. The C3S data set has been primarily derived from the satellite instruments SCIAMACHY on ENVISAT and TANSO-FTS onboard GOSAT. In addition, XCO₂ from NASA's OCO-2 mission has been used for some products (EMMA and OBS4MIPS). These instruments are shortly described in the following. Other satellites are also shortly mentioned, which are planned to be used for future versions of our data products.

1.1.2.1 SCIAMACHY/ENVISAT

SCIAMACHY (SCanning Imaging Absorption spectroMeter for Atmospheric ChartographY) was a spectrometer on ESA's ENVISAT satellite (2002-2012). SCIAMACHY (*Burrows et al., 2005; Bovensmann et al., 1999*) covers the spectral region from the ultra-violet to the SWIR spectral region (240 nm - 2380 nm) at moderate spectral resolution (0.2 nm - 1.5 nm) and observes the Earth's atmosphere in various viewing geometries (nadir, limb and solar and lunar occultation). For a good general overview on SCIAMACHY see also <https://en.wikipedia.org/wiki/SCIAMACHY>. SCIAMACHY permits the retrieval of XCO₂ (e.g., *Reuter et al., 2011; Schneising et al., 2011*) and XCH₄ (e.g., *Schneising et al., 2011; Frankenberg et al., 2011*) from the appropriate spectral regions in the SWIR (around 1.6 µm) and the NIR (O₂ A-band at 760 nm used to obtain the dry-air column using the known dry-air mixing ratio of atmospheric oxygen). The ground pixel size is typically 30 km along track times 60 km across track and the swath width is about 960 km. There are no across-track gaps between the ground pixels but there are gaps along-track as SCIAMACHY operates only part of the time (approx. 50%) in nadir observation mode.



1.1.2.2 TANSO-FTS/GOSAT

TANSO-FTS is a Fourier-Transform-Spectrometer (FTS) onboard the Japanese GOSAT satellite (Kuze *et al.*, 2009, 2014, 2016). The Greenhouse Gases Observing Satellite "IBUKI" (GOSAT) is the world's first spacecraft in orbit dedicated to measure the concentrations of carbon dioxide and methane from space. The spacecraft was launched successfully on January 23, 2009, and has been operating properly since then. GOSAT covers the relevant CO₂, CH₄ and O₂ absorption bands in the NIR and SWIR spectral region as needed for accurate XCO₂ and XCH₄ retrieval (in addition GOSAT also covers a large part of the Thermal Infrared (TIR) spectral region). The spectral resolution of TANSO-FTS is much higher compared to SCIAMACHY and also the ground pixels are smaller (10 km compared to several 10 km for SCIAMACHY). However, in contrast to SCIAMACHY, the GOSAT scan pattern consists of non-consecutive individual ground pixels, i.e., the scan pattern is not gap-free. For a good general overview about GOSAT see also <http://www.gosat.nies.go.jp/en/>.

GOSAT-2 has been successfully launched on 29 October 2018. OSAT-2 XCO₂ and XCH₄ retrievals are not yet included in the C3S GHG CDR.

1.1.2.3 OCO-2

NASA's Orbiting Carbon Observatory 2 (OCO-2) mission (Crisp *et al.*, 2004; Boesch *et al.*, 2011) has been successfully launched in July 2014. The OCO-2 Project primary science objective is to collect the first space-based measurements of atmospheric carbon dioxide with the precision, resolution and coverage needed to characterize its sources and sinks and quantify their variability over the seasonal cycle. During its two-year mission, OCO-2 will fly in a sun-synchronous, near-polar orbit with a group of Earth-orbiting satellites with synergistic science objectives whose ascending node crosses the equator near 13:30 hours Mean Local Time (MLT). Near-global coverage of the sunlit portion of Earth is provided in this orbit over a 16-day (233-revolution) repeat cycle. OCO-2's single instrument incorporates three high-resolution grating spectrometers, designed to measure the near-infrared absorption of reflected sunlight by carbon dioxide and molecular oxygen. OCO-2 covers similar spectral bands as SCIAMACHY and GOSAT but OCO-2 has much smaller ground pixels (km scale) but the swath width is much smaller (approx. 10 km) compared to SCIAMACHY. OCO-2 delivers XCO₂ but not XCH₄. Details on OCO-2 are also given on <https://oco.jpl.nasa.gov/>.

1.1.2.4 TanSat

The Chinese TanSat satellite (<https://en.wikipedia.org/wiki/TanSat>) has been successfully launched in December 2016. The TanSat satellite and instrument is very similar as OCO-2. As OCO-2, TanSat delivers XCO₂ but not XCH₄. TanSat XCO₂ retrievals are not yet included in the C3S GHG CDR.



1.1.2.5 Sentinel-5-Precursor (S5P)

ESA's Sentinel-5-Precursor (S5P) mission (*Veeffkind et al., 2012*) has been launched in October 2017. S5P permits XCH₄ retrievals (*Butz et al., 2012, Hu et al., 2018*) at about 7 km and using a wide swath of about 2600 km. Details on S5P can also be found on <https://earth.esa.int/web/guest/missions/esa-future-missions/sentinel-5P>. S5P XCH₄ retrievals are not yet included in the C3S GHG CDR.

1.1.2.6 Other instruments

Several other satellites are expected to be launched in the future, e.g., the active laser-based mission MERLIN (Methane Remote Sensing Lidar Mission, see [https://de.wikipedia.org/wiki/Merlin_\(Satellit\)](https://de.wikipedia.org/wiki/Merlin_(Satellit))).



1.1.3 XCO₂


As explained, XCO₂ is the column-averaged dry-air mixing ratio (mole fraction) of atmospheric CO₂. A XCO₂ value of, for example, 400 ppm at a given location means that 400 CO₂ molecules are present in the atmosphere above that location per one million air molecules excluding water molecules.

XCO₂ can be retrieved from instruments such as SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT using Optimal Estimation (Rodgers, 2000) or DOAS (Buchwitz et al., 2000) retrieval algorithms as shown in various publications (e.g., Buchwitz et al., 2005; Butz et al., 2011; Cogan et al., 2011; Reuter et al., 2011; 2013; Schneising et al., 2011; Yoshida et al., 2013). These products have been validated using Total Carbon Column Observing Network (TCCON) (Wunch et al., 2010, 2011, 2015) XCO₂ ground based observations (e.g., Dils et al., 2014).

In this document we describe the latest versions of these data products.

As an example, Figure 3 shows time series of satellite-derived XCO₂. As can be seen, XCO₂ is increasing by about 2 ppm/year primarily due to burning of fossil fuels and shows a pronounced seasonal cycle, primarily due to uptake and release of CO₂ by the terrestrial biosphere.

Figure 3 – Satellite-derived northern mid-latitude XCO₂ time series. Shown are four time series, each corresponding to one of the four individual satellite sensor Level 2 XCO₂ products, which are described in this document.



Michael.Buchwitz@up.physik.uni-bremen.de 16-July-2019



1.1.4 XCH₄


As explained, XCH₄ is the column-averaged dry-air mixing ratio (mole fraction) of atmospheric CH₄. A XCH₄ value of, for example, 1800 ppb at a given location means that 1800 CH₄ molecules are present in the atmosphere above that location per one billion air molecules excluding water molecules.

XCH₄ can be retrieved from instruments such as SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT using Optimal Estimation (Rodgers, 2000) or DOAS (Buchwitz *et al.*, 2000) retrieval algorithms as shown in various publications (e.g., Buchwitz *et al.*, 2005; Butz *et al.*, 2011; Frankenberg *et al.*, 2011; Schneising *et al.*, 2011; Parker *et al.*, 2011; Scheper *et al.*, 2012; Yoshida *et al.*, 2013). These products have been validated using Total Carbon Column Observing Network (TCCON) (Wunch *et al.*, 2010, 2011, 2015) XCH₄ ground based observations (e.g., Dils *et al.*, 2014).

In this document we describe the latest versions of these data products.

As an example, Figure 4 shows time series of satellite-derived XCH₄. As can be seen, XCH₄ is increasing since 2007 by about 7 ppb/year. The reason for this is not entirely clear (several potential reasons are discussed in the scientific literature).

Figure 4 – Satellite-derived northern mid-latitude XCH₄ time series. Shown are six time series, each corresponding to one of the four individual satellite sensor Level 2 XCH₄ products, which are described in this document.





1.1.5 List of XCO₂ and XCH₄ data products

Table 5 and Table 6 list the XCO₂ and XCH₄ data products, respectively.

As can be seen from Table 5, for each individual sensor Level 2 XCO₂ product two products have been generated using two different retrieval algorithms (OCFP is University of Leicester's Full Physics (FP) algorithm and SRFP is SRON's retrieval algorithm, also known as RemoTeC).

Products with comment « Existing GHG-CCI product » are the latest versions of Level 2 products, which have been generated in the framework of the GHG-CCI project (<http://www.esa-ghg-cci.org/>). They are available via the C3S CDS but are also available from the GHG-CCI website (<http://www.esa-ghg-cci.org/>) including documentation. They have been used within project C3S_312a_Lot6 to generate the merged Level 2 and Level 3 EMMA and OBS4MIPS products but the individual sensor L2 products have not been regenerated. They have been provided for C3S « as is » and are available via the C3S CDS.

Table 5 - Overview XCO₂ data products.

Product ID	Level	Sensor(s)	(Planned) Availability	Comments
CO2_GOS_OCFP	2	GOSAT	Oct. 2017: 2009-2016 Oct. 2018: 2009-2017 Dec. 2019: 2009-2018	
CO2_GOS_SRFP	2	GOSAT	Oct. 2017: 2009-2016 Oct. 2018: 2009-2017 Dec. 2019: 2009-2018	
CO2_SCI_BESD	2	SCIAMACHY	Oct. 2017: 2003-2012	Existing GHG-CCI product
CO2_SCI_WFMD	2	SCIAMACHY	Oct. 2017: 2002-2012	Existing GHG-CCI product
XCO2_EMMA	2	Merged SCIAMACHY, GOSAT, OCO-2	Oct. 2017: 2003-2016 Oct. 2018: 2003-2017 Dec. 2019: 2003-2018	
XCO2_OBS4MIPS	3	Merged SCIAMACHY & GOSAT	Oct. 2017: 2003-2016 Oct. 2018: 2003-2017 Dec. 2019: 2003-2018	



As can be seen from Table 6, for each individual sensor Level 2 XCH₄ product four products will be generated from GOSAT using four different retrieval algorithms using two « Full Physics » (FP) and two « Proxy » (PR) algorithms. For a discussion of FP versus PR algorithms see also, for example, *Schepers et al., 2012*. Each type of algorithm has different advantages and disadvantages. Typically, the PR products contain more data and therefore somewhat better spatio-temporal coverage (as quality filtering can be less strict) but the PR algorithms rely on a CO₂ model to correct for XCO₂ variations. FP products contain less data points but the advantage of this product is that it is independent of a CO₂ model.

Table 6 - Overview XCH₄ data products.

Product ID	Level	Sensor(s)	(Planned) Availability	Comments
CH4_GOS_OCPR	2	GOSAT	Oct. 2017: 2009-2016 Oct. 2018: 2009-2017 Dec. 2019: 2009-2018	
CH4_GOS_SRPR	2	GOSAT	Oct. 2017: 2009-2016 Oct. 2018: 2009-2017 Dec. 2019: 2009-2018	
CH4_GOS_OCFP	2	GOSAT	Oct. 2017: 2009-2016 Oct. 2018: 2009-2017 Dec. 2019: 2009-2018	
CH4_GOS_SRFP	2	GOSAT	Oct. 2017: 2009-2016 Oct. 2018: 2009-2017 Dec. 2019: 2009-2018	
CH4_SCI_WFMD	2	SCIAMACHY	Oct. 2017: 2002-2011	Existing GHG-CCI product
CH4_SCI_IMAP	2	SCIAMACHY	Oct. 2017: 2003-2012	Existing GHG-CCI product
XCH4_EMMA	2	Merged SCIAMACHY & GOSAT	Oct. 2017: 2003-2016 Oct. 2018: 2003-2017 Dec. 2019: 2003-2018	
XCH4_OBS4MIPS	3	Merged SCIAMACHY & GOSAT	Oct. 2017: 2003-2016 Oct. 2018: 2003-2017 Dec. 2019: 2003-2018	



1.2 Mid-tropospheric mixing ratios of CO₂ and CH₄

1.2.1 Overview

Satellite radiance observations in the thermal infrared (TIR) spectral region in nadir (downlooking) observation viewing mode are sensitive to atmospheric CO₂ and CH₄ mixing ratio changes in the mid and upper tropospheric region. They can thus be interpreted in terms of integrated mid-tropospheric columns, with typical sensitivity between 5 and 12 km.

In the following, the 2 hyperspectral infrared sounders AIRS and IASI are shortly described.

1.2.2 Instruments

1.2.2.1 AIRS

The Atmospheric Infrared Sounder (AIRS) is a polar orbiting nadir-viewing high-resolution infrared sounder operating in a cross-track-scanning mode. It was launched onboard the EOS Aqua satellite in May 2002, with two operational microwave sounders, AMSU and HSB, and is operational since September 2002. It is a high-spectral resolution, grating multispectral infrared sounder with 2378 channels. Its spectral domain ranges from 650 cm⁻¹ to 2665 cm⁻¹ (15.4 μm and 3.8 μm), with a spectral resolving power of 1200 (i.e., a spectral resolution ranging from 0.5 cm⁻¹ to 2 cm⁻¹). This domain is divided into three spectral bands, from 650 to 1135 cm⁻¹, from 1215 to 1615 cm⁻¹ and from 2180 to 2665 cm⁻¹. AIRS cross-track scanning is 1650 km and covers 70% of the earth every day. The instantaneous field of view (IFOV) is sampled by 3×3 circular pixels whose ground resolution is 13 km at nadir. Measurements from the three instruments are analyzed jointly to filter out the effects of clouds from the IR data in order to derive clear-column air-temperature profiles and surface temperatures with high vertical resolution and accuracy (1 K per 1 km layer in the troposphere).

1.2.2.2 IASI

The Infrared Atmospheric Sounding Interferometer (IASI) is a high resolution Fourier Transform Spectrometer based on a Michelson Interferometer coupled to an integrated imaging system that measures infrared radiation emitted from the Earth. Developed by the Center National d'Etudes Spatiales (CNES) in collaboration with the European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT), IASI was launched in October 2006 onboard the polar orbiting Meteorological Operational Platform (Metop-A), and in September 2012 onboard Metop-B. A third IASI will be launched onboard Metop-C. IASI provides 8461 spectral samples, ranging from 645 cm⁻¹ to 2760 cm⁻¹ (15.5 μm and 3.6 μm), with a spectral sampling of 0.25 cm⁻¹, and a spectral resolution of 0.5 cm⁻¹ after apodisation ('Level 1c' spectra). IASI is an across track scanning system,



whose swath width is of 2200 km, allowing global coverage twice a day. The IFOV is sampled by 2×2 circular pixels whose ground resolution is 12 km at nadir. IASI has demonstrated the possibility to retrieve or detect several chemistry and climate variables from hyperspectral infrared observation: for instance water vapour (H₂O), carbon dioxide (CO₂), carbon monoxide (CO), methane (CH₄), ozone (O₃), sulfur dioxide (SO₂), hydrogen sulfide (H₂S), ammonia (NH₃), nitric acid (HNO₃), volatile organic compounds (VOCs) and aerosols (*Hilton et al., 2012; Clarisse et al., 2011*) on regional and global scales. IASI enables the monitoring of key gases for climate and atmospheric chemistry in near real time and has also highlighted the benefit of high-performance infrared sounders for numerical weather prevision (NWP) applications.

1.2.3 CO₂

Mid-tropospheric columns of CO₂ can be retrieved from hyperspectral infrared sounders such as AIRS and IASI (*Chédin et al., 2003; Crevoisier et al., 2003*) using non-linear inference scheme (*Crevoisier et al., 2009a*).

Products have been validated using aircraft measurements, mostly from the Comprehensive Observation Network for TRace gases by AirLiner (CONTRAIL) program (*Machida et al., 2008; Matsueda et al. 2008*).

1.2.4 CH₄

Mid-tropospheric columns of CH₄ can be retrieved from the hyperspectral infrared sounder IASI (*Crevoisier et al., 2003, 2013*) using non-linear inference scheme (*Crevoisier et al., 2009b*).

Products have been validated using aircraft measurements, from the Comprehensive Observation Network for TRace gases by AirLiner (CONTRAIL) program (*Machida et al., 2008; Matsueda et al. 2008*) and the HIAPER Pole-to-Pole Observations (HIPPO) project (*Wofsy et al., 2012*), as well as from balloon measurements from AirCores (*Membrane et al., 2016*).



1.2.5 List of mid-tropospheric CO₂ and CH₄ data products

Table 7 lists the CO₂ and CH₄ mid/upper troposphere data products.

The product with comment « Existing GHG-CCI product » is the latest versions of AIRS CO₂ Level 2 products, which has been generated in the framework of the GHG-CCI project (<http://www.esa-ghg-cci.org/>). This product exists and is available from the GHG-CCI website (<http://www.esa-ghg-cci.org/> -> CRDP (Data)). It has been provided for C3S essentially « as is » but converted (from ASCII) to NetCDF format (all products listed in Table 7 are available in NetCDF format).

Table 7 - Overview mid/upper troposphere CO₂ and CH₄ data products.

Product ID	Level	Sensor(s)	(Planned) Availability	Comments
CO2_IASA_NLIS	2	IASI / Metop-A	Oct. 2017: 2007-2015	
CH4_IASA_NLIS	2	IASI / Metop-A	Oct. 2017: 2007-2015	
CO2_IASB_NLIS	2	IASI / Metop-B	Oct. 2017: 2013-2016 Oct. 2018: 2013-2017 Dec. 2019: 2013-2018	
CH4_IASB_NLIS	2	IASI / Metop-B	Oct. 2017: 2013-2016 Oct. 2018: 2013-2017 Dec. 2019: 2013-2018	
CO2_AIR_NLIS	2	AIRS	Oct. 2017: 2003-2007	Existing GHG-CCI product



2. Product validation methodology

2.1 Description of reference data used for validation

2.1.1 Reference data for validation of the XCO₂ and XCH₄ Level 2 products

2.1.1.1 TCCON network

For validation of satellite XCO₂ and XCH₄ retrievals the Total Carbon Column Observing Network (TCCON, <http://www.tccon.caltech.edu/>) has been established (e.g., *Wunch et al., 2010, 2011, 2015*).

This network is the core network used for validation of the satellite XCO₂ and XCH₄ retrievals. Nevertheless, there are also some limitation as explained in Sect. 2.2.1.4.1.

TCCON provides XCO₂ and XCH₄ data products as retrieved from ground-based Fourier Transform Infrared (FTIR) observations based on direct sun observations. Currently, there are about 20 TCCON sites (see Sect. 2.2.1.4.1).

The TCCON data products can essentially be directly compared with the satellite-derived XCO₂ and XCH₄ data products and TCCON data products have been used for this purpose extensively in the past as shown in many studies and publications. A short overview about these activities is given in Sect. 2.2.1.1.

2.1.1.2 Traceability to standard

As explained in this document, the satellite-derived XCO₂ and XCH₄ data products will be validated by comparison with TCCON XCO₂ and XCH₄ data products, which in turn have been calibrated against the World Meteorological Organization (WMO) *in situ* trace gas measurement scales (see *Wunch et al., 2010*). This approach ensures that the satellite XCO₂ and XCH₄ retrievals are linked to the WMO standards for atmospheric CO₂ and CH₄ measurements.



2.1.2 Reference data for validation of the mid/upper tropospheric CO₂ and CH₄ products

2.1.2.1 Reference data overview

For validation of mid/upper tropospheric CO₂ and CH₄, no remote sensing ground based measurements (such as TCCON) is available. Use is thus made of sparse airborne (aircrafts and balloons) measurements: averaging kernels associated to the retrieved columns are applied to vertical profiles measured by in-situ instruments and the resulting column is compared to columns measured from space.

Validation thus relies on:

- aircraft data acquired either during regular measurements onboard commercial airliners: CONTRAIL, IAGOS in the future.
- aircraft regular measurements made by research groups: NOAA aircraft network in the US and Canada.
- aircraft research campaigns: HIPPO, CoMet in the future.
- Balloon measurements: AirCores at various locations (Timmins, Kiruna, Sodankulä, Trainou-Orléans).

2.1.2.2 Traceability to standard

As explained in the following sections, the satellite mid/upper tropospheric CO₂ and CH₄ will be validated by comparison with aircraft and balloon measurements, which are calibrated against the World Meteorological Organization (WMO) scales. This ensures that the satellite retrievals are linked to WMO standards for atmospheric CO₂ and CH₄.



2.2 Description of product validation methodology

2.2.1 Methods for validation of XCO₂ and XCH₄ Level 2 products

In this section, the validation methodology is described. In the following sections the described methods are applied to the newly generated data sets.

2.2.1.1 Overview validation of GHG-CCI pre-cursor / pre-operational products

Past versions of satellite XCO₂ and XCH₄ retrievals as obtained from SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT have been extensively validated using TCCON as described in various peer-reviewed scientific publications (e.g., *Buchwitz et al., 2013a, 2016; Butz et al., 2010; Cogan et al., 2011; Dils et al., 2004; Parker et al., 2011; Reuter et al., 2011; Schneising et al., 2011; Yoshida et al., 2013*), project related reports (e.g., *Buchwitz et al., 2017*) and other documents (e.g., *Buchwitz et al., 2016a, 2017a; Reuter et al., 2016, 2017a*).

The latest version of the satellite XCO₂ and XCH₄ retrievals as generated within the GHG-CCI project (<http://www.esa-ghg-cci.org/>) of ESA's Climate Change Initiative is called "Climate Research Data Package No. 4" (CRDP4) and is available from the main data products website of the GHG-CCI website (<http://www.esa-ghg-cci.org/> -> CRDP (Data) or directly via http://www.esa-ghg-cci.org/sites/default/files/documents/public/documents/GHG-CCI_DATA.html). The quality assessment of that data set is described in the Product Validation and Intercomparison Report, version 5, PVIRv5 (*Buchwitz et al., 2017*). That GHG-CCI CRDP4 data set is the pre-cursor data set, which has been extended for C3S in the context of the C3S_312a_Lot6 project.

As shown in document PVIRv5 (*Buchwitz et al., 2017*) the validation of the GHG-CCI CRDP4 pre-cursor XCO₂ and XCH₄ data products has been carried out by comparison with TCCON ground-based XCO₂ and XCH₄ retrievals. The assessments have been carried out quasi independently by different individuals / teams using (somewhat) different methods (using all or only a sub-set of the TCCON sites, using different criteria for spatio-temporal co-location, using different methods to compute "relative systematic error" and "year-to-year bias variability, using "direct comparison" or the Ensemble Median Algorithm (EMMA, *Reuter et al., 2013*)) to check and ensure robustness of the findings. Overall it had been found that quite similar overall quality assessment results have been obtained using the different methods (see PVIRv5 for details), i.e., robust conclusions have been obtained.

The quality assessment was based on the computation of several quantities (metrics). The most important ones are:

- Single ground pixel random error (or "single measurement precision", 1-sigma): Computed as the standard deviation of the difference of the single satellite measurement with TCCON.
- Mean bias (per site and globally): Computed as the mean difference of the satellite measurements with TCCON (satellite minus TCCON).



- “Relative systematic error” (or “relative accuracy” or “relative bias”): To estimate this quantity the “spatial bias” had been computed as standard deviation of the biases as obtained at the various individual TCCON sites. This value is reported in several peer-reviewed publications (e.g., *Dils et al., 2014*) but does not consider temporal biases. To also address temporal biases *Dils et al., 2014*, also computed the quantities “seasonal mean bias”, “seasonality” and “seasonal relative accuracy” (SRA).
- Stability: Linear bias trend (drift): Computed from the slope (and the error of the slope) as obtained by fitting a straight line to satellite minus TCCON differences.
- Stability: Year-to-year bias variability: Computed as maximum minus minimum bias difference of smoothed (using a one year running average) satellite minus TCCON differences.

2.2.1.2 Methods to be applied to the C3S ECV CDR data set

The quality assessments, which have been carried out for the newly generated C3S products, are similar as past assessments, which have been carried out for the pre-cursor products (see previous sub-section). However, there are some important differences, in particular those related to Target Requirements (TR) assessments, which have not been carried out for the pre-cursor products. The C3S assessment method is described in the following sub-sections.

2.2.1.2.1 Quantitative assessment methods

For each data product the following quantities have been determined:

Single ground pixel random error (or “single measurement precision”, 1-sigma):

Computed as the standard deviation of the difference of the single satellite retrievals (i.e., for individual ground pixels) with the co-located TCCON reference value. See also document PVIRv5 (*Buchwitz et al., 2017*) for an assessment of this quantity using the pre-cursor products.

Reported uncertainties (“Uncertainty ratio”):

The satellite-derived Level 2 XCO₂ and XCH₄ data products contain an uncertainty estimate for each single observation. This uncertainty is meant to be the statistical uncertainty (1-sigma, dominated by the random error component of the uncertainty due to instrument noise) associated with that single observations. To assess the quality of these uncertainty estimates they are compared with the standard deviation of satellite minus TCCON retrievals at the various TCCON sites. It is expected that the mean value of the reported uncertainty is similar in magnitude (agreement within a few 10%) as the standard deviation of the difference to TCCON (this should be the case if the reported uncertainty is correct and if the comparison method does not introduce additional errors). Therefore, one expects that the “Uncertainty ratio”, i.e., the ratio of the mean value of the reported uncertainty and the standard deviation of satellite minus TCCON differences is close to unity. Although the exact interpretation of this ratio is difficult, it needs to be determined and reported. See also document PVIRv5 (*Buchwitz et al., 2017*) for an assessment of this quantity using the pre-cursor products.

Mean bias:

Computed as the mean difference of satellite minus TCCON retrievals. See also document PVIRv5 (Buchwitz *et al.*, 2017) for an assessment of this quantity using the pre-cursor products.

“Relative systematic error” (or “relative accuracy” or “relative bias” or simply “accuracy”):

To estimate this quantity two values and a combined value are computed and reported:

- The first number is the “spatial bias” computed as standard deviation of the biases as obtained at the various individual TCCON sites. This value is reported in several peer-reviewed publications (e.g., Dils *et al.*, 2014) but does not consider temporal biases (to address this, Dils *et al.*, 2014, computed several quantities: “seasonal mean bias”, “seasonality” and “seasonal relative accuracy”).
- The second number is the “spatio-temporal bias” for a seasonal time scale. There are several options how to compute this number and how to combine it with the first number to get an overall single number for “relative accuracy” and the used method how to exactly compute these numbers has not been fully specified (the most appropriate method may depend on the number of data points, i.e., on the instrument and the applied retrieval algorithm).
- For the QA/QC results presented in this document (and which has been applied to all satellite products discussed in this document) the “spatio-temporal bias”, has been computed as the root-sum-square (RSS) value of the (overall) “spatial bias” and the (overall) “seasonal bias”, i.e., by quadratically adding two numbers.
- The (overall) seasonal bias has been computed as the of the seasonal biases obtained at the individual TCCON sites. The seasonal bias at a given TCCON site has been computed as the standard deviation of the biases in the four (or at least three) seasons. The overall seasonal bias has therefore been computed similarly as the “seasonality” (parameter “Seas”) reported in Dils *et al.*, 2014.
- Because of the used RSS adding method, the “spatio-temporal bias” is always larger than the “spatial bias”. The “spatio-temporal bias” is a positive (or strictly speaking a non-negative) number, and is identified with “relative accuracy” (as it considers spatial and temporal biases).
- However, also other methods are used to compute “spatio-temporal bias” / “relative accuracy”, e.g., by the data provider (DP) method and by the EMMA method (see data quality summary Figure 1 for XCO₂ and Figure 2 for XCH₄), where the results from all assessment methods are presented. In any case, for the combined value, i.e., for “relative accuracy”, always the larger of the two individual values (“spatial bias” and “spatio-temporal bias”) has been used to report the overall value for “relative accuracy” .

Stability: Linear bias trend (Long term drift):

Computed from the slope as obtained by fitting a straight line to satellite minus TCCON differences using the entire time series. Also the 1-sigma uncertainty needs to be reported as obtain from the slope fit error.

Stability: Year-to-year bias variability:

Computed as maximum minus minimum bias difference of smoothed (using a one year running average) satellite minus TCCON differences.



2.2.1.2.2 Qualitative assessment methods

As the TCCON network is quite sparse it is important for quality assessment of the global satellite-derived data product to also use a number of other (more qualitative) assessment methods.

Therefore also the following activities have been carried out:

- Generation of global maps and (regional) time series figures to obtain an overview about the entire data set.
- Comparisons with global models (in particular those assimilating accurate surface CO₂ and CH₄ measurements).

2.2.1.3 Methods for comparison of the achieved performance with the user requirements

The results obtained with the “Quantitative assessment methods” are compared with the Target Requirements (TRs) as given in the Target Requirement Document (TRD) (*TRD GHG, 2017*).

In order to obtain a statement if a certain TR is met or not - or if it is “partially met” - several uncertainties need to be considered as good as possible:

- The uncertainty of the estimated parameter (e.g., the uncertainties of the obtained values for “accuracy” and “stability”).
- The uncertainty of the reference data (here: TCCON) (if not already included in the uncertainty of the obtained values for “accuracy” and “stability”).
- The uncertainty of the comparison method (e.g., considering imperfect collocation of the satellite data and the reference data) (if not already included in the uncertainty of the obtained values for “accuracy” and “stability”).

The following discussion is limited to “accuracy” and “stability” as these are the most critical / important data quality “figures of merit” and because TRs have been defined for them.

The TRs are the following (see also Target Requirement Document (*TRD GHG, 2017*)):

- (Relative) Accuracy XCO₂: < 0.5 ppm (1-sigma)
- Stability XCO₂: < 0.5 ppm/year
- (Relative) Accuracy XCH₄: < 10 ppb (1-sigma)
- Stability XCH₄: < 3 ppb/year

(Relative) Accuracy:

As explained earlier, the term “accuracy” as used here means “relative accuracy” or “relative bias”. The reason for this is that a possible “global offset” is not critical for the main application of the data products, which is to use them to obtain information on (regional) sources and sinks. What is critical is the bias difference between different locations and time periods (“spatio-temporal bias”).



Nevertheless, the “global offset” (a single number per product) has been determined and is reported in this document (and can be taken into account by the users if needed).

“Accuracy” is essentially estimated from standard deviations of the biases at TCCON validation sites. The estimated value is therefore a positive (strictly speaking a non-negative) number. It is assumed for the following (in line with the description as given in Sect. 2.2.1.2.1) that the value obtained for accuracy has been estimated (for each product and each applied assessment method) assuming error free TCCON observations and an error free comparison method (these errors are considered in a later step).

In order to compute the probability that the accuracy requirement is met, it is required to have at least a rough estimate of the uncertainty (“UNC_ACC”) of the reported achieved accuracy value (“ACC”). This uncertainty comes from the uncertainty of the reference data (here TCCON) and the uncertainty of the comparison method (e.g., colocation method and its representativity error).

The uncertainty of the TCCON reference data (see *Wunch et al., 2010*, but also the discussions of this uncertainty related to the use of TCCON data for the validation of satellite retrievals in *Buchwitz et al., 2015, 2016*, and *Dils et al., 2014*) is:

- TCCON uncertainty XCO₂: 0.4 ppm (1-sigma)
- TCCON uncertainty XCH₄: 4 ppb (1-sigma)

These uncertainties are enhanced by 50% to also consider other error sources, especially error of the comparison method such as the “representativity error”. The assumed uncertainty (“UNC_ACC”) of the estimated accuracy value (“ACC”) are therefore:

- UNC_ACC XCO₂: 0.6 ppm (1-sigma)
- UNC_ACC XCH₄: 6 ppb (1-sigma)

In summary, we now have ACC +/- UNC_ACC (1-sigma) for the estimated relative accuracy or spatio-temporal bias and its uncertainty. These values are interpreted as the mean and the standard deviation of an underlying probability density function (pdf).

ACC is a non-negative number and the Target Requirement (TR) for accuracy defines an “acceptable range” or interval of “acceptable” accuracy values: [0, TR[, i.e., in order to meet the requirements ACC shall be smaller than TR but will be larger than (or equal to) zero. Because of this “non-negativity”, ACC cannot be distributed according to a Gaussian (“normal”) probability density function (pdf) (esp. if the mean is much smaller than the standard deviation) but it may be reasonable to assume that the overall distribution is a lognormal distribution (<https://de.mathworks.com/help/stats/lognormal-distribution.html>), with parameters selected such that the lognormal pdf is very similar as a Gaussian pdf if the mean is on the order or larger than the standard deviation.



The probability density function (pdf) of the lognormal distribute is:

$$y = f(x|\mu, \sigma) = \frac{1}{x\sigma\sqrt{2\pi}} \exp\left\{-\frac{(\log x - \mu)^2}{2\sigma^2}\right\}, \quad \text{for } x > 0. \quad \text{Eq. (1)}$$



The lognormal distribution has parameters μ and σ , which are related to parameters mean $m = \text{ACC}$ and variance $v = \text{UNC_ACC}$ as follows:

$$\begin{aligned} \mu &= \log(m^2 / \sqrt{v + m^2}) \\ \sigma &= \sqrt{\log(v/m^2 + 1)} \end{aligned} \quad \text{Eq. (2)}$$

The cumulative distribution function (cdf) of the lognormal distribution is:

$$p = F(x|\mu, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} \int_0^x \frac{1}{t} \exp\left\{-\frac{(\log t - \mu)^2}{2\sigma^2}\right\} dt, \quad \text{for } x > 0. \quad \text{Eq. (3)}$$

This function is used to compute the probability, that the accuracy requirement is met, see Figure 5 for XCO_2 and Figure 6 for XCH_4 .

Figure 5 - Probability that the XCO₂ accuracy TR is met as a function of the achieved accuracy.Figure 6 - Probability that the XCH₄ accuracy TR is met as a function of the achieved accuracy.



Stability:

For the TR assessment, the stability assessment is limited to “Linear bias trend / drift” (i.e., the year-to-year bias variability is also determined as explained above but not used for the TR assessment).

As for “accuracy” it is assumed that the value for stability has been obtained assuming error free TCCON observations and an error free comparison method. In contrast to “accuracy” it is assumed that the uncertainty of the stability value is known (it corresponds to the (1-sigma) slope (SLO) error of the linear fit). The result of the stability assessment is: $STA \pm UNC_SLO$.

To consider the uncertainty of the reference data we assume that the TCCON data approximately meet the following stability requirements:

- XCO_2 stability: 0.2 ppm/year
- XCH_4 stability: 1 ppb/year

These uncertainties need to be added quadratically (via Root-Sum-Square (RSS)) to UNC_SLO to obtain the overall uncertainty UNC_STA .

As shown in Table S-1 for XCO_2 and Table S-2 for XCH_4 in column “Long-term drift” in document PVIRv5 (Buchwitz *et al.*, 2017) typical values for $STA \pm UNC_SLO$ are (if the uncertainty is converted to 1-sigma):

- XCO_2 : $+0.1 \pm 0.07$ (1-sigma) ppm/year
- XCH_4 : -0.8 ± 0.4 (1-sigma) ppb/year

These values are listed here only for illustration (the exact value depends on product and assessment method).

Quadratically adding the assumed TCCON uncertainty gives for this example for $STA \pm UNC_STA$:


- XCO_2 : $+0.1 \pm 0.21$ (1-sigma) ppm/year
- XCH_4 : -0.8 ± 1.08 (1-sigma) ppb/year

In contrast to ACC, STA can also be negative and we use a Gaussian probability density function $N(x, \text{mean}=STA, \text{sigma}=UNC_STA)$ to compute the probability that the stability TR is met. This probability is the integral of N over the interval as defined by the stability TR requirement, i.e., interval $] -TR, +TR[$, or simply the difference between two different values of the cumulative distribution function $Nc(x, \text{mean}=STA, \text{sigma}=UNC_STA)$ (namely at $x=TR$ and $x=-TR$). The probability P that the stability TR is met for XCO_2 for a given value of STA is therefore for this example: $P(STA) = Nc(+0.5, \text{mean}=+0.1, \text{sigma}=0.2) - Nc(-0.5, \text{mean}=+0.1, \text{sigma}=0.2) = 97\%$. This means that in these cases it is almost certain that the stability TR is met.


Figure 7 shows how the used probability functions look like.



Figure 7 – Probability functions used to obtain the probability that the stability requirement is met. Top: for XCO₂. Bottom: for XCH₄.



Michael.Buchwitz@iup.physik.uni-bremen.de 31-Jul-2019



Michael.Buchwitz@iup.physik.uni-bremen.de 31-Jul-2019

2.2.1.4 Known limitations

2.2.1.4.1 TCCON

The TCCON network consists of about 20 TCCON sites (see Figure 8). It is relatively dense in the USA, in Europe and in Japan but overall the TCCON network is relatively sparse (e.g., no or only very few sites in Russia, South America and Africa) and does not cover all conditions, which affect or can affect the quality of the satellite XCO₂ and XCH₄ retrievals (e.g., deserts due to their high surface albedo combined with potentially high amounts of specific aerosol types such as desert dust storm mineral aerosols).

Figure 8 - Location of TCCON sites. Source: <http://www.tccon.caltech.edu/>.



The TCCON network is the core network for the validation of the satellite XCO₂ and XCH₄ retrievals and is therefore absolute essential for this part of the C3S service.

It would therefore be highly beneficial for this service

- if the TCCON network would be expanded to better cover all geophysical conditions relevant for the quality assessment of the satellite retrievals.
- if the TCCON XCO₂ and XCH₄ retrievals would be available faster (current availability: one year after observation).
- if the quality of the TCCON retrievals would be further improved (if possible) as the current data quality (approx. 0.4 ppm for XCO₂ (1-sigma) and 4 ppb for XCH₄ (1-sigma)) is on the order of the required data quality of the satellite retrievals.



As a minimum, it needs to be guaranteed that the existing network remains in place but unfortunately even this is currently not guaranteed.

2.2.2 Methods for validation of XCO₂ and XCH₄ Level 3 Obs4MIPs products

The gridded Level 3 XCO₂ and XCH₄ products are in Obs4MIPs format.

They have been generated such that the products likely get approval from the Obs4MIPs committee (<https://www.earthsystemcog.org/projects/obs4mips/>). The C3S Obs4MIPs products are updates of pre-cursor products generated in the framework of ESA's GHG-CCI project.

The main applications of these products are comparisons with climate models as shown in, e.g., *Lauer et al., 2017*, presenting a comparison of the version 1 XCO₂ Obs4MIPs data product (see also *Reuter et al., 2016*). The version 1 XCH₄ Obs4MIPs product is described in *Buchwitz et al., 2016a*. In February 2017, version 2 of the XCO₂ and XCH₄ Obs4MIPs data products has been generated in the framework of the GHG-CCI project (<http://www.esa-ghg-cci.org/>) covering the time period 2003-2015 (*Buchwitz et al., 2017a; Reuter et al., 2017*).

These products have now been re-generated for C3S and they are extended in time (now covering 2003-2016 (version 3)).

The XCO₂ and XCH₄ Obs4MIPs products are based on the XCO₂ and XCH₄ Level 2 products described in this document. The quality of the Obs4MIPs products therefore depends on the quality of the underlying Level 2 products.

Note that the data quality user requirements for the XCO₂ and XCH₄ products (*TRD GHG, 2017*) are requirements for Level 2 products. Explicit data quality requirement for Level 3 products do not exist. Nevertheless, quality assessments similar as for the Level 2 products have been carried out including TR assessments.



2.2.3 Methods for validation of CO₂ and CH₄ Level 2 mid/upper troposphere products

2.2.3.1 Overview of existing methods as applied to pre-cursor data sets

Past versions of satellite mid/upper tropospheric CO₂ and CH₄ obtained from IASI have been validated using aircraft or, more recently, balloon measurements of atmospheric profiles.

The previous version of the satellite mid/upper tropospheric CO₂ and CH₄ IASI retrievals as generated within the GHG-CCI project (<http://www.esa-ghg-cci.org/>) of ESA's Climate Change Initiative is called "Climate Research Data Package No. 4" (CRDP4) and is available from the main data products website of the GHG-CCI website (<http://www.esa-ghg-cci.org/> -> CRDP (Data) or directly via http://www.esa-ghg-cci.org/sites/default/files/documents/public/documents/GHG-CCI_DATA.html). The quality assessment of this data set is described in the Product Validation and Intercomparison Report, version 5, PVIRv5 (*Buchwitz et al., 2017*). This GHG-CCI CRDP4 data set is the pre-cursor data set, which will be extended for C3S in the context of the C3S_312a_Lot6 project.

As shown in document PVIRv5 (*Buchwitz et al., 2017*) the validation of the GHG-CCI CRDP4 pre-cursor CO₂ and CH₄ mid/upper tropospheric data products has been carried out by comparison with aircraft and balloon-borne AirCores in-situ profile measurements. These comparisons have enabled to validate global trend, growth rate and amplitude of the seasonal cycle. However, due to the scarcity of the measurements, quantity such as single retrieval precision or accuracy remains limited and may be derived only in specific regions where enough measurements are available.

2.2.3.2 Methods applied to the C3S ECV CDR data set

2.2.3.2.1 Quantitative assessment methods

Essentially the same methods have been applied as described in Sect. 2.2.1.2.1 for the XCO₂ and XCH₄ data products, when the number of available aircraft or AirCore measurements of vertical profiles allows the computation of the quantities.

2.2.3.2.2 Qualitative assessment methods

The same methods have been applied as described in Sect. 2.2.1.2.2 for the XCO₂ and XCH₄ data products.

2.2.3.3 Methods for comparison of the achieved performance with the user requirements

Essentially the same methods have been applied as described in Sect. 2.2.1.2.1 for the XCO₂ and XCH₄ data products.



2.2.3.4 Known limitations

The main limitation is the scarcity of measurements in the mid and upper troposphere of CO₂ and CH₄. Moreover, aircraft profiles are generally available up to 6-8 km, which means that the above part of the profile need to be taken from atmospheric transport simulation. This could result in a regional/seasonal bias, which is not well known. Recently developed AirCores, which provide 0-30 km profiles of CO₂ and CH₄ by flying under meteorological balloons, provides a means to fully validate the gas columns retrieved from space, provided that enough measurements are available (less than 20 profiles are currently available worldwide).

For this service, it would thus be highly beneficial:

- if AirCores could be launched regularly at various locations (for instance at existing TCCON/ICOS stations).
- if extensive aircraft campaigns could be organized to collect information in several places where no measurements are currently available (tropical and boreal regions).
- if measurements from IAGOS could include CO₂ and CH₄.



3. Validation results

3.1 Validation results for Level 2 XCO₂ products

In this section, the validation method as explained in the previous section is applied to the XCO₂ and XCH₄ Level 2 individual sensor pre-cursor products, which have been generated in the framework of the ESA project GHG-CCI (*Buchwitz et al., 2016*). The main purpose of this section is to illustrate the method and to show which data quality can be expected from the to be generated C3S XCO₂ and XCH₄ Level 2 data products. The used products are from the latest GHG-CCI data set called “Climate Research Data Package No. 4” (CRDP4, see *Buchwitz et al., 2017*).

For each data product a set of well defined “figures of merit” (FoMs) need to be computed to summarize the validation results and to compute the probability that the TR is met as explained in Sect. 2.2. This can be done using different methods depending on, for example, the chosen co-location criteria and other “filters” such as required number of successful co-locations required to “accept” a certain set of FoM (if the number of co-locations is too small than the obtained FoMs may not be regarded as significant or robust enough).

In the following sub-sections results from one of the methods are presented. This method is a method developed and implemented at Univ. Bremen for the validation of all C3S XCO₂ and XCH₄ Level 2 data products. For the final validation also other methods will be used, in particular the methods applied by each data provider to validate their own data set(s). The same “ensemble approach” for validation has also been used for the GHG-CCI products in the framework of the GHG-CCI project (see *Buchwitz et al., 2017*).

The validation results as shown in this document are based on the TCCON products from 22 TCCON sites: Eureka, NyAlesund, Sodankyla, East Trout Lake, Bialystok, Bremen, Karlsruhe, Paris, Orleans, Garmisch, Park Falls, Lamont, Tsukuba, Edwards, Caltech, Saga, Burgos, Ascension Island, Darwin, Reunion Island, Wollongong, Lauder- Detailed information on each site can be found at <https://tcccon-wiki.caltech.edu/Sites>. The used TCCON version is GGG2014 (data access: 27-May-2019).

Co-location criteria:

- Temporal: +/- 2 hours
- Spatial: +/- 2° latitude, +/- 4° longitude



3.1.1 Validation results for product CO2_SCI_BESD

As a first step, the satellite product is compared with the corresponding TCCON product at each TCCON site separately. Only results from those sites are accepted for further processing if comparisons at least 30 days are possible (note that one day corresponds to one satellite overpass).

Figure 9 shows the comparison at the TCCON site Lamont (“LAM”), Oklahoma, USA. Please see the figure caption for a detailed explanation of the Figures of Merit (FoMs) resulting from this comparison.

As can be seen from Figure 9 also FoMs for seasonal bias and stability are computed. These FoMs are only computed if the time series is “long enough” (at least 3 years) with, for example, a sufficient number of co-locations per season (at least 10 days) and per year (at least 20 days). For Lamont these conditions are fulfilled.

From the results obtained at the individual TCCON sites a single “Product Quality Summary Figure” is produced which is shown as **Figure 10** for product CO2_SCI_BESD. The top right part shows a table listing the FoMs as obtained for the individual TCCON sites (the Lamont (LAM) results are shown in **Figure 9**). Listed are

- the TCCON site ID (e.g., LAM_01 for Lamont),
- the random error or single measurement precision (in ppm, 1-sigma),
- the uncertainty ratio “UncR”, which is the ratio of the reported XCO₂ uncertainty (as reported in the data product for each individual satellite ground pixel) and the estimated uncertainty as computed from the standard deviation of the difference of the individual observations to TCCON (note that a value not too far away from 1.0 is expected for reliable, i.e., “good quality” reported uncertainties),
- the bias in terms of mean bias and seasonal bias (see Figure 9) and
- FoMs characterising stability in terms of drift and year-to-year bias variability (see caption Figure 9 for details).

The FoMs obtained from the individual sites are used to compute “overall quality FoMs” listed directly below the table of the individual TCCON site results. These overall quality FoMs are obtained by computing (i) the “Mean” and (ii) the standard deviation (“StdDev”).

A subset of these FoMs is used to report the final FoMs for the CO2_SCI_BESD product, which are listed in the yellow marked box in the bottom right of **Figure 10**:

- Single measurement precision (1-sigma)
- Uncertainty ratio (“UncR”)
- Relative accuracy computed as standard deviation of the site-to-site biases as a measure of “regional bias” and also as seasonal bias to include a time dependence
- The global offset or mean bias
- The linear drift component of stability and its 1-sigma uncertainty
- The year-to-year bias component of stability and its 1-sigma uncertainty



Also listed are the probabilities that the accuracy TR and the stability (drift) TR is met (see Sect. 2.2.1.3 for details).

These final FoMs are used for **Table 8**, which summarizes the quality assessment results for this product.

Figure 9 - Comparison of satellite XCO₂ product CO2_SCI_BESD (red symbols in top panel) with TCCON XCO₂ (black symbols in top panel). Top: Daily satellite and TCCON XCO₂ (the number of days is listed (in blue) as Ndays). Also listed are the following figures of merit (in blue): the systematic error (mean bias satellite single observations minus TCCON), mean value of the single observation random error, the number of satellite observations (Nobs) used for the comparisons, the uncertainty ratio "UncRatio", which is the ratio of the reported uncertainty (1-sigma, per ground pixel) and the estimated uncertainty as computed from satellite minus TCCON differences, and the linear correlation coefficient of the daily averaged data ("R(daily)"). Bottom: Daily differences satellite minus TCCON (red symbols). The blue symbols show the 3-monthly biases. The light green line shows the biases at yearly resolution (obtained by smoothing the daily biases). The dark green line shows the linear trend. The corresponding numbers are listed at the bottom (reported as mean value and standard deviation) using the same colors as used for the x-y plot: daily bias (in red), 3-monthly bias and overall seasonal bias (blue), linear trend (dark green) and year-to-year bias variability (light green; here the reported value is the peak-to-peak difference and its estimated uncertainty in ppm/year).








Figure 10 - Product Quality Summary Figure for product CO2_SCI_BESD. Please see the main text for a detailed explanation.



2004/06 - 2012/03



Michael.Buchwitz@iup.physik.uni-bremen.de 4-Sep-2019 apricor= -0.13 TCcor= 0.24

Sgl.meas.precision (1-sigma): 2.00 UncR: 0.98
 Global offset (mean bias): -0.19 (single) / -0.23 (daily)
 Rel.accuracy: Spat.: 0.28 Seas.: 0.67 Spat.-temp.: 0.73
 Stability: Drift (+/-1-sigma): 0.02 +/- 0.12
 Stability: Year-to-year (+/-1-sigma): 1.28 +/- 0.36
 TR: Accuray: p(ACC<0.5ppm; 0.73 +/- 0.60): 43%
 Stability (drift): p(STA:+/-0.5ppm/yr; 0.02 +/- 0.24): 97%



Table 8 - Product Quality Summary Table for product CO2_SCI_BESD as obtained by comparison with TCCON reference data using the QA/QC assessment method. The listed requirements are the threshold (T) requirements as given in *TRD GHG, 2017*. For precision (i.e., single observation statistical uncertainty or random error) also the corresponding breakthrough (B) and goal (G) requirements are listed. For the achieved performance of (relative) “Accuracy” two values are listed: The first one is the spatial component of the bias and the second one is the spatio-temporal bias, computed by also considering seasonal biases. The spatio-temporal bias is our estimate of “relative accuracy”. TR refers to “target requirement” and reported is the probability that the corresponding TR is met, i.e., the probabilities that accuracy is better than 0.5 ppm and stability is better than 0.5 ppm/year. Note that for this PQAR document a much more challenging method has been used to compute spatio-temporal bias and (relative) accuracy, compared to previous versions of this document.

Product Quality Summary Table for Product: CO2_SCI_BESD Level: 2, Version: 02.01.02, Time period covered: 1.2003 – 3.2012				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppm]	2.0	< 8 (T) < 3 (B) < 1 (G)	-	-
Uncertainty ratio) in [-]: Ratio reported uncertainty to standard deviation of satellite-TCCON difference	0.98	-	-	No requirement but value close to unity expected for a high quality data product.
Mean bias [ppm]	-0.19	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppm]	Spatial – spatiotemporal: 0.28 – 0.73	< 0.5	Probability that accuracy TR is met: 43%	-
Stability: Drift [ppm/year]	0.02 +/- 0.12 (1-sigma)	< 0.5	Probability that stability TR is met: 97%	-
Stability: Year-to-year bias variability [ppm/year]	1.3 +/- 0.4 (1-sigma)	< 0.5	-	-




3.1.2 Validation results for product CO2_SCI_WFMD

Similar figures as shown in 3.1.1 for product CO2_SCI_BESD are shown in this section but for product CO2_SCI_WFMD.

The Product Quality Summary Table for product CO2_SCI_WFMD is shown as Table 9.

Figure 11 - As Figure 9 but for product CO2_SCI_WFMD.



Michael.Buchwitz@iup.physik.uni-bremen.de 4-Sep-2019 C3S3 coloc:2/2/4 apricor: -0.15 TCcor: -0.02



Figure 12 - As Figure 10 but for product CO2_SCI_WFMD.




Table 9 - Product Quality Summary Table for product CO2_SCI_WFMD.

Product Quality Summary Table for Product: CO2_SCI_WFMD Level: 2, Version: 4.0, Time period covered: 10.2002 – 4.2012				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppm]	2.8	< 8 (T) < 3 (B) < 1 (G)	-	-
Uncertainty ratio) in [-]: Ratio reported uncertainty to standard deviation of satellite-TCCON difference	1.18	-	-	No requirement but value close to unity expected for a high quality data product.
Mean bias [ppm]	0.67	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppm]	Spatial – spatiotemporal: 0.65 – 1.00	< 0.5	Probability that accuracy TR is met: 16%	-
Stability: Drift [ppm/year]	-0.10 +/- 0.19 (1-sigma)	< 0.5	Probability that stability TR is met: 91%	-
Stability: Year-to-year bias variability [ppm/year]	2.11 +/- 0.60 (1-sigma)	< 0.5	-	-




3.1.3 Validation results for product CO2_GOS_OCFP

Similar figures as shown in 3.1.1 for product CO2_SCI_BESD are shown in this section but for product CO2_GOS_OCFP.

The Product Quality Summary Table for product CO2_GOS_OCFP is shown as Table 9.

Figure 13 - As Figure 9 but for product CO2_GOS_OCFP.



Michael.Buchwitz@iup.physik.uni-bremen.de 4-Sep-2019 C3S3 coloc:2/2/4 apricor: 0.02 TCcor: -0.11



Figure 14 - As Figure 10 but for product CO2_GOS_OCFP.




Table 10 - Product Quality Summary Table for product CO2_GOS_OCFP.

Product Quality Summary Table for Product: CO2_GOS_OCFP Level: 2, Version: 7.2, Time period covered: 4.2009 – 12.2018				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppm]	1.8	< 8 (T) < 3 (B) < 1 (G)	-	-
Uncertainty ratio) in [-]: Ratio reported uncertainty to standard deviation of satellite-TCCON difference	1.07	-	-	No requirement but value close to unity expected for a high quality data product.
Mean bias [ppm]	0.27	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppm]	Spatial – spatiotemporal: 0.57 – 0.75	< 0.5	Probability that accuracy TR is met: 41%	-
Stability: Drift [ppm/year]	-0.07 +/- 0.12 (1-sigma)	< 0.5	Probability that stability TR is met: 96%	-
Stability: Year-to-year bias variability [ppm/year]	1.77 +/- 0.46 (1-sigma)	< 0.5	-	-




3.1.4 Validation results for product CO2_GOS_SRFP

Similar figures as shown in 3.1.1 for product CO2_SCI_BESD are shown in this section but for product CO2_GOS_SRFP.

The Product Quality Summary Table for product CO2_GOS_SRFP is shown as Table 11.

Figure 15 - As Figure 9 but for product CO2_GOS_SRFP.



Michael.Buchwitz@iup.physik.uni-bremen.de 4-Sep-2019 C3S3 coloc:2/2/4 apricor: -0.08 TCcor: -0.04



Figure 16 – As Figure 10 but for product CO2_GOS_SRFP.




Table 11 - Product Quality Summary Table for product CO2_GOS_SRFP.

Product Quality Summary Table for Product: CO2_GOS_SRFP Level: 2, Version: 2.3.8, Time period covered: 4.2009 – 12.2018				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppm]	2.1	< 8 (T) < 3 (B) < 1 (G)	-	-
Uncertainty ratio) in [-]: Ratio reported uncertainty to standard deviation of satellite-TCCON difference	0.94	-	-	No requirement but value close to unity expected for a high quality data product.
Mean bias [ppm]	0.22	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppm]	Spatial – spatiotemporal: 0.85 – 1.03	< 0.5	Probability that accuracy TR is met: 14%	-
Stability: Drift [ppm/year]	-0.16 +/- 0.16 (1-sigma)	< 0.5	Probability that stability TR is met: 90%	-
Stability: Year-to-year bias variability [ppm/year]	2.31 +/- 0.61 (1-sigma)	< 0.5	-	-



3.1.5 Validation results for product XCO₂_EMMA

Similar figures as shown in 3.1.1 for product CO₂_SCI_BESD are shown in this section but for product XCO₂_EMMA.

The Product Quality Summary Table for product XCO₂_EMMA is shown as Table 11.

Figure 17 - As Figure 9 but for product XCO₂_EMMA.





Figure 18 - As Figure 10 but for product XCO₂_EMMA.Table 12 - Product Quality Summary Table for product XCO₂_EMMA.

Product Quality Summary Table for Product: XCO ₂ _EMMA Level: 2, Version: 4.1, Time period covered: 1.2003 – 12.2018				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppm]	1.28	< 8 (T) < 3 (B) < 1 (G)	-	-
Uncertainty ratio) in [-]: Ratio reported uncertainty to standard deviation of satellite-TCCON difference	1.09	-	-	No requirement but value close to unity expected for a high quality data product.
Mean bias [ppm]	0.30	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppm]	Spatial – spatiotemporal: 0.60 – 0.78	< 0.5	Probability that accuracy TR is met: 36%	-
Stability: Drift [ppm/year]	-0.06 +/- 0.09 (1-sigma)	< 0.5	Probability that stability TR is met: 97%	-
Stability: Year-to-year bias variability [ppm/year]	1.51 +/- 0.36 (1-sigma)	< 0.5	-	-



3.2 Validation results of Level 2 XCH₄ products

In this section the validation method as explained in the previous section is applied to the XCO₂ and XCH₄ Level 2 individual sensor pre-cursor products, which have been generated in the framework of the ESA project GHG-CCI (*Buchwitz et al., 2016*). The main purpose of this section is to illustrate the method and to show which data quality can be expected from the to be generated C3S XCO₂ and XCH₄ Level 2 data products. The used products are from the latest GHG-CCI data set called “Climate Research Data Package No. 4” (CRDP4, see *Buchwitz et al., 2017*).

For each data product a set of well defined “figures of merit” (FoMs) need to be computed to summarize the validation results and to compute the probability that the TR is met as explained in Sect. 2.2. This can be done using different methods depending on, for example, the chosen co-location criteria and other “filters” such as required number of successful co-locations required to “accept” a certain set of FoM (if the number of co-locations is too small than the obtained FoMs may not be regarded as significant or robust enough).

In the following sub-sections results from one of the methods are presented. This method is a method developed and implemented at Univ. Bremen for the validation of all C3S XCO₂ and XCH₄ Level 2 data products. For the final validation also other methods will be used, in particular the methods applied by each data provider to validate their own data set(s). The same “ensemble approach” for validation has also been used for the GHG-CCI products in the framework of the GHG-CCI project (see *Buchwitz et al., 2017*).

The validation results as shown in this document are based on the TCCON products from 22 TCCON sites: Eureka, NyAlesund, Sodankyla, East Trout Lake, Bialystok, Bremen, Karlsruhe, Paris, Orleans, Garmisch, Park Falls, Lamont, Tsukuba, Edwards, Caltech, Saga, Burgos, Ascension Island, Darwin, Reunion Island, Wollongong, Lauder- Detailed information on each site can be found at <https://tcccon-wiki.caltech.edu/Sites>. The used TCCON version is GGG2014 (data access: 27-May-2019).

Co-location criteria:

- Temporal: +/- 2 hours
- Spatial: +/- 2° latitude, +/- 4° longitude




3.2.1 Validation results for product CH₄_SCI_WFMD


Similar figures as shown in 3.1.1 for product CO₂_SCI_BESD are shown in this section but for product CH₄_SCI_WFMD.

The Product Quality Summary Table for product CH₄_SCI_WFMD is shown as Table 13.


Figure 19 - As Figure 9 but for product CH₄_SCI_WFMD.



Michael.Buchwitz@iup.physik.uni-bremen.de 4-Sep-2019 C3S3 coloc:2/2/4 apricor: 1.01 TCcor: 0.77

Figure 20 - As **Figure 10** but for product CH₄_SCI_WFMD.

2004/06 - 2011/12



Michael.Buchwitz@iup.physik.uni-bremen.de 4-Sep-2019 apricor= -0.3 TCcor= 0.7

Sgl.meas.precision (1-sigma): 87.6 UncR: 0.97

Global offset (mean bias): 3.8 (single) / 3.10 (daily)

Rel.accuracy: Spat.: 11.6 Seas.: 9.4 Spat.-temp.: 14.95

Stability: Drift (+/-1-sigma): -0.08 +/- 2.66

Stability: Year-to-year (+/-1-sigma): 29.56 +/- 7.43

TR: Accuray: p(ACC<10ppb; 14.95 +/- 6.00): 19%

Stability (drift): p(STA:+/-3ppb/yr; -0.08 +/- 2.84): 71%



Table 13 - Product Quality Summary Table for product CH4_SCI_WFMB as obtained by comparison with TCCON reference data. The listed requirements are the threshold (T) requirements as given in *TRD GHG, 2017*. For precision (i.e., single observation statistical uncertainty or random error) also the corresponding breakthrough (B) and goal (G) requirements are listed. For the achieved performance of “Accuracy” two values are listed: The first one is the spatial component of the bias (computed as the standard deviation of the bias as the TCCON sites), the second one is the temporal (seasonal) component of the bias. The probability that the accuracy TR is met is computed using the largest of the two accuracy values.

Product Quality Summary Table for Product: CH4_SCI_WFMD Level: 2, Version: 4.0, Time period covered: 10.2002 – 12.2011				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppb]	88	< 34 (T) < 17 (B) < 9 (G)	-	-
Uncertainty ratio) in [-]: Ratio reported uncertainty to standard deviation of satellite-TCCON difference	0.97	-	-	No requirement but value close to unity expected for a high quality data product.
Mean bias [ppb]	3.8	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppb]	Spatial – spatiotemporal: 11.6 – 14.9	< 10	Probability that accuracy TR is met: 19%	-
Stability: Linear bias trend [ppb/year]	-0.08 +/- 2.66 (1-sigma)	< 3	Probability that stability TR is met: 71%	-
Stability: Year-to-year bias variability [ppb/year]	30 +/- 8 (1-sigma)	< 3	-	-



3.2.2 Validation results for product CH₄_SCI_IMAP

Similar figures as shown in 3.1.1 for product CO₂_SCI_BESD are shown in this section but for product CH₄_SCI_IMAP.

The Product Quality Summary Table for product CH₄_SCI_IMAP is shown as Table 14.

Figure 21 - As Figure 9 but for product CH₄_SCI_IMAP.





Figure 22 - As Figure 10 but for product CH₄_SCI_IMAP.Table 14 - Product Quality Summary Table for product CH₄_SCI_IMAP.

Product Quality Summary Table for Product: CH ₄ _SCI_IMAP Level: 2, Version: 7.2, Time period covered: 1.2003 – 4.2012				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppb]	48	< 34 (T) < 17 (B) < 9 (G)	-	-
Uncertainty ratio) in [-]: Ratio reported uncertainty to standard deviation of satellite-TCCON difference	1.36	-	-	No requirement but value close to unity expected for a high quality data product.
Mean bias [ppb]	4.6	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppb]	Spatial – spatiotemporal: 7.0 – 9.8	< 10	Probability that accuracy TR is met: 62%	-
Stability: Linear bias trend [ppb/year]	0.60 +/- 0.59 (1-sigma)	< 3	Probability that stability TR is met: 98%	-
Stability: Year-to-year bias variability [ppb/year]	26 +/- 7 (1-sigma)	< 3	-	-




3.2.3 Validation results for product CH₄_GOS_OCPR


Similar figures as shown in 3.1.1 for product CO₂_SCI_BESD are shown in this section but for product CH₄_GOS_OCPR.

The Product Quality Summary Table for product CH₄_GOS_OCPR is shown as Table 15.

Figure 23 - As Figure 9 but for product CH₄_GOS_OCPR.



Michael.Buchwitz@iup.physik.uni-bremen.de 4-Sep-2019 C3S3 coloc:2/2/4 apricor: 1.06 TCcor: 0.20

Figure 24 - As Figure 10 but for product CH₄_GOS_OCPR.Table 15 - Product Quality Summary Table for product CH₄_GOS_OCPR.

Product Quality Summary Table for Product: CH ₄ _GOS_OCPR Level: 2, Version: 7.2, Time period covered: 4.2009 – 12.2018				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppb]	13	< 34 (T) < 17 (B) < 9 (G)	-	-
Uncertainty ratio) in [-]: Ratio reported uncertainty to standard deviation of satellite-TCCON difference	0.88	-	-	No requirement but value close to unity expected for a high quality data product.
Mean bias [ppb]	3.0	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppb]	Spatial – spatiotemporal: 4.3 – 4.8	< 10	Probability that accuracy TR is met: 89%	-
Stability: Linear bias trend [ppb/year]	0.73 +/- 0.96 (1-sigma)	< 3	Probability that stability TR is met: 95%	-
Stability: Year-to-year bias variability [ppb/year]	11 +/- 3 (1-sigma)	< 3	-	-




3.2.4 Validation results for product CH₄_GOS_SRPR


Similar figures as shown in 3.1.1 for product CO₂_SCI_BESD are shown in this section but for product CH₄_GOS_SRPR.

The Product Quality Summary Table for product CH₄_GOS_SRPR is shown as Table 16.

Figure 25 - As Figure 9 but for product CH₄_GOS_SRPR.



Michael.Buchwitz@iup.physik.uni-bremen.de 4-Sep-2019 C3S3 coloc:2/2/4 apricor: -1.06 TCcor: 2.28

Figure 26 - As Figure 10 but for product CH₄_GOS_SRPR.Table 16 - Product Quality Summary Table for product CH₄_GOS_SRPR.

Product Quality Summary Table for Product: CH ₄ _GOS_SRPR Level: 2, Version: 2.3.9, Time period covered: 4.2009 – 12.2018				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppb]	13	< 34 (T) < 17 (B) < 9 (G)	-	-
Uncertainty ratio) in [-]: Ratio reported uncertainty to standard deviation of satellite-TCCON difference	0.95	-	-	No requirement but value close to unity expected for a high quality data product.
Mean bias [ppb]	-1.3	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppb]	Spatial – spatiotemporal: 4.0 – 4.6	< 10	Probability that accuracy TR is met: 89%	-
Stability: Linear bias trend [ppb/year]	0.67 +/- 0.96 (1-sigma)	< 3	Probability that stability TR is met: 95%	-
Stability: Year-to-year bias variability [ppb/year]	11 +/- 3 (1-sigma)	< 3	-	-




3.2.5 Validation results for product CH₄_GOS_OCFP


Similar figures as shown in 3.1.1 for product CO₂_SCI_BESD are shown in this section but for product CH₄_GOS_OCFP.

The Product Quality Summary Table for product CH₄_GOS_OCFP is shown as Table 17.

Figure 27 - As Figure 9 but for product CH₄_GOS_OCFP.



Michael.Buchwitz@iup.physik.uni-bremen.de 4-Sep-2019 C3S3 coloc:2/2/4 apricor: 0.93 TCcor: 0.18

Figure 28 - As Figure 10 but for product CH₄_GOS_OCFP.Table 17 - Product Quality Summary Table for product CH₄_GOS_OCFP.

Product Quality Summary Table for Product: CH ₄ _GOS_OCFP Level: 2, Version: 7.2, Time period covered: 4.2009 – 12.2018				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppb]	13	< 34 (T) < 17 (B) < 9 (G)	-	-
Uncertainty ratio) in [-]: Ratio reported uncertainty to standard deviation of satellite-TCCON difference	1.12	-	-	No requirement but value close to unity expected for a high quality data product.
Mean bias [ppb]	2.7	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppb]	Spatial – spatiotemporal: 4.5 – 5.2	< 10	Probability that accuracy TR is met: 88%	-
Stability: Linear bias trend [ppb/year]	-0.10 +/- 1.37 (1-sigma)	< 3	Probability that stability TR is met: 92%	-
Stability: Year-to-year bias variability [ppb/year]	13 +/- 3 (1-sigma)	< 3	-	-




3.2.6 Validation results for product CH₄_GOS_SRF


Similar figures as shown in 3.1.1 for product CO₂_SCI_BESD are shown in this section but for product CH₄_GOS_OCFP.

The Product Quality Summary Table for product CH₄_GOS_SRF is shown as Table 18.

Figure 29 - As Figure 9 but for product CH₄_GOS_SRF.



Michael.Buchwitz@iup.physik.uni-bremen.de 4-Sep-2019 C3S3 coloc:2/2/4 apricor: 1.28 TCcor: 0.50

Figure 30 – As Figure 10 but for product CH₄_GOS_SRFP.Table 18 - Product Quality Summary Table for product CH₄_GOS_SRFP.

Product Quality Summary Table for Product: CH ₄ _GOS_SRFP Level: 2, Version: 2.3.8, Time period covered: 4.2009 – 12.2018				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppb]	12	< 34 (T) < 17 (B) < 9 (G)	-	-
Uncertainty ratio) in [-]: Ratio reported uncertainty to standard deviation of satellite-TCCON difference	0.94	-	-	No requirement but value close to unity expected for a high quality data product.
Mean bias [ppb]	4.1	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppb]	Spatial – spatiotemporal: 4.4 – 5.6	< 10	Probability that accuracy TR is met: 86%	-
Stability: Linear bias trend [ppb/year]	-0.03 +/- 0.72 (1-sigma)	< 3	Probability that stability TR is met: 99%	-
Stability: Year-to-year bias variability [ppb/year]	9.5 +/- 2.3 (1-sigma)	< 3	-	-




3.2.7 Validation results for product XCH₄_EMMA

Similar figures as shown in 3.1.1 for product CO₂_SCI_BESD are shown in this section but for product CH₄_GOS_OCFP.

The Product Quality Summary Table for product CH₄_GOS_SRFP is shown as Table 18.

Figure 31 - As Figure 9 but for product XCH₄_EMMA. The large scatter before mid 2009 is due to the worse performance of SCIAMACHY compared to GOSAT.



Michael.Buchwitz@iup.physik.uni-bremen.de 4-Sep-2019 C3S3 coloc:2/2/4 apricor: 0.86 TCcor: 1.05



Figure 32 – As **Figure 10** but for product XCH4_EMMA. The “outliers” are due to SCIAMACHY only data before mid 2009.




Table 19 - Product Quality Summary Table for product XCH4_EMMA.

Product Quality Summary Table for Product: XCH4_EMMA Level: 2, Version: 3.1, Time period covered: 1.2003 – 12.2018				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppb]	21.2	< 34 (T) < 17 (B) < 9 (G)	-	-
Uncertainty ratio) in [-]: Ratio reported uncertainty to standard deviation of satellite-TCCON difference	1.01	-	-	No requirement but value close to unity expected for a high quality data product.
Mean bias [ppb]	-4.0	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppb]	Spatial – spatiotemporal: 5.2 – 5.6	< 10	Probability that accuracy TR is met: 86%	-
Stability: Linear bias trend [ppb/year]	0.53 ± 0.79 (1-sigma)	< 3	Probability that stability TR is met: 97%	-
Stability: Year-to-year bias variability [ppb/year]	13 ± 3 (1-sigma)	< 3	-	-




3.3 Validation results for Level 3 XCO₂ product

In order to validate this product, it has been compared with Total Carbon Column Observation Network (TCCON, *Wunch et al., 2011*) ground-based XCO₂ retrievals using version GGG2014 (*Wunch et al., 2015*).

The validation has been done similarly as for the Level 2 products but with some exception, e.g., the monthly mean product has been directly compared with monthly mean TCCON data.

Figure 33 shows an overview about all validation results.

Table 20 shows the product quality summary table for this product.

Figure 33 – Overview validation results product XCO₂_OBS4MIPS.

Michael.Buchwitz@iup.physik.uni-bremen.de 26-Jul-2019 C3S3 coloc:5/5 apricor: 0.00

Table 20 – Product Quality Summary Table for product XCO₂_OBS4MIPS.

Product Quality Summary Table for Product: XCO ₂ _OBS4MIPS Level: 3, Version: 3.1, Time period covered: 1.2003 – 12.2018				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Overall uncertainty [ppm]	1.2	-	-	No requirement but small value expected for a high quality data product.
Mean bias [ppm]	0.18	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppm]	Spatio-temporal bias: 0.7 +/- 0.6 (1-sigma)	< 0.5	Probability that accuracy TR is met: 74%	-
Stability: Linear bias trend [ppm/year]	-0.06 +/- 0.20 (1-sigma)	< 0.5	Probability that stability TR is met: 98%	-




3.4 Validation results for Level 3 XCH₄ products

In order to validate this product, it has been compared with Total Carbon Column Observation Network (TCCON, *Wunch et al., 2011*) ground-based XCO₂ retrievals using version GGG2014 (*Wunch et al., 2015*).

The validation has been done similarly as for the Level 2 products but with some exception, e.g., the monthly mean product has been directly compared with monthly mean TCCON data.

Figure 34 shows an overview about all validation results.

Table 21 shows the product quality summary table for this product.

Figure 34 – Overview validation results product XCH₄_OBS4MIPS.Table 21 – Product Quality Summary Table for product XCH₄_OBS4MIPS.

Product Quality Summary Table for Product: XCH ₄ _OBS4MIPS Level: 3, Version: 3.1, Time period covered: 1.2003 – 12.2018				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Overall uncertainty [ppm]	8.7	-	-	No requirement but small value expected for a high quality data product.
Mean bias [ppb]	-2.9	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppb]	Spatio-temporal bias: 4.9 +/- 6.0 (1-sigma)	< 10	Probability that accuracy TR is met: 89%	-
Stability: Linear bias trend [ppb/year]	-0.25 +/- 1.00 (1-sigma)	< 3	Probability that stability TR is met: 100%	-



3.5 Validation results for Level 2 mid-tropospheric products

Detailed validation results are given in Annex E to this document. A summary of the validation results is given in Table 22 - Table 24.

Summary quality IASI CO₂ products:

The single measurement precision of product CO2_IASA_NLIS (from IASI on Metop-A) is 1 ppm. The mean bias (global offset) is 0.57 ppm. The product appears to meet the “relative systematic error” requirement of better than 0.5 ppm: the estimated relative accuracy is in the range 0.46-0.49 ppm. The product is also very stable (-0.01 +/- 0.01 ppm/year (1-sigma)) meeting the requirement for long-term drift. The performance of product CO2_IASB_NLIS (from IASI on Metop-B) seems to be similar.

Summary quality IASI CH₄ products:

The single measurement precision of product CH4_IASA_NLIS (from IASI on Metop-A) is 12 ppb. The mean bias (global offset) is -1.3 ppb. The product appears to meet the “relative systematic error” requirement of better than 10 ppb: the estimated relative accuracy is 5.2 ppb. The product appears to be very stable but a quantitative analysis could not be carried out due to lack of reference data. The performance of product CH4_IASB_NLIS (from IASI on Metop-B) seems to be similar.

For product CO2_AIRS_NLIS (from project GHG-CCI) the estimated performance is: single measurement precision: 1.3 ppm, mean bias: -0.43 ppm.



Table 22 - Product Quality Summary Table for product CO2_IASA_NLIS.

Product Quality Summary Table for Product: CO2_IASA_NLIS Level: 2, Version: 8.0, Time period covered: 7.2007 – 5.2015				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppm]	0.99	< 8 (T) < 3 (B) < 1 (G)	-	-
Mean bias [ppm]	0.57	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: Relative systematic error [ppm]	Spatial – spatiotemporal: 0.46 / 0.49	< 0.5	Probability that accuracy TR is met: 70%	-
Stability: Drift [ppm/year]	-0.01 ± 0.01 (1-sigma)	< 0.5	Probability that stability TR is met: 100%	-
Stability: Year-to-year bias variability [ppm/year]	2.64 ± 0.79 (1-sigma)	< 0.5	-	-



Table 23 - Product Quality Summary Table for products CH4_IASA_NLIS (NC stands for Not computed due to lack of available data).

Product Quality Summary Table for Product: CH4_IASA_NLIS Level: 2, Version: 8.4, Time period covered: 7.2007 – 5.2015				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppb]	11.9	< 34 (T) < 17 (B) < 9 (G)	-	-
Mean bias [ppb]	-1.3	-	-	No requirement but value close to zero expected for a high quality data product.
Accuracy: relative systematic error [ppb]	5.2	< 10	Probability that accuracy TR is met: 90%	-
Stability: Linear bias trend [ppb/year]	NC	< 3	NC	Time series of available aircraft/AirCore obs are not long enough to compute these 2 parameters
Stability: Year-to-year bias variability [ppb/year]	NC	< 3	-	

Table 24 - Product Quality Summary Table for products CO2_AIRS_NLIS.

Product Quality Summary Table for Product: CO2_AIRS_NLIS Level: 2, Version: 3.0, Time period covered: 4.2003 – 7.2007				
Parameter [unit]	Achieved performance	Requirement	TR	Comments
Single measurement precision (1-sigma) in [ppb]	1.32	< 8 (T) < 3 (B) < 1 (G)	-	-
Mean bias [ppb]	-0.43	-	-	No requirement but value close to zero expected for a high quality data product.



4. Application(s) specific assessments

The new data products described and validated in this document and its ANNEXes have not yet been used for application specific assessments in terms of peer-reviewed publications.

Note however, that the ESA GHG-CCI project pre-cursor data sets have been used for many applications related to the natural and anthropogenic sources and sinks of atmospheric carbon dioxide and methane. Please see the publication list as given on the GHG-CCI website <http://www.esa-ghg-cci.org/?q=node/85> (where also links to nearly all publications are given) and see also the references as given in Section REFERENCES.

The previous C3S data set (CDR1, 2003-2016) has been used for the peer-reviewed publication *Buchwitz et al., 2018*. They analysed the initial C3S XCO₂ Obs4MIPs data product to compute and investigate annual mean XCO₂ growth rates. Their study can be summarized as follows:

“The growth rate of atmospheric carbon dioxide (CO₂) reflects the net effect of emissions and uptake resulting from anthropogenic and natural carbon sources and sinks. Annual mean CO₂ growth rates have been determined globally and for selected latitude bands from satellite retrievals of column-average dry-air mole fractions of CO₂, i.e., XCO₂, for the years 2003 to 2016. The global XCO₂ growth rates agree with National Oceanic and Atmospheric Administration (NOAA) growth rates from CO₂ surface observations within the uncertainty of the satellite-derived growth rates (mean difference \pm standard deviation: 0.0 ± 0.24 ppm/year; R: 0.87). This new and independent data set confirms record large growth rates around 3 ppm/year in 2015 and 2016, which are attributed to the 2015/2016 El Niño. Based on a comparison of the satellite-derived growth rates with human CO₂ emissions from fossil fuel combustion and with El Niño Southern Oscillation (ENSO) indices, we estimate by how much the impact of ENSO dominates the impact of fossil fuel burning related emissions in explaining the variance of the atmospheric CO₂ growth rate.”

Using the new XCO₂_OBS4MIPS version 4.1 data product the satellite-derived XCO₂ growth rate time series has been updated and extended using the method described in *Buchwitz et al., 2018*. The results are shown in Figure 35. As can be seen from panel (b), the satellite-derived growth rates agree with the NOAA growth rates within the error bar (1-sigma) of the satellite-derived growth rates as also shown for the previous data set (see the detailed discussion in *Buchwitz et al., 2018*). As can be seen, the estimated XCO₂ year 2018 growth rate is 2.1 ± 0.5 ppm/year (NOAA: 2.43 ± 0.08 ppm/year). For details see Reuter et al., 2019.

The method of *Buchwitz et al., 2018*, has now also been applied to the satellite methane observations using product XCH₄_OBS4MIPS version 4.1. The results are shown in Figure 36. As can be seen, the agreement with NOAA is much better for 2010 and later years, i.e., for the time period where GOSAT data have been added. Before 2010 the product is entirely based on SCIAMACHY retrievals, which are less precise and accurate compared to the GOSAT retrievals. Note also that the satellite product is based on an ensemble of several satellite-derived methane products since 2010 and this is expected to also increase the quality of this data product. During the 2010-2018 time period the average XCH₄ increase is 7.9 ± 0.2 ppb. For details see Reuter et al., 2019.



Figure 35 – (a) Monthly mean XCO₂ time series 2003-2018., (b) Corresponding annual growth rates (red) compared to NOAA growth rates obtained from surface CO₂ observations (blue; source: https://www.esrl.noaa.gov/gmd/ccgg/trends/gl_gr.html; access: 30-Jul-2019). From: Reuter et al., 2019.







Figure 36 – (a) Monthly mean XCH_4 time series 2003-2018., (b) Corresponding annual growth rates (red) compared to NOAA growth rates obtained from surface CH_4 observations (blue; source: ftp://aftp.cmdl.noaa.gov/products/trends/ch4/ch4_gr_gl.txt; access: 30-Jul-2019). From: Reuter et al., 2019.





5. Compliance with user requirements

5.1 Level 2 XCO₂ and XCH₄ products

XCO₂:

Figure 1 shows a summary of the achieved performance in terms of single measurement precision, (relative) accuracy (in terms of spatial and spatio-temporal biases) and stability (in terms of linear bias drift / trend).

As can be seen, the achieved single observation random error (or precision) is on the order of 1.5 ppm and better than 3 ppm for all products. This is better than the required breakthrough requirement (B) of better than 3 ppm but somewhat worse than the goal (G) requirement of better than 1 ppm.

The systematic error (relative accuracy) threshold (T) requirement is “better than 0.5 ppm”. The achieved performance is around 0.7 ppm +/- a few 0.1 ppm, depending on product and assessment method. The probability that the threshold requirement is met is between 36% and 64%, depending on product.

Stability is very good. No significant linear bias drift has been detected. The probability that the threshold (T) stability requirement of 0.5 ppm/year is met is larger than 87% for all products.

XCH₄:

Figure 2 shows a summary of the achieved performance in terms of single measurement precision, (relative) accuracy (in terms of spatial and spatio-temporal biases) and stability (in terms of linear bias drift / trend).

As can be seen, the achieved single observation random error (or precision) is better than 17 ppb, which is the breakthrough (B) requirement, for the GOSAT and the EMMA products. For SCIAMACHY the precision is worse (50-90 ppb).

The systematic error (relative accuracy) threshold (T) requirement is “better than 10 ppb”. The achieved performance is around 5 ppb for the GOSAT and the EMMA products. For SCIAMACHY the achieved accuracy is around 10 ppb.

Stability is very good for all products. No significant linear bias drift has been detected. The probability that the threshold (T) stability requirement of 3 ppb/year is met is larger than 85% for all products.



5.2 Level 3 XCO₂ and XCH₄ products

The quality assessment results for the XCO₂ product XCO2_OBS4MIPS are:

The estimated accuracy is 0.7 +/- 0.6 ppm and the probability that the 0.5 ppm requirement is met is 47%.

The linear bias trend is -0.06 +/- 0.2 ppm/year and the probability that the 0.5 ppm/year requirement is met is 98%.

Overall, this product has therefore reasonable accuracy and high stability.

The quality assessment results for the XCH₄ product XCH4_OBS4MIPS are:

The estimated accuracy is 4.9 +/- 6 ppb and the probability that the 10 ppb requirement is met is 89%.

The linear bias trend is 0.25 +/- 1 ppb/year and the probability that the 3 ppm/year requirement is met is 100%.

Overall, this product has therefore very good accuracy and high stability.



5.3 Level 2 mid-tropospheric products

Summary quality IASI CO₂ products:

The single measurement precision of product CO2_IASA_NLIS (from IASI on Metop-A) is 1 ppm. The mean bias (global offset) is 0.57 ppm. The product appears to meet the “relative systematic error” requirement of better than 0.5 ppm: the estimated relative accuracy is in the range 0.46-0.49 ppm. The product is also very stable (-0.01 +/- 0.01 ppm/year (1-sigma)) meeting the requirement for long-term drift. The performance of product CO2_IASB_NLIS (from IASI on Metop-B) seems to be similar.

Summary quality IASI CH₄ products:

The single measurement precision of product CH4_IASA_NLIS (from IASI on Metop-A) is 12 ppb. The mean bias (global offset) is -1.3 ppb. The product appears to meet the “relative systematic error” requirement of better than 10 ppb: the estimated relative accuracy is 5.2 ppb. The product appears to be very stable but a quantitative analysis could not be carried out due to lack of reference data. The performance of product CH4_IASB_NLIS (from IASI on Metop-B) seems to be similar.

For product CO2_AIRS_NLIS (from project GHG-CCI) the estimated performance is: single measurement precision: 1.3 ppm, mean bias: -0.43 ppm.



References

- Alexe et al., 2015:** Alexe, M., P. Bergamaschi, A. Segers, R. Detmers, A. Butz, O. Hasekamp, S. Guerlet, R. Parker, H. Boesch, C. Frankenberg, R. A. Scheepmaker, E. Dlugokencky, C. Sweeney, S. C. Wofsy, and E. A. Kort, [Inverse modeling of CH₄ emissions for 2010–2011 using different satellite retrieval products from GOSAT and SCIAMACHY](#), Atmos. Chem. Phys., 15, 113–133, doi:10.5194/acp-15-113-2015, 2015.
- ATBD GHG, 2017:** Buchwitz, M., Aben, I., Anand, J., Armante, R., Boesch, H., Crevoisier, C., Detmers, R. G., Hasekamp, O. P., Reuter, M., Schneising-Weigel, O., Algorithm Theoretical Basis Document (ATBD) – Main document, C3S project C3S_312a_Lot6_IUP-UB – Greenhouse Gases, 2017.
- ATBD GHG, 2018:** Buchwitz, M., Aben, I., Anand, J., Armante, R., Boesch, H., Crevoisier, C., Detmers, R. G., Hasekamp, O. P., Reuter, M., Schneising-Weigel, O., Algorithm Theoretical Basis Document (ATBD) – Main document, C3S project C3S_312a_Lot6_IUP-UB – Greenhouse Gases, v2.0, 2018.
- ATBD GHG, 2019:** Buchwitz, M., Aben, I., J., Armante, R., Boesch, H., Crevoisier, C., Di Noia, A., Hasekamp, O. P., Reuter, M., Schneising-Weigel, O., Wu, L., Algorithm Theoretical Basis Document (ATBD) – Main document for Greenhouse Gas (GHG: CO₂ & CH₄) data set CDR 3 (2003-2018), C3S project C3S_312b_Lot2_DLR, v3.0, 2019.
- Bergamaschi et al., 2009:** Bergamaschi, P., Frankenberg, C., Meirink, J. F., Krol, M., Villani, M. G., Houweling, S., Dentener, F., Dlugokencky, E. J., Miller, J. B., Gatti, L. V., Engel, A., and Levin, I.: Inverse modeling of global and regional CH₄ emissions using SCIAMACHY satellite retrievals, J. Geophys. Res., 114, D22301, doi:10.1029/2009JD012287, 2009.
- Bergamaschi et al., 2013:** Bergamaschi, P., Houweling, H., Segers, A., et al., [Atmospheric CH₄ in the first decade of the 21st century: Inverse modeling analysis using SCIAMACHY satellite retrievals and NOAA surface measurements](#), J. Geophys. Res., 118, 7350–7369, doi:10.1002/jrgd.50480, 2013.
- Boesch et al., 2011:** Boesch, H., D. Baker, B. Connor, D. Crisp, and C. Miller, Global characterization of CO₂ column retrievals from shortwave-infrared satellite observations of the Orbiting Carbon Observatory-2 mission, Remote Sensing, 3 (2), 270–304, 2011.
- Bovensmann et al., 1999:** Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noël, S., Rozanov, V. V., Chance, K. V., Goede, A. H. P. (1999), SCIAMACHY - Mission objectives and measurement modes, J. Atmos. Sci., 56 (2), 127–150, 1999.
- Bovensmann et al., 2010:** Bovensmann, H., Buchwitz, M., Burrows, J. P., Reuter, M., Krings, T., Gerilowski, K., Schneising, O., Heymann, J., Tretner, A., and Erzinger, J.: A remote sensing technique for global monitoring of power plant CO₂ emissions from space and related applications, Atmos. Meas. Tech., 3, 781–811, 2010.
- Buchwitz et al., 2000:** Buchwitz, M., Rozanov, V. V., and Burrows, J. P.: A near-infrared optimized DOAS method for the fast global retrieval of atmospheric CH₄, CO, CO₂, H₂O, and N₂O total column amounts from SCIAMACHY Envisat-1 nadir radiances, J. Geophys. Res. 105, 15,231–15,245, 2000.
- Buchwitz et al., 2005:** Buchwitz, M., R. de Beek, J. P. Burrows, H. Bovensmann, T. Warneke, J. Notholt, J. F. Meirink, A. P. H. Goede, P. Bergamaschi, S. Körner, M. Heimann, and A. Schulz,



Atmospheric methane and carbon dioxide from SCIAMACHY satellite data: Initial comparison with chemistry and transport models, *Atmos. Chem. Phys.*, 5, 941-962, 2005.

Buchwitz et al., 2007: Buchwitz, M., O. Schneising, J. P. Burrows, H. Bovensmann, M. Reuter, J. Notholt, First direct observation of the atmospheric CO₂ year-to-year increase from space, *Atmos. Chem. Phys.*, 7, 4249-4256, 2007.

Buchwitz et al., 2013a: Buchwitz, M., M. Reuter, O. Schneising, H. Boesch, S. Guerlet, B. Dils, I. Aben, R. Armante, P. Bergamaschi, T. Blumenstock, H. Bovensmann, D. Brunner, B. Buchmann, J. P. Burrows, A. Butz, A. Chédin, F. Chevallier, C. D. Crevoisier, N. M. Deutscher, C. Frankenberg, F. Hase, O. P. Hasekamp, J. Heymann, T. Kaminski, A. Laeng, G. Lichtenberg, M. De Mazière, S. Noël, J. Notholt, J. Orphal, C. Popp, R. Parker, M. Scholze, R. Sussmann, G. P. Stiller, T. Warneke, C. Zehner, A. Bril, D. Crisp, D. W. T. Griffith, A. Kuze, C. O'Dell, S. Oshchepkov, V. Sherlock, H. Suto, P. Wennberg, D. Wunch, T. Yokota, Y. Yoshida, The Greenhouse Gas Climate Change Initiative (GHG-CCI): comparison and quality assessment of near-surface-sensitive satellite-derived CO₂ and CH₄ global data sets, *Remote Sensing of Environment*, doi:10.1016/j.rse.2013.04.024, <http://authors.elsevier.com/sd/article/S0034425713003520>, 2013.

Buchwitz et al., 2013b: Buchwitz, M., Reuter, M., Bovensmann, H., Pillai, D., Heymann, J., Schneising, O., Rozanov, V., Krings, T., Burrows, J. P., Boesch, H., Gerbig, C., Meijer, Y., and Loesch, A.: Carbon Monitoring Satellite (CarbonSat): assessment of atmospheric CO₂ and CH₄ retrieval errors by error parameterization, *Atmos. Meas. Tech.*, 6, 3477-3500, 2013.

Buchwitz et al., 2014: Buchwitz, M., et al.: ESA Climate Change Initiative (CCI) Product Specification Document (PSD) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG), 6-June-2014, Version 3, 2014. Link: http://www.esa-ghg-cci.org/index.php?q=webfm_send/160

Buchwitz et al., 2015: Buchwitz, M., Reuter, M., Schneising, O., Boesch, H., Guerlet, S., Dils, B., Aben, I., Armante, R., Bergamaschi, P., Blumenstock, T., Bovensmann, H., Brunner, D., Buchmann, B., Burrows, J.P., Butz, A., Chédin, A., Chevallier, F., Crevoisier, C.D., Deutscher, N.M., Frankenberg, C., Hase, F., Hasekamp, O.P., Heymann, J., Kaminski, T., Laeng, A., Lichtenberg, G., De Mazière, M., Noël, S., Notholt, J., Orphal, J., Popp, C., Parker, R., Scholze, M., Sussmann, R., Stiller, G.P., Warneke, T., Zehner, C., Bril, A., Crisp, D., Griffith, D.W.T., Kuze, A., O'Dell, C., Oshchepkov, S., Sherlock, V., Suto, H., Wennberg, P., Wunch, D., Yokota, T., Yoshida, Y., The Greenhouse Gas Climate Change Initiative (GHG-CCI): comparison and quality assessment of near-surface-sensitive satellite-derived CO₂ and CH₄ global data sets. *Remote Sens. Environ.* 162:344–362, <http://dx.doi.org/10.1016/j.rse.2013.04.024>, 2015.

Buchwitz et al., 2016: Buchwitz, M., Reuter, M., Schneising, O., Hewson, W., Detmers, R. G., Boesch, H., Hasekamp, O. P., Aben, I., Bovensmann, H., Burrows, J. P., Butz, A., Chevallier, F., Dils, B., Frankenberg, C., Heymann, J., Lichtenberg, G., De Mazière, M., Notholt, J., Parker, R., Warneke, T., Zehner, C., Griffith, D. W. T., Deutscher, N. M., Kuze, A., Suto, H., and Wunch, D.: Global satellite observations of column-averaged carbon dioxide and methane: The GHG-CCI XCO₂ and XCH₄ CRDP3 data, *Remote Sensing of Environment* (in press), Special Issue on Essential Climate Variables, DOI: 10.1016/j.rse.2016.12.027, (link: <http://dx.doi.org/10.1016/j.rse.2016.12.027>), 2016.

Buchwitz et al., 2016a: Buchwitz, M.; Reuter, M.; Aben, I.; Boesch, H.; Butz, A.; Detmers, R.G.; Frankenberg, C.; Hasekamp, O.P.; Parker, R.; Schneising, O.; Somkuti, P., ESA Greenhouse Gases Climate Change Initiative (GHG-CCI): Merged SCIAMACHY and GOSAT Level 3 gridded atmospheric



column-average methane (XCH₄) product in Obs4MIPs format, Centre for Environmental Data Analysis, 10 October 2016, doi:10.5285/C965E4AC-D2AF-4BAA-9E99-A234E9BA0193, link: http://www.esa-ghg-cci.org/?q=webfm_send/331, pp. 11, 2016.

Buchwitz et al., 2017: ESA Climate Change Initiative (CCI) Product Validation and Intercomparison Report (PVIR) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG) for data set Climate Research Data Package No. 4 (CRDP#4), Version 5.0, 9. Feb. 2017, link: http://www.esa-ghg-cci.org/?q=webfm_send/352, 2017.

Buchwitz et al., 2017a: Buchwitz, M.; Reuter, M.; Aben, I.; Boesch, H.; Butz, A.; Detmers, R.G.; Frankenberg, C.; Hasekamp, O.P.; Parker, R.; Schneising, O.; Somkuti, P., ESA Greenhouse Gases Climate Change Initiative (GHG-CCI): Merged SCIAMACHY and GOSAT Level 3 gridded atmospheric column-average methane (XCH₄) product in Obs4MIPs format version 2 (CRDP#4), Technical Note, link: http://www.esa-ghg-cci.org/?q=webfm_send/349, pp. 11, 1 February 2017, 2017.

Buchwitz et al., 2018: Buchwitz, M., Reuter, M., Schneising, O., Noël, S., Gier, B., Bovensmann, H., Burrows, J. P., Boesch, H., Anand, J., Parker, R. J., Somkuti, P., Detmers, R. G., Hasekamp, O. P., Aben, I., Butz, A., Kuze, A., Suto, H., Yoshida, Y., Crisp, D., and O'Dell, C., Computation and analysis of atmospheric carbon dioxide annual mean growth rates from satellite observations during 2003–2016, *Atmos. Chem. Phys.*, *Atmos. Chem. Phys.*, 18, 17355–17370, <https://doi.org/10.5194/acp-18-17355-2018>, 2018.

Burrows et al., 1995: Burrows, J. P., Hölzle, E., Goede, A. P. H., Visser, H., and Fricke, W., SCIAMACHY—Scanning Imaging Absorption Spectrometer for Atmospheric Chartography, *Acta Astronaut.*, 35(7), 445–451, doi:10.1016/0094-5765(94)00278-t, 1995.

Butz et al., 2011: Butz, A., Guerlet, S., Hasekamp, O., et al., Toward accurate CO₂ and CH₄ observations from GOSAT, *Geophys. Res. Lett.*, doi:10.1029/2011GL047888, 2011.

Butz et al., 2012: Butz, A., Galli, A., Hasekamp, O., Landgraf, J., Tol, P., and Aben, I.: Remote Sensing of Environment, TROPOMI aboard Sentinel-5 Precursor : Prospective performance of CH₄ retrievals for aerosol and cirrus loaded atmospheres, 120, 267–276, doi:10.1016/j.rse.2011.05.030, 2012.

Chédin et al. 2003: Chédin, A., Saunders, R., Hollingsworth, A., Scott, N. A., Matricardi, M., Etcheto, J., Clerbaux, C., Armante, R. and Crevoisier, C.: The feasibility of monitoring CO₂ from high resolution infrared sounders. *J. Geophys. Res.*, 108, ACH 6-1–6-19, doi: 10.1029/2001JD001443, 2003.

Chevallier et al., 2005: Chevallier, F., R. J. Engelen, and P. Peylin, The contribution of AIRS data to the estimation of CO₂ sources and sinks. *Geophys. Res. Lett.*, 32, L23801, doi:10.1029/2005GL024229, 2005.

Chevallier et al., 2007: Chevallier, F., F.-M. Bréon, and P. J. Rayner, Contribution of the Orbiting Carbon Observatory to the estimation of CO₂ sources and sinks: Theoretical study in a variational data assimilation framework. *J. Geophys. Res.*, 112, D09307, doi:10.1029/2006JD007375, 2007.

Chevallier et al., 2009a: Chevallier, F., R. J. Engelen, C. Carouge, T. J. Conway, P. Peylin, C. Pickett-Heaps, M. Ramonet, P. J. Rayner and I. Xueref-Remy, AIRS-based vs. surface-based estimation of carbon surface fluxes. *J. Geophys. Res.*, 114, D20303, doi:10.1029/2009JD012311, 2009.



- Chevallier et al., 2009b:** Chevallier, F., S. Maksyutov, P. Bousquet, F.-M. Bréon, R. Saito, Y. Yoshida, and T. Yokota, On the accuracy of the CO₂ surface fluxes to be estimated from the GOSAT observations. *Geophys. Res. Lett.*, 36, L19807, doi:10.1029/2009GL040108, 2009.
- Chevallier et al., 2010:** Chevallier, F., Feng, L., Boesch, H. Palmer, P., and Rayner, P., On the impact of transport model errors for the estimation of CO₂ surface fluxes from GOSAT observations, *Geophys. Res. Lett.*, 37, L21803, 2010.
- Chevallier et al., 2014:** Chevallier, F., Palmer, P.I., Feng, L., Boesch, H., O'Dell, C.W., Bousquet, P., [Towards robust and consistent regional CO₂ flux estimates from in situ and space-borne measurements of atmospheric CO₂](#), *Geophys. Res. Lett.*, 41, 1065-1070, DOI: 10.1002/2013GL058772, 2014.
- Chevallier et al., 2016b:** Chevallier, F., et al., Climate Assessment Report (CAR), ESA Climate Change Initiative (CCI) GHG-CCI project, Version 3, 3 May 2016, link: http://www.esa-ghg-cci.org/?q=webfm_send/318, 2016.
- Ciais et al., 2014:** Ciais, P., Dolman, A. J., Bombelli, A., et al.: Current systematic carbon cycle observations and needs for implementing a policy-relevant carbon observing system, *Biogeosciences*, 11, 3547-3602, www.biogeosciences.net/11/3547/2014/, doi:10.5194/bg-11-3547-2014, 2014.
- Ciais et al., 2015:** Ciais, P., et al.: Towards a European Operational Observing System to Monitor Fossil CO₂ emissions - Final Report from the expert group, <http://www.copernicus.eu/main/towards-european-operational-observing-system-monitor-fossil-co2-emissions>, pp. 68, October 2015, 2015.
- CMUG-RBD, 2010:** Climate Modelling User Group Requirements Baseline Document, Deliverable 1.2, Number D1.2, Version 1.3, 2 Nov 2010.
- Cogan et al., 2011:** Cogan, A. J., Boesch, H., Parker, R. J., et al., Atmospheric carbon dioxide retrieved from the Greenhouse gases Observing SATellite (GOSAT): Comparison with ground-based TCCON observations and GEOS-Chem model calculations, *J. Geophys. Res.*, 117, D21301, doi:10.1029/2012JD018087, 2012.
- Corbin et al., 2008:** Corbin, K. D., A. S. Denning, L. Lu, J.-W. Wang, and I. T. Baker, Possible representation errors in inversions of satellite CO₂ retrievals, *J. Geophys. Res.*, 113, D02301, doi:10.1029/2007JD008716, 2008.
- Cressot et al., 2014:** Cressot, C., F. Chevallier, P. Bousquet, et al., On the consistency between global and regional methane emissions inferred from SCIAMACHY, TANSO-FTS, IASI and surface measurements, *Atmos. Chem. Phys.*, 14, 577-592, 2014.
- Crevoisier et al., 2004:** Crevoisier, C., S. Heilliette, A. Chédin, S. Serrar, R. Armante, and N. A. Scott, Midtropospheric CO₂ concentration retrieval from AIRS observations in the tropics, *Geophys. Res. Lett.*, 31, L17106, doi:10.1029/2004GL020141, 2004.
- Chevallier et al., 2005:** Chevallier, F., R. J. Engelen, and P. Peylin, The contribution of AIRS data to the estimation of CO₂ sources and sinks. *Geophys. Res. Lett.*, 32, L23801, doi:10.1029/2005GL024229, 2005.



- Crevoisier et al., 2004:** Crevoisier, C., S. Heilliette, A. Chédin, S. Serrar, R. Armante, and N. A. Scott, Midtropospheric CO₂ concentration retrieval from AIRS observations in the tropics, *Geophys. Res. Lett.*, 31, L17106, doi:10.1029/2004GL020141, 2004.
- Crevoisier et al., 2009:** Crevoisier, C., Chédin, A., Matsueda, H., et al., First year of upper tropospheric integrated content of CO₂ from IASI hyperspectral infrared observations, *Atmos. Chem. Phys.*, 9, 4797-4810, 2009.
- Crevoisier et al. 2009b:** Crevoisier, C., Nobileau, D., Fiore, A., Armante, R., Chédin, A., and Scott, N. A.: Tropospheric methane in the tropics – first year from IASI hyperspectral infrared observations, *Atmos. Chem. Phys.*, 9, 6337–6350, doi:10.5194/acp-9-6337-2009, 2009b.
- Crevoisier et al., 2013:** Crevoisier, C., Nobileau, D., Armante, R., et al., The 2007–2011 evolution of tropical methane in the mid-troposphere as seen from space by MetOp-A/IASI, *Atmos. Chem. Phys.*, 13, 4279-4289, 2013.
- Crisp et al., 2004:** Crisp, D., Atlas, R. M., Breon, F.-M., Brown, L. R., Burrows, J. P., Ciais, P., Connor, B. J., Doney, S. C., Fung, I. Y., Jacob, D. J., Miller, C. E., O'Brien, D., Pawson, S., Randerson, J. T., Rayner, P., Salawitch, R. S., Sander, S. P., Sen, B., Stephens, G. L., Tans, P. P., Toon, G. C., Wennberg, P. O., Wofsy, S. C., Yung, Y. L., Kuang, Z., Chudasama, B., Sprague, G., Weiss, P., Pollock, R., Kenyon, D., and Schroll, S.: The Orbiting Carbon Observatory (OCO) mission, *Adv. Space Res.*, 34, 700-709, 2004.
- Detmers et al., 2015:** Detmers, R. G., O. Hasekamp, I. Aben, S. Houweling, T. T. van Leeuwen, A. Butz, J. Landgraf, P. Koehler, L. Guanter, and B. Poulter, [Anomalous carbon uptake in Australia as seen by GOSAT](#), *Geophys. Res. Lett.*, 42, doi:10.1002/2015GL065161, 2015.
- Dils et al., 2014:** B. Dils, M. Buchwitz, M. Reuter, O. Schneising, H. Boesch, R. Parker, S. Guerlet, I. Aben, T. Blumenstock, J. P. Burrows, A. Butz, N. M. Deutscher, C. Frankenberg, F. Hase, O. P. Hasekamp, J. Heymann, M. De Mazière, J. Notholt, R. Sussmann, T. Warneke, D. Griffith, V. Sherlock, D. Wunch :The Greenhouse Gas Climate Change Initiative (GHG-CCI): Comparative validation of GHG-CCI SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT CO₂ and CH₄ retrieval algorithm products with measurements from the TCCON network, *Atmos. Meas. Tech.*, 7, 1723-1744, 2014.
- ESA-CCI-GHG-URDv2.1:** Chevallier, F., et al., User Requirements Document (URD), ESA Climate Change Initiative (CCI) GHG-CCI project, Version 2.1, 19 Oct 2016, link: http://www.esa-ghg-cci.org/?q=webfm_send/344, 2016.
- Frankenberg et al., 2011:** Frankenberg, C., Aben, I., Bergamaschi, P., et al., Global column-averaged methane mixing ratios from 2003 to 2009 as derived from SCIAMACHY: Trends and variability, *J. Geophys. Res.*, doi:10.1029/2010JD014849, 2011.
- GCOS-154:** Global Climate Observing System (GCOS): SYSTEMATIC OBSERVATION REQUIREMENTS FOR SATELLITE-BASED DATA PRODUCTS FOR CLIMATE - 2011 Update - Supplemental details to the satellite-based component of the “Implementation Plan for the Global Observing System for Climate in Support of the UNFCCC (2010 Update)”, December 2011, prepared by World Meteorological Organization (WMO), Intergovernmental Oceanographic Commission, United Nations Environment Programme (UNEP), International Council for Science, Doc.: GCOS 154, link: <http://cci.esa.int/sites/default/files/gcos-154.pdf>, 2011.



GCOS-200: The Global Observing System for Climate: Implementation Needs, World Meteorological Organization (WMO), GCOS-200 (GOOS-214), pp. 325, link:

http://unfccc.int/files/science/workstreams/systematic_observation/application/pdf/gcos_ip_10oct2016.pdf, 2016.

Guerlet et al., 2013: Guerlet, S., S. Basu, A. Butz, M. Krol, P. Hahne, S. Houweling, O. P. Hasekamp and I. Aben, [Reduced carbon uptake during the 2010 Northern Hemisphere summer from GOSAT](#), Geophys. Res. Lett., doi: 10.1002/grl.50402, 2013.

Hayman et al., 2014: Hayman, G. D., O'Connor, F. M., Dalvi, M., Clark, D. B., Gedney, N., Huntingford, C., Prigent, C., Buchwitz, M., Schneising, O., Burrows, J. P., Wilson, C., Richards, N., Chipperfield, M., Comparison of the HadGEM2 climate-chemistry model against in-situ and SCIAMACHY atmospheric methane data, Atmos. Chem. Phys., 14, 13257-13280, doi:10.5194/acp-14-13257-2014, 2014.

Hollmann et al., 2013: Hollmann, C.J. Merchant, R. Saunders, C. Downy, M. Buchwitz, A. Cazenave, E. Chuvieco, P. Defourny, G. de Leeuw, R. Forsberg, T. Holzer-Popp, F. Paul, S. Sandven, S. Sathyendranath, M. van Roozendaal, W. Wagner, [The ESA Climate Change Initiative: satellite data records for essential climate variables](#), Bulletin of the American Meteorological Society (BAMS), 0.1175/BAMS-D-11-00254.1, pp. 12, 2013.

Houweling et al., 2004: Houweling, S., Breon, F.-M., Aben, I., Rödenbeck, C., Gloor, M., Heimann, M. and Ciais, P.: Inverse modeling of CO₂ sources and sinks using satellite data: A synthetic inter-comparison of measurement techniques and their performance as a function of space and time, Atmos. Chem. Phys., 4, 523-538, 2004.

Houweling et al., 2005: Houweling, S., Hartmann, W., Aben, I., Schrijver, H., Skidmore, J., Roelofs, G.-J., and Breon, F.-M.: Evidence of systematic errors in SCIAMACHY-observed CO₂ due to aerosols, Atmos. Chem. Phys., 5, 3003–3013, 2005.

Houweling et al., 2015: Houweling, S., D. Baker, S. Basu, H. Boesch, A. Butz, F. Chevallier, F. Deng, E. J. Dlugokencky, L. Feng, A. Ganshin, O. Hasekamp, D. Jones, S. Maksyutov, J. Marshall, T. Oda, C.W. O'Dell, S. Oshchepkov, P. I. Palmer, P. Peylin, Z. Poussi, F. Reum, H. Takagi, Y. Yoshida, and R. Zhuravlev, [An intercomparison of inverse models for estimating sources and sinks of CO₂ using GOSAT measurements](#), J. Geophys. Res. Atmos., 120, 5253–5266, doi:10.1002/2014JD022962, 2015.

Hu et al., 2018: Hu, H., J. Landgraf, R. Detmers, T. Borsdorff, J. Aan de Brugh, I. Aben, A. Butz, O. Hasekamp, Toward Global Mapping of Methane With TROPOMI: First Results and Intersatellite Comparison to GOSAT, Geophys. Res. Lett., Vol. 45, Issue 8, 3682-3689, <https://doi.org/10.1002/2018GL077259>, 2018.

Hungerschofer et al., 2010: Hungerschofer, K., Breon, F.-M., Peylin, P., Chevallier, F., Rayner, P., Klonecki, A., Houweling, S., and Marshall, J., Evaluation of various observing systems for the global monitoring of CO₂ surface fluxes, Atmos. Chem. Phys., 10, 10503-10520, 2010.

IPCC, 2013: Climate Change 2013: The Physical Science Basis, Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Report on Climate Change, <http://www.ipcc.ch/report/ar5/wg1/>, 2013.



- JCGM, 2008:** JCGM/WG 1, Working Group 1 of the Joint Committee for Guides in Metrology, Evaluation of measurement data – Guide to the expression of uncertainty in measurement, http://www.bipm.org/utis/common/documents/jcgm/JCGM_100_2008_E.pdf, 2008.
- Kirschke et al., 2013:** Kirschke, S., Bousquet, P., Ciais, P., et al.: Three decades of global methane sources and sinks, *Nat. Geosci.*, 6, 813–823, doi:10.1038/ngeo1955, 2013.
- Kuze et al., 2009:** Kuze, A., Suto, H., Nakajima, M., and Hamazaki, T. (2009), Thermal and near infrared sensor for carbon observation Fourier-transform spectrometer on the Greenhouse Gases Observing Satellite for greenhouse gases monitoring, *Appl. Opt.*, 48, 6716–6733, 2009.
- Kuze et al., 2014:** Kuze, A., Taylor, T., Kataoka, F., Bruegge, C., Crisp, D., Harada, M., Helmlinger, M., Inoue, M., Kawakami, S., Kikuchi, N., Mitomi, Y., Murooka, J., Naitoh, M., O'Brien, D., O'Dell, C., Ohyama, H., Pollock, H., Schwandner, F., Shiomi, K., Suto, H., Takeda, T., Tanaka, T., Urabe, T., Yokota, T., and Yoshida, Y. (2014), Long-term vicarious calibration of GOSAT short-wave sensors: techniques for error reduction and new estimates of radiometric degradation factors, *IEEE T. Geosci. Remote*, 52, 3991–4004, doi:10.1109/TGRS.2013.2278696, 2014.
- Kuze et al., 2016:** Kuze, A., Suto, H., Shiomi, K., Kawakami, S., Tanaka, M., Ueda, Y., Deguchi, A., Yoshida, J., Yamamoto, Y., Kataoka, F., Taylor, T. E., and Buijs, H. L.: Update on GOSAT TANSO-FTS performance, operations, and data products after more than 6 years in space, *Atmos. Meas. Tech.*, 9, 2445–2461, doi:10.5194/amt-9-2445-2016, 2016.
- Lauer et al., 2017:** Lauer, A., V. Eyring, M. Righi, M. Buchwitz, P. Defourny, M. Evaldsson, P. Friedlingstein, R. de Jeu, G. de Leeuw, A. Loew, C. J. Merchant, B. Müller, T. Popp, M. Reuter, S. Sandven, D. Senfleben, M. Stengel, M. Van Roozendaal, S. Wenzel, U. Willén, Benchmarking CMIP5 models with a subset of ESA CCI Phase 2 data using the ESMValTool, *Remote Sensing of Environment*, DOI: 10.1016/j.rse.2016.12.027, in press, pp. 31, 2017.
- Machida et al. 2008:** Machida, T., Matsueda, H., Sawa, Y., Nakagawa, Y., Hirokuni, K., Kondo, N., Goto, K., Nakazawa, T., Ishikawa, K., and Ogawa, T.: Worldwide measurements of atmospheric CO₂ and other trace gas species using commercial airlines, *J. Atmos. Ocean. Tech.*, 25(10), 1744–1754, doi:10.1175/2008JTECHA1082.1, 2008.
- Massart et al., 2016:** Massart, S., A. Agustí-Panareda, J. Heymann, M. Buchwitz, F. Chevallier, M. Reuter, M. Hilker, J. P. Burrows, N. M. Deutscher, D. G. Feist, F. Hase, R. Sussmann, F. Desmet, M. K. Dubey, D. W. T. Griffith, R. Kivi, C. Petri, M. Schneider, V. A. Velasco, [Ability of the 4-D-Var analysis of the GOSAT BESD XCO₂ retrievals to characterize atmospheric CO₂ at large and synoptic scales](#), *Atmos. Chem. Phys.*, 16, 1653–1671, doi:10.5194/acp-16-1653-2016, 2016.
- Matsueda et al. 2008:** Matsueda, H., Machida, T., Sawa, Y., Nakagawa, Y., Hirokuni, K., Ikeda, H., Kondo, N., and Goto, K.: Evaluation of atmospheric CO₂ measurements from new flask air sampling of JAL airliner observation, *Pap. Meteorol. Geophys.*, 59, 1–17, 2008.
- McNorton et al., 2016:** McNorton, J., E. Gloor, C. Wilson, G. D. Hayman, N. Gedney, E. Comyn-Platt, T. Marthews, R. J. Parker, H. Boesch, and M. P. Chipperfield, [Role of regional wetland emissions in atmospheric methane variability](#), *Geophys. Res. Lett.*, 43, doi:10.1002/2016GL070649, 2016.
- Meirink et al., 2006:** Meirink, J.-F., Eskes, H. J., and Goede, A. P. H., Sensitivity analysis of methane emissions derived from SCIAMACHY observations through inverse modelling, *Atmos. Chem. Phys.*, 6, 1275–1292, 2006.



- Membrive et al. 2016:** Membrive, O., Crevoisier, C., Sweeney, C., Danis, F., Hertzog, A., Engel, A., Bönisch, H., and Picon, L.: AirCore-HR: A high resolution column sampling to enhance the vertical description of CH₄ and CO₂, *Atmos. Meas. Tech. Discuss.*, doi:10.5194/amt-2016-236, 2016.
- Miller et al., 2007:** Miller, C. E., Crisp, D., DeCola, P. L., et al.: Precision requirements for space-based XCO₂ data, *J. Geophys. Res.*, 112, D10314, doi:10.1029/2006JD007659, 2007.
- Nisbet et al., 2014:** Nisbet, E., Dlugokencky, E., and Bousquet, P.: Methane on the rise – again, *Science*, 343, 493–495, doi:10.1126/science.1247828, 2014.
- Pandey et al., 2016:** Pandey, S., S. Houweling, M. Krol, I. Aben, F. Chevallier, E. J. Dlugokencky, L. V. Gatti, E. Gloor, J. B. Miller, R. Detmers, T. Machida, T. Roeckmann, [Inverse modeling of GOSAT-retrieved ratios of total column CH₄ and CO₂ for 2009 and 2010](#), *Atmos. Chem. Phys.*, 16, 5043–5062, doi:10.5194/acp-16-5043-2016, 2016.
- Parker et al., 2011:** Parker, R., Boesch, H., Cogan, A., et al., Methane Observations from the Greenhouse gases Observing SATellite: Comparison to ground-based TCCON data and Model Calculations, *Geophys. Res. Lett.*, doi:10.1029/2011GL047871, 2011.
- Parker et al., 2016:** Parker, R. J., H. Boesch, M. J. Wooster, D. P. Moore, A. J. Webb, D. Gaveau, and D. Murdiyarso, [Atmospheric CH₄ and CO₂ enhancements and biomass burning emission ratios derived from satellite observations of the 2015 Indonesian fire plumes](#), *Atmos. Chem. Phys.*, *Atmos. Chem. Phys.*, 16, 10111–10131, doi:10.5194/acp-16-10111-2016, 2016.
- Pillai et al., 2016:** Pillai, D., Buchwitz, M., Gerbig, C., Koch, T., Reuter, M., Bovensmann, H., Marshall, J., and Burrows, J. P.: Tracking city CO₂ emissions from space using a high resolution inverse modeling approach: A case study for Berlin, Germany, *Atmos. Chem. Phys.*, 16, 9591–9610, doi:10.5194/acp-16-9591-2016, 2016.
- PUGS GHG, 2017:** Buchwitz, M., Aben, I., Anand, J., Armante, R., Boesch, H., Crevoisier, C., Detmers, R. G., Hasekamp, O. P., Reuter, M., Schneising-Weigel, O., Product User Guide and Specification (PUGS) – Main document, C3S project C3S_312a_Lot6_IUP-UB – Greenhouse Gases, version 1, 2017.
- PUGS GHG, 2018:** Buchwitz, M., Aben, I., Anand, J., Armante, R., Boesch, H., Crevoisier, C., Detmers, R. G., Hasekamp, O. P., Reuter, M., Schneising-Weigel, O., Product User Guide and Specification (PUGS) – Main document, C3S project C3S_312a_Lot6_IUP-UB – Greenhouse Gases, version 2, 2018.
- PUGS GHG, 2019:** Buchwitz, M., Aben, I., J., Armante, R., Boesch, H., Crevoisier, C., Di Noia, A., Hasekamp, O. P., Reuter, M., Schneising-Weigel, O., Wu, L., Product User Guide and Specification (PUGS) – Main document for Greenhouse Gas (GHG: CO₂ & CH₄) data set CDR 3 (2003–2018), C3S project C3S_312b_Lot2_DLR, v3.0, 2019.
- Rayner and O'Brien, 2001:** Rayner, P. J., and O'Brien, D.M.: The utility of remotely sensed CO₂ concentration data in surface inversions, *Geophys. Res. Lett.*, 28, 175–178, 2001.
- Reuter et al. 2011:** Reuter, M., Bovensmann, H., Buchwitz, M., Burrows, J. P., Connor, B. J., Deutscher, N. M., Griffith, D.W. T., Heymann, J., Keppel-Aleks, G., Messerschmidt, J., and et al.: Retrieval of atmospheric CO₂ with enhanced accuracy and precision from SCIAMACHY: Validation with FTS measurements and comparison with model results., *Journal of Geophysical Research*, 116, doi:10.1029/2010JD015047, URL <http://dx.doi.org/10.1029/2010JD015047>, 2011.



Reuter et al., 2013: Reuter, M. H. Bösch, H. Bovensmann, A. Bril, M. Buchwitz, A. Butz, J. P. Burrows, C. W. O'Dell, S. Guerlet, O. Hasekamp, J. Heymann, N. Kikuchi, S. Oshchepkov, R. Parker, S. Pfeifer, O. Schneising, T. Yokota, and Y. Yoshida, A joint effort to deliver satellite retrieved atmospheric CO₂ concentrations for surface flux inversions: The ensemble median algorithm EMMA, *Atmos. Chem. Phys.*, 13, 1771-1780, 2013.

Reuter et al., 2014a: Reuter, M., M. Buchwitz, A. Hilboll, A. Richter, O. Schneising, M. Hilker, J. Heymann, H. Bovensmann and J. P. Burrows, Decreasing emissions of NO_x relative to CO₂ in East Asia inferred from satellite observations, *Nature Geoscience*, 28 Sept. 2014, doi:10.1038/ngeo2257, pp.4, 2014.

Reuter et al., 2014b: Reuter, M., M. Buchwitz, M. Hilker, J. Heymann, O. Schneising, D. Pillai, H. Bovensmann, J. P. Burrows, H. Bösch, R. Parker, A. Butz, O. Hasekamp, C. W. O'Dell, Y. Yoshida, C. Gerbig, T. Nehrkorn, N. M. Deutscher, T. Warneke, J. Notholt, F. Hase, R. Kivi, R. Sussmann, T. Machida, H. Matsueda, and Y. Sawa, Satellite-inferred European carbon sink larger than expected, *Atmos. Chem. Phys.*, 14, 13739-13753, doi:10.5194/acp-14-13739-2014, 2014.

Reuter et al., 2016: Reuter, M.; Buchwitz, M.; Aben, I.; Boesch, H.; Butz, A.; Detmers, R.G.; Hasekamp, O.P.; Heymann, J.; Parker, R.; Schneising, O.; Somkuti, P., ESA Greenhouse Gases Climate Change Initiative (GHG_cci): Merged SCIAMACHY and GOSAT Level 3 gridded atmospheric column-average carbon dioxide (XCO₂) product in Obs4MIPs format. Centre for Environmental Data Analysis, 10 October 2016, doi:10.5285/3FAE8371-0CBB-4B21-9EA6-7A1FC293C4A2, link: http://www.esa-ghg-cci.org/?q=webfm_send/330, pp. 11, 2016.

Reuter et al., 2017: Reuter, M., M. Buchwitz, M. Hilker, J. Heymann, H. Bovensmann, J. Burrows, S. Houweling, Y. Liu, R. Nassar, F. Chevallier, P. Ciais, J. Marshall, and M. Reichstein, 2016: How much CO₂ is taken up by the European terrestrial biosphere ?, *Bull. Amer. Meteor. Soc.* doi:10.1175/BAMS-D-15-00310.1, 24 April 2017, 665-671, 2017.

Reuter et al., 2017a: Reuter, M.; Buchwitz, M.; Aben, I.; Boesch, H.; Butz, A.; Detmers, R.G.; Hasekamp, O.P.; Heymann, J.; Parker, R.; Schneising, O.; Somkuti, P., ESA Greenhouse Gases Climate Change Initiative (GHG-CCI): Merged SCIAMACHY and GOSAT Level 3 gridded atmospheric column-average carbon dioxide (XCO₂) product in Obs4MIPs format version 2 (CRDP#4), Technical Note, link: http://www.esa-ghg-cci.org/?q=webfm_send/348, pp. 11, 1 February 2017, 2017.

Reuter et al., 2019: Reuter, M., Buchwitz, M., Schneising, O., Noel, S., Bovensmann, H., Burrows, J. P., Boesch, H., Di Noia, A., Anand, J., Parker, R. J., Somkuti, P., Wu, L., Hasekamp, O. P., Aben, I., Kuze, A., Suto, H., Shiomi, K., Yoshida, Y., Morino, I., Crisp, D., O'Dell, C., Notholt, J., Petri, C., Warneke, T., Velasco, V., Deutscher, N. M., Griffith, D. W. T., Kivi, R., Pollard, D., Hase, F., Sussmann, R., Te, Y. V., Strong, K., Roche, S., Sha, M. K., De Maziere, M., Feist, D. G., Iraci, L. T., Roehl, C., Retscher, C., and Schepers, D.: Ensemble-based satellite-derived carbon dioxide and methane column-averaged dry-air mole fraction data sets (2003-2018) for carbon and climate applications, *Atmos. Meas. Tech. Discuss.*, <https://doi.org/10.5194/amt-2019-398>, in review, 2019.

Rodgers, 2000: Rodgers C. D.: *Inverse Methods for Atmospheric Sounding: Theory and Practice*, World Scientific Publishing, 2000.

Ross et al., 2013: Ross, A. N., Wooster, M. J., Boesch, H., Parker, R., First satellite measurements of carbon dioxide and methane emission ratios in wildfire plumes, *Geophys. Res. Lett.*, 40, 1-5, doi:10.1002/grl.50733, 2013.



- Schaefer et al., 2016:** Schaefer, H., Mikaloff Fletcher, S. E., Veidt, C., Lassey, K. R., Brailsford, G. W., Bromley, T. M., Dlugokencky, E. J., Michel, S. E., Miller, J. B., Levin, I., Lowe, D. C., Martin, R. J., Vaughn, B. H., and White, J. W. C.: A 21st-century shift from fossil-fuel to biogenic methane emissions indicated by $^{13}\text{CH}_4$, *Science*, Vol. 352, Issue 6281, pp. 80-84, doi 10.1126/science.aad2705, 2016.
- Shindell et al., 2013:** Shindell, D. T., Pechony, O., Voulgarakis, A., et al. (2013), Interactive ozone and methane chemistry in GISS-E2 historical and future climate simulations, *Atmos. Chem. Phys.*, 13, 2653–2689, doi:10.5194/acp-13-2653-2013, 2013.
- Schepers et al., 2012:** Schepers, D., Guerlet, S., Butz, A., Landgraf, J., Frankenberg, C., Hasekamp, O., Blavier, J.-F., Deutscher, N. M., Griffith, D. W. T., Hase, F., Kyro, E., Morino, I., Sherlock, V., Sussmann, R., Aben, I. (2012), Methane retrievals from Greenhouse Gases Observing Satellite (GOSAT) shortwave infrared measurements: Performance comparison of proxy and physics retrieval algorithms, *J. Geophys. Res.*, 117, D10307, doi:10.1029/2012JD017549, 2012.
- Schneising et al., 2011:** Schneising, O., Buchwitz, M., Reuter, M., et al., Long-term analysis of carbon dioxide and methane column-averaged mole fractions retrieved from SCIAMACHY, *Atmos. Chem. Phys.*, 11, 2881-2892, 2011.
- Schneising et al., 2014a:** Schneising, O., Reuter, M., Buchwitz, M., Heymann, J., Bovensmann, H., and Burrows, J. P., Terrestrial carbon sink observed from space: variation of growth rates and seasonal cycle amplitudes in response to interannual surface temperature variability, *Atmos. Chem. Phys.*, 14, 133-141, 2014.
- Schneising et al., 2014b:** Schneising, O., Burrows, J. P., Dickerson, R. R., Buchwitz, M., Reuter, M., Bovensmann, H., Remote sensing of fugitive methane emissions from oil and gas production in North American tight geologic formations, *Earth's Future*, 2, DOI: 10.1002/2014EF000265, pp. 11, 2014.
- TRD GHG, 2017:** Buchwitz, M., Aben, I., Anand, J., Armante, R., Boesch, H., Crevoisier, C., Detmers, R. G., Hasekamp, O. P., Reuter, M., Schneising-Weigel, O., Target Requirement Document, Copernicus Climate Change Service (C3S) project on satellite-derived Essential Climate Variable (ECV) Greenhouse Gases (CO_2 and CH_4) data products (project C3S_312a_Lot6), Version 1.3, 20-October-2017, pp. 53, 2017.
- Turner et al., 2015:** Turner, A. J., D. J. Jacob, K. J. Wecht, J. D. Maasakkers, S. C. Biraud, H. Boesch, K. W. Bowman, N. M. Deutscher, M. K. Dubey, D. W. T. Griffith, F. Hase, A. Kuze, J. Notholt, H. Ohyama, R. Parker, V. H. Payne, R. Sussmann, V. A. Velazco, T. Warneke, P. O. Wennberg, and D. Wunch, Estimating global and North American methane emissions with high spatial resolution using GOSAT satellite data, *Atmos. Chem. Phys.*, 15, 7049-7069, doi:10.5194/acp-15-7049-2015, 2015.
- Turner et al., 2016:** Turner, A. J., D. J. Jacob, J. Benmergui, S. C. Wofsy, J. D. Maasakkers, A. Butz, O. Hasekamp, and S. C. Biraud, A large increase in U.S. methane emissions over the past decade inferred from satellite data and surface observations, *Geophys. Res. Lett.*, 43, 2218–2224, doi:10.1002/2016GL067987, 2016.
- Veefkind et al. 2012:** Veefkind, J. P., Aben, I., McMullan, K., Förster, H., De Vries, J., Otter, G., Claas, J., Eskes, H. J., De Haan, J. F., Kleipool, Q., Van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P. F.:



TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications. *Rem. Sens. Environment*, 120:70–83, 2012.

Velazco et al. 2011: Velazco, V. A., Buchwitz, M., Bovensmann, H., Reuter, M., Schneising, O., Heymann, J., Krings, T., Gerilowski, K., and Burrows, J. P.: Towards space based verification of CO₂ emissions from strong localized sources: fossil fuel power plant emissions as seen by a CarbonSat constellation, *Atmos. Meas. Tech.*, 4, 2809–2822, 2011.

Wenzel et al., 2016: Wenzel, S., Cox, P. M., Eyring, V., Friedlingstein, P.: Projected land photosynthesis constrained by changes in the seasonal cycle of atmospheric CO₂, *Nature*, 499–501, doi:10.1038/nature18722, 2016.

Wofsy et al. 2012: Wofsy, S. C., Daube, B. C., Jimenez, R., et al.: HIPPO Merged 10-second Meteorology, Atmospheric Chemistry, Aerosol Data (R 20121129), Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA, http://dx.doi.org/10.3334/CDIAC/hippo_010 (Release 29 November 2012), 2012.

Wunch et al. 2010: Wunch, D., Toon, G. C., Wennberg, P. O., Wofsy, S. C., Stephens, B. B., Fischer, M. L., Uchino, O., Abshire, J. B., Bernath, P., Biraud, S. C., Blavier, J.-F. L., Boone, C., Bowman, K. P., Browell, E. V., Campos, T., Connor, B. J., Daube, B. C., Deutscher, N. M., Diao, M., Elkins, J. W., Gerbig, C., Gottlieb, E., Griffith, D. W. T., Hurst, D. F., Jiménez, R., Keppel-Aleks, G., Kort, E. A., Macatangay, R., Machida, T., Matsueda, H., Moore, F., Morino, I., Park, S., Robinson, J., Roehl, C. M., Sawa, Y., Sherlock, V., Sweeney, C., Tanaka, T., and Zondlo, M. A.: Calibration of the Total Carbon Column Observing Network using aircraft profile data, *Atmospheric Measurement Techniques*, 3, 1351–1362, doi:10.5194/amt-3-1351-2010, URL <http://www.atmos-meas-tech.net/3/1351/2010/>, 2010.

Wunch et al. 2011: Wunch, D., Toon, G. C., Blavier, J.-F. L., Washenfelder, R. A., Notholt, J., Connor, B. J., Griffith, D. W. T., Sherlock, V., and Wennberg, P. O.: The Total Carbon Column Observing Network (TCCON), *Philosophical Transactions of the Royal Society of London, Series A: Mathematical, Physical and Engineering Sciences*, 369, 2087–2112, doi:10.1098/rsta.2010.0240, 2011.

Wunch et al. 2015: Wunch, D., Toon, G.C., Sherlock, V., Deutscher, N.M., Liu, X., Feist, D.G., Wennberg, P.O., The Total Carbon Column Observing Network's GGG2014 Data Version. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA (available at: doi:10.14291/tccon.ggg2014.documentation.R0/1221662), 2015.

Yoshida et al. 2013: Yoshida, Y., Kikuchi, N., Morino, I., Uchino, O., Oshchepkov, S., Bril, A., Saeki, T., Schutgens, N., Toon, G. C., Wunch, D., Roehl, C. M., Wennberg, P. O., Griffith, D. W. T., Deutscher, N. M., Warneke, T., Notholt, J., Robinson, J., Sherlock, V., Connor, B., Rettinger, M., Sussmann, R., Ahonen, P., Heikkinen, P., Kyrö, E., Mendonca, J., Strong, K., Hase, F., Dohe, S., and Yokota, T.: Improvement of the retrieval algorithm for GOSAT SWIR XCO₂ and XCH₄ and their validation using TCCON data, *Atmos. Meas. Tech.*, 6, 1533–1547, doi:10.5194/amt-6-1533-2013, 2013.



6. Acknowledgement

We acknowledge previous funding by the European Space Agency (ESA) via Climate Change Initiative (CCI) project GHG-CCI. This funding significantly enhanced the quality of the retrieval algorithms and related documentation. This resulted in more mature data products as needed for an operational project such as the Copernicus Climate Change Service (C3S). We also acknowledge the availability of GOSAT data products via the ESA GOSAT Third Party Mission (TPM) archive.

We are also very grateful to the GOSAT team in Japan comprising the Japan Aerospace Exploration Agency (JAXA), the National Institute for Environmental Studies (NIES), and the Ministry of the Environment (MOE) for providing access to the GOSAT Level 1 and Level 2 data products via the GOSAT Data Archive Service (GDAS) hosted by NIES.

We also acknowledge the availability of OCO-2 Level 1 and Level 2 (XCO₂) data products from NASA, which have been used for the generation on the XCO₂_EMMA and XCO₂_OBS4MIPS products. These products also include OCO-2 XCO₂ retrieved at Univ. Bremen with the FOCAL algorithm. The FOCAL activities would not have been possible without funding from University of Bremen, from the EU H2020 projects CHE (grant agreement ID: 776186) and VERIFY (Grant agreement ID: 776810) and from ESA via project GHG-CCI+.

Last but not least we acknowledge the availability of TCCON data via the TCCON data archive (<https://tccondata.org/>).



7. List of ANNEXes

The ANNEXes to this main document are the following ANNEXes A – E valid for data set CDR 3 (2003-2018):

7.1 ANNEX A: PQAR for products CO₂_GOS_OCFP, CH₄_GOS_OCFP, CH₄_OCPR

Describes the validation of the GOSAT XCO₂ and XCH₄ Level 2 products generated by University of Leicester, UK.

7.2 ANNEX B: PQAR for products CO₂_GOS_SRFP, CH₄_GOS_SRFP

Describes the validation of the GOSAT XCO₂ and XCH₄ Full Physics (FP) Level 2 products generated by SRON, The Netherlands.

7.3 ANNEX C: PQAR for product CH₄_GOS_SRPR

Describes the validation of the GOSAT XCH₄ Proxy (PR) Level 2 product generated by SRON, The Netherlands.

7.4 ANNEX D: PQAR for products XCO₂_EMMA, XCH₄_EMMA, XCO₂_OBS4MIPS, XCH₄_OBS4MIPS

Describes the validation of the multi-sensor multi-algorithms merged XCO₂ and XCH₄ Level 2 and 3 products generated by University of Bremen, Germany.

7.5 ANNEX E: PQAR for IASI CO₂ and CH₄ and AIRS CO₂ products

Describes the validation of the mid-tropospheric CO₂ and CH₄ products from the IASI instrument series and AIRS generated by LMD/CNRS, France.

These ANNEXes and the corresponding data products are / will be available via the Copernicus Climate Data Store (CDS):

<https://cds.climate.copernicus.eu/#!/home>

See also Copernicus Climate Change Service (C3S):

<https://climate.copernicus.eu/>



ECMWF - Shinfield Park, Reading RG2 9AX, UK

Contact: info@copernicus-climate.eu