

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 1
		Version 3 Final
		6 June 2014

ESA Climate Change Initiative (CCI)

Product Specification Document (PSD)

for the Essential Climate Variable (ECV)

Greenhouse Gases (GHG)

- Description of Common Parameters for core (ECA) products -

Written by: GHG-CCI project team

with contributions from NASA/ACOS/OCO-2 team members

Lead authors: M. Buchwitz, IUP, Univ. Bremen, and R. Detmers, SRON Netherlands
Institute for Space Research

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 2
		Version 3 Final
		6 June 2014

Change log:

Version Nr.	Date	Status	Reason for change
Version 1 - Final	February 28, 2011	Final PSDv1	1 st version
Version 2 - Final	March 12, 2012	Final PSDv2	2 nd version
<i>Version 3 – Draft 1</i>	<i>May 2, 2014</i>	<i>Draft submitted to project team for comments</i>	<i>To describe improved data format for core products (ECA products)</i>
<i>Version 3 – Draft 2</i>	<i>May 13, 2014</i>	<i>Improved draft</i>	<i>Inputs from co-authors added</i>
<i>Version 3 – Draft 3</i>	<i>May 16, 2014</i>	<i>Improved draft</i>	<i>Inputs from co-authors added</i>
<i>Version 3 – Draft 4</i>	<i>May 19, 2014</i>	<i>Improved draft</i>	<i>Inputs from co-authors added</i>
<i>Version 3 – Draft 5</i>	<i>May 22, 2014</i>	<i>Improved Draft</i>	<i>Inputs from co-authors added</i>
<i>Version 3 – Draft 6</i>	<i>May 27, 2014</i>	<i>Improved Draft</i>	<i>Inputs from co-authors added</i>
Version 3 – Final	June 6, 2014	Final PSDv3	Inputs from SRON added

Authors:

- GHG-CCI project team members: H. Boesch (Univ. Leicester), M. Buchwitz (IUP, Univ. Bremen), F. Chevallier (LSCE), R. Detmers (SRON), B. Dils (BIRA-IASB), C. Frankenberg (JPL, USA), O. Hasekamp (SRON), S. Houweling (SRON), G. Lichtenberg (DLR), J. Marshall (MPI-BGC), R. Parker (Univ. Leicester), M. Reuter (IUP, Univ. Bremen), O. Schneising (IUP, Univ. Bremen)
- NASA/ACOS/OCO-2 team members: C. O'Dell (Colorado State University, USA), C. Frankenberg (JPL, USA)

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 3
		Version 3 Final
		6 June 2014

Table of Contents

1	Overview.....	4
2	Instruments.....	7
2.1	SCIAMACHY/ENVISAT	7
2.2	TANSO/GOSAT	7
3	GHG-CCI core (ECA) products.....	8
3.1	File format and file naming convention	11
3.2	Common Parameters	12
3.3	How to use the Averaging Kernels (AKs)?	17
3.3.1	Introduction.....	17
3.3.2	Averaging kernel (AK) related parameters	19
3.3.3	How to use layer-based AKs?	20
3.3.4	How to use level-based AKs?	24
3.4	Algorithm-Specific Parameters	26
4	References	27
5	Acronyms.....	31

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 4
		Version 3 Final
		6 June 2014

1 Overview

This document is Product Specification Document (PSD) Version 3 (PSDv3), which is a deliverable of the ESA project GHG-CCI (<http://www.esa-ghg-cci.org>).

This version of the PSD incorporates the user requirements as described in the GHG-CCI User Requirements Document (URD) /**Buchwitz et al., 2011**/ as well as the Data Standards Requirements for CCI Data Producers /**Bennett and James, 2013**/.

The GHG-CCI project aims at generating and delivering the Essential Climate Variable (ECV) Greenhouse Gases (GHG) in-line with GCOS (Global Climate Observing System) requirements /**GCOS 2011**/, GHG-CCI addresses the ECV "Greenhouse Gases" defined as by GCOS follows: "Retrievals of greenhouse gases, such as CO₂ and CH₄, of sufficient quality to estimate regional sources and sinks".

Two kinds of algorithms are used within GHG-CCI to generate satellite-derived atmospheric carbon dioxide (CO₂) and methane (CH₄) data products:

- **ECV Core Algorithms (ECAs):** Algorithms to retrieve column-averaged dry air mole fractions of CO₂ and CH₄ from (currently) SCIAMACHY/ENVISAT and TANSO/GOSAT, i.e., XCO₂ (in ppm) and XCH₄ (in ppb). These instruments are used as "core instruments" because their measurements are sensitive to the lowest atmospheric layer and therefore provide information on the regional sources and sinks of CO₂ and CH₄. Please see **Table 1** for an overview about all products generated with ECAs.
- **Additional Constraints Algorithms (ACAs):** Algorithms to retrieve CO₂ and/or CH₄ information on layers above the Planetary Boundary Layer (PBL) such as IASI and MIPAS. This can provide additional constraints for inverse modelling of CO₂ and/or CH₄ regional fluxes by providing atmospheric CO₂ and/or CH₄ information for "upper layers".

In this document only the GHG-CCI ECA products are described. Furthermore, this PSD only describes the "common parameters" of all ECA products, i.e., those parameters, which are common to all ECA products. The individual ECA products typically have additional, algorithm specific parameters. **These specific parameters are not described in this document but are / will be described in dedicated Product User Guides (PUGs)** (see GHG-CCI website).

ACA product formats will not be described in this document but in dedicated PUGs (see GHG-CCI website).

Currently GHG-CCI is in Phase 2 (2014-2016). Previous versions of this document (PSDv1 and PSDv2) describe the products generated during Phase 1 of GHG-CCI (2010-2013). In particular PSDv2 describes the format of the first version of the GHG-CCI Climate Research Data Package (CRDP#1).

This document (**PSDv3**) describes the improved format for GHG-CCI core (=ECA) products to be used for the next version of the CRDP, which is **CRDP#2** (planned to be available in October 2014).

These products plus corresponding documentation will be available for download via the same website as used for all GHG-CCI products (including previous versions, i.e., CRDP#1):

<http://www.esa-ghg-cci.org/> -> **CRDP (Data)**

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 5
		Version 3 Final
		6 June 2014

An overview of all GHG-CCI data products (ECA and ACA products) including data access info etc. is given on the **GHG-CCI main data products website**:

<http://www.esa-ghg-cci.org/> -> CRDP (Data) -> GHG-CCI Data Products Main Website

For a short overview of all ECA products please see **Table 1**. As can be seen, **Table 1** lists baseline and alternative algorithms.

The purpose of the alternative products is to “challenge the baseline” but also to provide alternative products for ensemble applications. If it will be shown in the future that the alternative algorithm/product is better than the baseline, the baseline will change.

Focus of GHG-CCI Phase 2 (2014-2016) will be on the **baseline products** (full documentation, product format meeting all requirements, data available in time as planned, entire planned time series processed, etc.); the **alternative and EMMA products** will also be further developed but with lower priority. For the actual status please visit the GHG-CCI main data products website (see above).

The main purpose of this document is to describe the new data format for ECA products (limited to common parameters) to be used for the CRDP#2 (planned release October 2014). All baseline products will be in this new format but not necessarily all alternative products and the EMMA product (although GHG-CCI will aim at achieving this).

Product format changes in the future (if any) will be reflected in future updates to this document.

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 6
		Version 3 Final
		6 June 2014

ECA products: GHG-CCI Core Products						
Product Main ID	Product	Product details	Sensor	Algorithm		Comment
				Baseline	Alternative (*)	
XCO2_SCI	XCO ₂	Level 2, CO ₂ column-averaged dry-air mole fraction (in ppm)	SCIAMACHY / ENVISAT	BESD	WFMD	
XCO2_GOS	XCO ₂	Level 2, CO ₂ column-averaged dry-air mole fraction (in ppm)	TANSO / GOSAT	OCFP (UoL-FP), SRFP (RemoTeC) (baseline not yet defined (#))		
XCO2_EMMA	XCO ₂	Level 2, CO ₂ column-averaged dry-air mole fraction (in ppm)	SCIAMACHY and GOSAT	EMMA (*) Experimental product currently covering mid 2009 – mid 2010		Merged ensemble product
XCH4_SCI	XCH ₄	Level 2, CH ₄ column-averaged dry-air mole fraction (in ppb)	SCIAMACHY / ENVISAT	WFMD, IMAP (baseline not yet defined (#))		Note: degradation issues after October 2005 (most severe after mid-2010)
XCH4_GOS_FP	XCH ₄	Level 2, CH ₄ column-averaged dry-air mole fraction (in ppb)	TANSO / GOSAT	SRFP (RemoTeC)	OCFP (UoL-FP)	"Full Physics" (FP) algorithm product
XCH4_GOS_PR	XCH ₄	Level 2, CH ₄ column-averaged dry-air mole fraction (in ppb)	TANSO / GOSAT	OCPR (UoL-PR)	SRPR (RemoTeC)	"ProxY" (PR) algorithm product

Table 1: Overview GHG-CCI ECV Core Algorithm (ECA) products. For details please see **Table 2** and **Table 3**. (*) See main text for potential restrictions on alternative algorithms/products and EMMA. (#) Evaluation during GHG-CCI Phase 1 revealed some significant differences but it was not possible to clearly identify which algorithm/product is better (please see GHG-CCI Algorithm Selection Report /GHG-CCI ASR/ and Product Validation and Intercomparison Report /GHG-CCI PVIRv2/ for details).

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 7
		Version 3 Final
		6 June 2014

2 Instruments

This section provides a brief description of the two core GHG-CCI instruments, SCIAMACHY on ENVISAT and TANSO-FTS onboard GOSAT.

2.1 SCIAMACHY/ENVISAT

The SCanning Imaging Absorption spectroMeter for Atmospheric CHartographY (SCIAMACHY, 2002-2012) was a passive remote sensing spectrometer, which observed backscattered, reflected, transmitted or emitted radiation from the atmosphere and Earth's surface, in the wavelength range between approximately 240 and 2380 nm. SCIAMACHY was part of the scientific payload of the ENVISAT satellite, which was launched on 1st of March 2002. A detailed instrument description is given in **/Bovensmann et al., 1999/**. Until 2009, when GOSAT was launched, SCIAMACHY was the only satellite instrument allowing for the measurement of the atmospheric carbon dioxide and methane concentration with significant sensitivity down to the Earth's surface, where (most of) the sources and sinks of CO₂ and CH₄ are located.

2.2 TANSO/GOSAT

The Japanese Greenhouse gases Observing SATellite (GOSAT) **/Kuze et al., 2009/** was launched on 23rd January 2009 by JAXA, the Japanese Space Agency. GOSAT provides the first dedicated global measurements of total column CO₂ and CH₄ from its Short-Wave-Infrared (SWIR) bands **/Yoshida et al., 2013/**. GOSAT is equipped with two instruments, the Thermal And Near Infrared Sensor for carbon Observations - Fourier Transform Spectrometer (TANSO-FTS) as well as a dedicated Cloud and Aerosol Imager (TANSO-CAI).

The TANSO-FTS instrument has four spectral bands with a high spectral resolution 0.3 cm⁻¹, three of which operate in the SWIR at around 0.76, 1.6 and 2.0 µm providing sensitivity to near-surface CO₂ and CH₄ concentration variations with the fourth channel operating in the thermal infrared between 5.5 and 14.3 µm providing mid-tropospheric sensitivity **/Saitoh et al., 2009/**.

The measurement strategy of TANSO-FTS is optimised for the characterisation of continental-scale CO₂ and CH₄ sources and sinks with the aim of achieving a 0.3%-1% relative accuracy for 3-month averages of CO₂ at a 100-1000 km spatial resolution **/Kuze et al., 2009/**. The aim for CH₄ is to achieve an accuracy of better than 2% on the same spatial and temporal scales. In order to achieve this, TANSO-FTS utilises a pointing mirror to perform off-nadir measurements at the same location on each 3-day repeat cycle. The pointing mirror allows TANSO-FTS to observe up to ±35° across track and ±20° along-track. These measurements nominally consist of 5 across track points spaced ~100 km apart (although measurements are possible with 1, 3, 5, 7 or 9 across track points) with a ground footprint diameter of approximately 10.5 km and a 4 second exposure duration. Whilst the majority of data is limited to measurements over land, where the surface reflectance is high, TANSO-FTS also observes in sun-glint mode over the ocean within ±20° of the sub-solar latitude.

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 8
		Version 3 Final
		6 June 2014

3 GHG-CCI core (ECA) products

ECV Core Algorithms (ECAs) are algorithms to retrieve XCO₂ and/or XCH₄ from SCIAMACHY and/or TANSO-FTS. Several ECAs are further developed and assessed within GHG-CCI. At present at least two different ECAs (baseline and alternative algorithm) are used for each of the main (core) data products (i.e., XCO₂ and XCH₄ from SCIAMACHY and GOSAT). For details please see GHG-CCI website:

<http://www.esa-ghg-cci.org/> -> CRDP (Data) -> GHG-CCI Data Products Main Website

Table 2 provides an overview of the XCO₂ ECAs. **Table 3** shows the corresponding information for XCH₄.

The data products generated with ECAs contain a common set of parameters (described in this document) and algorithm specific parameters (not described in this document but in the corresponding Product User Guides (PUGs)).

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 9
		Version 3 Final
		6 June 2014

XCO ₂ Core Product Identifier	Algorithm	Institute	Technique	Reference
CO2_SCI_BESD	BESD: Bremen optimal ESTimation DOAS SCIAMACHY XCO ₂ algorithm	IUP, Univ. Bremen, Germany	Optimal Estimation	Reuter et al., 2010, 2011
CO2_SCI_WFMD (*)	WFM-DOAS (WFMD): Weighting Function Modified DOAS SCIAMACHY XCO ₂ algorithm	IUP, Univ. Bremen, Germany	Least-squares DOAS	Schneising et al., 2011, 2012
CO2_GOS_OCFP	OCFP or UoL-FP: University of Leicester's (OCO-based) GOSAT XCO ₂ algorithm	University of Leicester (UoL), UK	Optimal Estimation	Boesch et al., 2006, Connor et al., 2008
CO2_GOS_SRFP	SRFP or RemoTeC: SRON/KIT GOSAT XCO ₂ algorithm	SRON, Netherlands, in cooperation with KIT, Germany	Phillips-Tikhonov regularization	Butz et al., 2009, 2010; Guerlet et al., 2013a, 2013b
CO2_EMMA (*)	Ensemble algorithm	Lead: IUP, Univ. Bremen, Germany	Level 2 merging algorithm	Reuter et al., 2013

Table 2: Overview GHG-CCI ECV Core Algorithms (ECAs) for XCO₂. (*) Alternative algorithm/product with potential limitations (see **Sect. 1**).

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 10
		Version 3 Final
		6 June 2014

XCH ₄ Core Product Identifier	Algorithm	Institute	Technique	Reference
CH4_SCI_WFMD	WFM-DOAS (WFMD): Weighting Function Modified DOAS SCIAMACHY XCH ₄ algorithm	IUP, Univ. Bremen, Germany	Least-squares DOAS; PROxy (PR) method using CO ₂	Schneising et al., 2011, 2012
CH4_SCI_IMAP	IMAP-DOAS (IMAP): Iterative Maximum A Posteriori DOAS SCIAMACHY XCH ₄ algorithm	SRON, Netherlands, in cooperation with JPL, USA	Iterative maximum a posteriori DOAS method; PROxy (PR) method using CO ₂	Frankenberg et al., 2005, 2011
CH4_GOS_OCPR	OCPR or UoL-PR: University of Leicester's (OCO- based) GOSAT XCH ₄ proxy algorithm	University of Leicester (UoL), UK	Optimal Estimation; PROxy (PR) method using CO ₂	Parker et al., 2011
CH4_GOS_SRF	SRFP or RemoTeC: SRON/KIT GOSAT XCH ₄ full physics algorithm	SRON, Netherlands, in cooperation with KIT, Germany	Phillips-Tikhonov regularization; Full Physics (FP) method	Butz et al., 2009, 2010; Guerlet et al., 2013a
CH4_GOS_OCFP (*)	OCFP or UoL-PR: University of Leicester's (OCO- based) GOSAT XCH ₄ full physics algorithm	University of Leicester (UoL), UK	Optimal Estimation; Full Physics (FP) method	Parker et al., 2011
CH4_GOS_SRPR (*)	SRPR or RemoTeC: SRON/KIT GOSAT XCH ₄ proxy algorithm	SRON, Netherlands, in cooperation with KIT, Germany	Phillips-Tikhonov regularization; PROxy (PR) method using CO ₂	Butz et al., 2009, 2010

Table 3: Overview GHG-CCI ECV Core Algorithms (ECAs) for XCH₄. (*) Alternative algorithm/product with potential limitations (see **Sect. 1**).

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 11
		Version 3 Final
		6 June 2014

3.1 File format and file naming convention

User requirements for GHG-CCI core products are given in the GHG-CCI URDv1 **/Buchwitz et al., 2011/** and specific data format requirements are given in Data Standards Requirements for CCI Data Producers **/Bennett and James, 2013/**. Both documents have been considered for this version of the PSD.

The required file format is described in **/Bennett and James, 2013/**, requirement R-1:

<i>All CCI projects shall produce data according to the following minimum requirements for data products:</i>
CCI projects shall produce data according to the CCI Data Standards Requirements: • Produced in netCDF-4 (classic) format (see http://www.unidata.ucar.edu/software/netcdf/) • Conform with the CF (Climate and Forecasting) convention 3 (currently version 1.6) • CF standard names used for the main variables • Include the global attributes listed in /Bennett and James, 2013/
<i>Note: For netCDF-4 (classic) format, the mode must be set to NC_CLASSIC_MODEL, but compression can be invoked. If netCDF-4 cannot be used (e.g. user tools not yet mature enough), data may be produced in netCDF-3 format.</i>
<i>Files can be checked for compliance with the CF convention using the on-line checking tool : http://cfpcmdi.llnl.gov/conformance/compliance-checker</i>

These requirements have been considered for the products described in this document.

In particular, the netCDF-4 (classic) format is used and the products are in-line with CF (Climate and Forecasting) convention 3.

Concerning the requirement to use CF standard names for the main variables: The process to register "XCO₂" and "XCH₄" as standard names has been initiated (but at the time of writing of this document the registration process was still ongoing). Details on the file content are given in the following sections.

As required **/Bennett and James, 2013/** the following convention is used:

The **directory structure** consists of archive-root (ESACCI/GHG), sensor (satellite or satellite instrument), product (for example BESD_CO2 or SRFP_CH4), version (product version) and time (YYYY, i.e., year only).

Examples:

./ESACCI/GHG/SCIAMACHY/BESD_CO2/v02.00.08/2002/

./ESACCI/GHG/GOSAT/SRFP_CH4/v2.11/2012/

The **file names** consist of ESACCI, CCI project (GHG), processing level (L2), product type (CO2 / CH4), sensor (e.g., SCIAMACHY), algorithm (e.g., BESD or SRFP), date (YYYYMMDD), file version (fv#) and file name extension (.nc), separated by hyphens ("-").

Examples:

ESACCI-GHG-L2-CO2-SCIAMACHY-BESD-20021216-fv1.nc

ESACCI-GHG-L2-CH4-GOSAT-SRFP-20120909-fv1.nc

Each *.nc product file corresponds to one day of satellite observations.

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 12
		Version 3 Final
		6 June 2014

3.2 Common Parameters

The users of the GHG ECV data products, as represented by the GHG-CCI Climate Research Group (CRG), need data products which contain all the information required for surface flux inverse modelling such as retrieved XCO_2 and XCH_4 values for individual ground pixels, their errors, corresponding averaging kernels, *a-priori* profiles, etc. The GHG-CCI data products are (therefore) Level 2 products. For details please see the GHG-CCI User Guide /**Buchwitz et al., 2011**/).

In order to provide the desired information and to be in agreement with the CCI Data Standard Requirements /**Bennett and James, 2013**/ it is required that all ECA products contain at least the parameters listed in the tables below (**Table 4** and **Table 5**).

In order to use our products as easily as possible we have aimed at harmonizing our various products. The goal was to make sure that users can easily switch from one product to another. This has been achieved for all products and parameters with the exception of the averaging kernels and related parameters. These parameters are closely related to retrieval algorithm specific characteristics and require special consideration by the users of our products as is explained in detail in **Section 3.3**.

Dimensions in **Table 4** and **Table 5** are defined as follows:

- ***n***: number of satellite observations (ground pixels) (per file, i.e., for the given day of observations)
- For Averaging Kernel (AK) and related parameters:
 - As explained in **Section 3.3**, the AK and related parameters are provided for “layer-based AKs” and “level-based AKs”
 - For layer-based AK ***m*** is the number of layers which are defined by **$k = m+1$** pressure levels.
 - For level-based AK only levels are used. Here all parameters have the same number of elements, namely ***m*** levels. Here the number of pressure levels is also ***m*** (i.e., **$k = m$**).

Table 4 and **Table 5** present an overview about all common parameters including a short description of each parameter. A detailed description is given afterwards.

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 13
		Version 3 Final
		6 June 2014

Name	Type	Dimensions	Units	Short Description
Common parameters for XCO₂ products:				
xco2	Float	n	micromol per mol, abbreviated ppm, i.e., 10 ⁻⁶	Retrieved column-averaged dry-air mole fraction of atmospheric carbon dioxide (XCO ₂) in ppm.
xco2_uncertainty	Float	n	micromol per mol, abbreviated ppm, i.e., 10 ⁻⁶	Statistical uncertainty of XCO ₂ in ppm (1-sigma).
xco2_averaging_kernel	Float	n x m	[-]	XCO ₂ averaging kernel (a profile = vector for each single observation). Quantifies the altitude sensitivity of the XCO ₂ retrieval. (see Sect. 3.3)
co2_profile_apriori	Float	n x m	micromol per mol, abbreviated ppm, i.e., 10 ⁻⁶	A priori mole fraction profile of atmospheric CO ₂ in ppm. (see Sect. 3.3)
xco2_quality_flag	Byte	n	[-]	Quality flag for XCO ₂ retrieval. 0=good.
Common parameters for XCH₄ products:				
xch4	Float	n	nanomol per mol, abbreviated ppb, i.e., 10 ⁻⁹	Retrieved column-averaged dry-air mole fraction of atmospheric methane (XCH ₄) in ppb.
xch4_uncertainty	Float	n	nanomol per mol, abbreviated ppb, i.e., 10 ⁻⁹	Statistical uncertainty of XCH ₄ in ppb (1-sigma)
xch4_averaging_kernel	Float	n x m	[-]	XCH ₄ averaging kernel (a profile = vector for each single observation). Quantifies the altitude sensitivity of the XCH ₄ retrieval. (see Sect. 3.3)
ch4_profile_apriori	Float	n x m	nanomol per mol, abbreviated ppb, i.e., 10 ⁻⁹	A priori mole fraction profile of atmospheric CH ₄ in ppb. (see Sect. 3.3)
xch4_quality_flag	Byte	n	[-]	Quality flag for XCH ₄ retrieval, 0 = good.
Common parameters for XCO₂ and XCH₄ products: See continuation Table 5				

Table 4: Common parameters for the SCIAMACHY and TANSO XCO₂ and XCH₄ data products generated with ECAs. To be continued via **Table 5**.

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 14
		Version 3 Final
		6 June 2014

Name	Type	Dimensions	Units	Short Description
Common parameters for XCO₂ and XCH₄ products:				
solar_zenith_angle	Float	n	Degrees	Solar zenith angle
sensor_zenith_angle	Float	n	Degrees	Sensor zenith angle
time	Double	n	Seconds	Measurement time
longitude	Float	n	Degrees	Center longitude of the measurement
latitude	Float	n	Degrees	Center latitude of the measurement
pressure_levels	Float	n x k (k = m or k = m+1, see Sect. 3.3)	hPa	Vertical altitude coordinate in pressure units as used for averaging kernels (see Sect. 3.3)
pressure_weight	Float	n x m	[-]	Pressure weights as used for averaging kernels (see Sect. 3.3)

Table 5: Continuation of **Table 4**.

Description of each parameter:

xco2

Main XCO₂ parameter. Retrieved column-average dry-air mole fraction of atmospheric carbon dioxide (XCO₂) in ppm.

xco2_uncertainty

Statistical uncertainty of main XCO₂ parameter: 1-sigma uncertainty of the retrieved XCO₂ in ppm.

xco2_averaging_kernel

XCO₂ averaging kernel (for each observation: vertical profile = vector of dimension *m*).

Represents the sensitivity of the retrieved XCO₂ to atmospheric carbon dioxide mole fraction perturbations depending on pressure (height).

For details please see **Sect. 3.3**.

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 15
		Version 3 Final
		6 June 2014

co2_profile_apriori

A *a priori* mole fraction profile of atmospheric carbon dioxide in ppm needed to apply the XCO₂ averaging kernels.

For details please see **Sect. 3.3**.

xco2_quality_flag

Quality flag for XCO₂ retrieval. 0 = good. 1 = bad.

xch4

Main XCH₄ parameter. Retrieved column-average dry-air mole fraction of atmospheric methane (XCH₄) in ppb

xch4_uncertainty

Statistical uncertainty of main XCH₄ parameter: 1-sigma uncertainty of the retrieved XCH₄ in ppb.

xch4_averaging_kernel

XCH₄ averaging kernel (for each observation: vertical profile = vector of dimension *m*).

Represents the sensitivity of the retrieved XCH₄ to atmospheric methane mole fraction perturbations depending on pressure (height).

For details please see **Sect. 3.3**.

ch4_profile_apriori

A *a priori* mole fraction profile of atmospheric methane in ppb needed to apply the XCH₄ averaging kernels.

For details please see **Sect. 3.3**.

xch4_quality_flag

Quality flag for XCH₄ retrieval. 0 = good. 1 = bad.

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 16
		Version 3 Final
		6 June 2014

solar_zenith_angle

Solar zenith angle (SZA). Angle between the line of sight to the sun and the local vertical. SZA is a positive number (i.e., larger or equal to 0 deg).

sensor_zenith_angle

Sensor zenith angle is the angle between the line of sight from the observed ground pixel to the sensor and the local vertical. The sensor zenith angle is a positive number (i.e., larger or equal to 0 deg; 0 deg for exact nadir (downlooking) observation).

time

Measurement time in seconds since 01.01.1970 00:00:00.

longitude

Center longitude of the measurement. A number in the range -180 deg to +180 deg. 0 deg passes through Greenwich.

latitude

Center latitude of the measurement. A number in the range -90 deg (south pole) to +90 deg (north pole). 0 deg = equator.

pressure_levels

Pressure levels as used for the averaging kernels. Ordered from the bottom of the atmosphere to the top of the atmosphere (i.e., by decreasing pressure).

For details please see **Sect. 3.3**.

pressure_weight

Layer / level dependent weights needed to apply the averaging kernels.

For details please see **Sect. 3.3**.

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 17
		Version 3 Final
		6 June 2014

3.3 How to use the Averaging Kernels (AKs)?

3.3.1 Introduction

In order to compare the satellite-retrieved XCO_2 and XCH_4 data products with model simulations and for inverse modelling of surface fluxes (see, e.g., **/Bergamaschi et al., 2007/**) the altitude sensitivity of the satellite retrievals has to be taken into account. Information on the altitude sensitivity is provided by the satellite XCO_2 and XCH_4 averaging kernels and corresponding CO_2 and CH_4 *a priori* vertical profiles.

Also for validation purposes the averaging kernels have to be considered, see, e.g., **/Wunch et al., 2010, 2011/**, **/Dils et al., 2013/** and TCCON website (in particular https://tcccon-wiki.caltech.edu/Network_Policy/Data_Use_Policy/Auxiliary_Data).

All common variables described in the previous section (e.g., xco_2 , xco_2 _uncertainty, time, longitude, etc.) can be used identically for all GHG-CCI ECA products with the **exception** of the averaging kernels and related parameters, as these parameters are closely related to the retrieval algorithm used.

In this section it is explained how the averaging kernels and related parameters can be used.

How these parameters have been defined depends on the retrieval algorithm used to generate a certain product and it was not possible to fully harmonize their use, i.e., their use depends on the product.

The purpose of this section is to explain how to use the averaging kernels and their related parameters and for which data product which method is recommended.

There are two different averaging kernel (AK) categories:

Depending on product, the AKs are

- “layer-based” (IUP, Univ. Bremen, and SRON products)

or

- “level-based” (Univ. Leicester products).

In the following sub-sections more information on this is given including the information for which product which category is valid.

Note that user can also determine “automatically” or via inspection of the product files which category a given product belongs to:

- For “layer-based” products the vertical dimension of parameter **pressure_levels** is $m+1$, i.e., there is one entry more than for parameter **pressure_weight** (or any of the other parameters with a vertical dimension), which has m vertical entries, i.e., one entry less than parameter **pressure_levels**.
- For “level-based” products all parameters have m entries.

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 18
		Version 3 Final
		6 June 2014

In the following sub-sections the relevant parameters are listed and shortly explained (**Sect. 3.3.2**) followed by detailed explanations of how these parameters can be used for the layer-based AK products (**Sect. 3.3.3**) and for the level-based AK products (**Sect. 3.3.4**).

Important note:

The AK related parameters and how they can be used as described in this document is most interesting for users who want to use different products and prefer to easily switch from one product to another. The main purpose of the common parameters and methods described in this document is to provide the users with the parameters and formulas to do this. However, all products also contain additional parameters, not described in this document, but in the PUGs of the individual products (please see also the Algorithm Theoretical Basis Documents (ATBDs) of the individual algorithms used to generate the individual products). Using these additional parameters (and corresponding formulas) users may be able to obtain somewhat more accurate results (although the differences are expected to be very small).

	ESA Climate Change Initiative (CCI)	Page 19
	Product Specification Document Version 3 (PSDv3)	Version 3 Final
	for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	6 June 2014

3.3.2 Averaging kernel (AK) related parameters

For each single observation (ground pixel) several averaging kernel related parameters are contained in the satellite product files. These parameters are listed in **Table 6**.

For additional information and how to use these parameters see the following two sub-sections.

Parameter Name	Mathematical symbol	Dimension (*)	Unit	Explanation
pressure_levels	p	k	[hPa]	Pressure levels; note: $k = m + 1$ (for layer-based approach) or $k = m$ (for level-based approach)
pressure_weight	pw	m	[-]	Pressure weights for all layers / levels
xco2_averaging_kernel	AK	m	[-]	XCO ₂ averaging kernel
co2_profile_apriori	VMR	m	μmol/mol, abbreviated ppm (10 ⁻⁶)	CO ₂ <i>a priori</i> profile
xch4_averaging_kernel	AK	m	[-]	XCH ₄ averaging kernel
ch4_profile_apriori	VMR	m	nanomol/mol, abbreviated ppb (10 ⁻⁹)	CH ₄ <i>a priori</i> profile

Table 6: Overview of averaging kernel (AK) and related parameters. (*) Here the ground pixel dimension (n , see previous sections, **Table 4** and **Table 5**) is not shown, i.e., here each array is 1-dimensional (a vector). Each element corresponds to one atmospheric level or layer as explained in the following sections.

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 20
		Version 3 Final
		6 June 2014

3.3.3 How to use layer-based AKs?

In this section it is described how the common parameters related to averaging kernels (AKs) can be used to apply the satellite's AKs to model profiles in order to take the altitude sensitivity of the satellite's XCO₂ and XCH₄ retrievals into account.

As indicated at the end of **Sect. 3.3.1**, each product may (or may not) contain additional parameters and corresponding formulas, not described in this document (but in the corresponding PUG), which can be used to obtain somewhat more accurate results for a specific product (although the differences compared to the method described in this section are expected to be small).

For the layer-based approach the AKs and corresponding *a priori* CO₂ and CH₄ profiles are defined for layers and they correspond to layer averages. There are ***m*** layers, which are defined by ***k = m+1*** pressure levels.

The “AK layer-based approach”, which is explained in this sub-section, needs to be applied for the following GHG-CCI ECA products (all IUP, Univ. Bremen, and SRON products):

- CO2_SCI_BESD
- CO2_GOS_SRFP
- CO2_EMMA
- CH4_SCI_WFMD
- CH4_SCI_IMAP
- CH4_GOS_SRFP
- CH4_GOS_SRPR

As already described above:

Note that user can also determine “automatically” or via inspection of the product files which category a given product belongs to:

- For “layer-based” products the vertical dimension of parameter **pressure_levels** is ***m+1***, i.e., there is one entry more than for parameter **pressure_weight** (or any of the other parameters with a vertical dimension), which has ***m*** vertical entries, i.e., one entry less than parameter **pressure_levels**.
- For “level-based” products all parameters have ***m*** entries.

The layer-based approach is also described and used in **/Bergamaschi et al., 2007/**. Here a slightly modified version of their Eq. 2 is shown (here GHG = CO₂ or CH₄):

$$XGHG^{mod} = \sum_{i=1}^m [VMR_i^{apri} + AK_i(VMR_i^{mod} - VMR_i^{apri})] pw_i \quad \text{Eq. (1)}$$

- Here $XGHG^{mod}$ is the desired modelled XCO₂ or XCH₄ value, which corresponds to the satellite XCO₂ or XCH₄ retrievals.
- The sum is over the ***m*** atmospheric layers (located between pressure levels p_i and p_{i+1} with $i = 1 \dots m$). Here pressure is the “normal” or “total” or “wet” pressure (not the “dry pressure”, see below). Here $i = 1$ corresponds to the bottom of the atmosphere and $i = m+1$ corresponds to the top of the atmosphere.

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 21
		Version 3 Final
		6 June 2014

- pw_i is a layer-dependent weight (depending on algorithm/product this corresponds to $\Delta p_i/p_{surf}$ of **Bergamaschi et al., 2007** times a conversion factor for the conversion of wet to dry pressure).
- VMR_i^{apri} is the satellite *a priori* layer-averaged CO₂ or CH₄ volume mixing ratio (VMR) or, more precisely, Dry Mole Fraction (DMF), between pressure levels p_i and p_{i+1} (note: $p_i > p_{i+1}$).
- VMR_i^{mod} is the corresponding value of the model (CO₂ or CH₄) VMR (DMF) between pressure levels p_i and p_{i+1} .
- AK_i is the satellite XCO₂ or XCH₄ averaging kernel for layer i .

Note that in this equation all parameters are coming from the satellite product with the exception of VMR_i^{mod} .

Note that the described approach permits to use all satellite data as they are without the need to manipulate them, e.g., by interpolation. Only the model quantity VMR_i^{mod} needs to be computed.

For illustration and a short overview please see **Figure 1**.

For a modeler the receipt to compute $XGHG^{mod}$ is the following:

- For each satellite observation:
 - o Interpolate the model profiles to the location and time of the satellite observation.
 - o Compute for each satellite layer i , as defined by pressure levels p_i and p_{i+1} :
 - The layer-averaged model (CO₂ or CH₄) VMR (DMF), i.e., VMR_i^{mod}
 - o Apply the formula given above to compute the desired quantity $XGHG^{mod}$ (see also **Figure 1** and **Figure 2**).

Figure 1 and **Figure 2** explain how the parameters as provided via the satellite product files (**Table 6**) have to be used in order to apply **Eq. (1)**.

	ESA Climate Change Initiative (CCI)	Page 22
	Product Specification Document Version 3 (PSDv3)	Version 3 Final
	for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	6 June 2014

How to use „layer-based“ Averaging Kernels (AKs):

Parameters provided via the satellite product files are shown in blue. Modelers have to compute the layer-averaged model VMRs (= gas Dry Mole Fractions (DMF)) **co2_mod** or **ch4_mod** for all layers and use these formulas:

$$\begin{aligned}
 xco2_mod &= \sum_i [co2_profile_apriori(i) + (co2_mod(i) - co2_profile_apriori(i)) * xco2_averaging_kernel(i)] * pressure_weight(i) \\
 xch4_mod &= \sum_i [ch4_profile_apriori(i) + (ch4_mod(i) - ch4_profile_apriori(i)) * xch4_averaging_kernel(i)] * pressure_weight(i)
 \end{aligned}$$

Here the underlying mathematical formula (XGHG = XCO₂ or XCH₄):

$$XGHG^{mod} = \sum_{i=1}^m [VMR_i^{apri} + (VMR_i^{mod} - VMR_i^{apri}) AK_i] pw_i$$

↑
 Modelled XCO₂ or XCH₄
 Sum over *m* atmospheric layers
 e.g., via dimension of array xco2_averaging_kernel

↑
 Model CO₂ or CH₄ layer-averaged VMR (= DMF) for layer *i*
 co2_mod
 ch4_mod

↑
 Satellite retrieval *a priori* layer-averaged VMR (= DMF) for layer *i*
 co2_profile_apriori
 ch4_profile_apriori

↑
 Satellite retrieval AK value for layer *i*
 xco2_averaging_kernel
 xch4_averaging_kernel

↑
 Pressure weight for layer *i*
 pressure_weight

Figure 1: Overview how to compute XCO₂ or XCH₄ (= XGHG) using the „layer-based“ AK method. Additional explanations are given in **Figure 2**.

Parameters for layer-based AKs:




Figure 2: Additional explanations related to the parameters needed to use the “layer-based AK approach” (see also **Figure 1**).

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 24
		Version 3 Final
		6 June 2014

3.3.4 How to use level-based AKs?

For the level-based approach the AKs and corresponding *a priori* VMR (= DMF) profiles are defined on levels (not on layers).

The same parameters (variable names etc.) as provided via the satellite products files are used as for the layer-based approach described in the previous section but with a slightly different implementation to apply these parameters to compute the modelled XCO_2 or XCH_4 .

For the level-based approach all AK related arrays are given for m levels.

The “AK level-based approach”, which is explained in this sub-section, needs to be applied for the following GHG-CCI ECA products (all UoL products, i.e., all “OC” products):

- CO2_GOS_OCFP
- CH4_GOS_OCPR
- CH4_GOS_OCFP

As already described above:

Note that user can also determine “automatically” or via inspection of the product files which category a given product belongs to:

- For “layer-based” products the vertical dimension of parameter **pressure_levels** is $m+1$, i.e., there is one entry more than for parameter **pressure_weight** (or any of the other parameters with a vertical dimension), which has m vertical entries, i.e., one entry less than parameter **pressure_levels**.
- For “level-based” products all parameters have m entries.

For model comparisons and inverse modelling the following method is recommended in order to compute the modelled XCO_2 or XCH_4 .

The equation to apply the level-based averaging kernels to the model data is the same as for the layer-based approach (**Eq. 1**) but with the variables now all on levels, rather than layers. The key point is that the model data (co2_mod or ch4_mod in **Fig. 3**) must be interpolated onto the retrieval pressure levels (p_i). This interpolation should be done with care so as to conserve the total column amounts of $XGHG$.

For illustration and a short overview please see **Figure 13**.

For a modeler the recipe to compute $XGHG^{mod}$ is the following:

- For each satellite observation:
 - o Interpolate the model profiles to the location and time of the satellite observation.
 - o Compute for model data at each satellite retrieval pressure level i the model VMR, i.e., VMR_i^{mod}
 - o Apply the formula given above (**Eq. (1)**) to compute the desired quantity $XGHG^{mod}$ (see also **Figure 13** and **Figure 24**).

	ESA Climate Change Initiative (CCI)	Page 25
	Product Specification Document Version 3 (PSDv3)	Version 3 Final
	for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	6 June 2014

Figure 13 and **Figure 24** explain how the parameters as provided via the satellite product files (**Table 6**) have to be used in order to apply **Eq. (1)**.

How to use „level-based“ Averaging Kernels (AKs):

Parameters provided via the satellite product files are shown in blue. Modelers have to interpolate model-level VMRs (= gas Dry Mole Fractions (DMF)) **co2_mod** or **ch4_mod** for all levels and use these formulas:

$$\begin{aligned}
 xco2_mod &= \sum_i [co2_profile_apriori(i) + (co2_mod(i) - co2_profile_apriori(i)) * xco2_averaging_kernel(i) * pressure_weight(i)] \\
 xch4_mod &= \sum_i [ch4_profile_apriori(i) + (ch4_mod(i) - ch4_profile_apriori(i)) * xch4_averaging_kernel(i) * pressure_weight(i)]
 \end{aligned}$$

Here the underlying mathematical formula (XGHG = XCO₂ or XCH₄):

$$XGHG^{mod} = \sum_{i=1}^m [VMR_i^{apri} + (VMR_i^{mod} - VMR_i^{apri}) AK_i] pw_i$$

Modelled XCO₂ or XCH₄ (points to $XGHG^{mod}$)
 Sum over *m* atmospheric levels (points to $\sum_{i=1}^m$, e.g., via dimension of array *xco2_averaging_kernel*)
 Interpolated Model CO₂ or CH₄ VMR (= DMF) profile for level *i* (points to VMR_i^{mod} , **co2_mod**, **ch4_mod**)
 Satellite retrieval *a priori* VMR (= DMF) for level *i* (points to VMR_i^{apri} , **co2_profile_apriori**, **ch4_profile_apriori**)
 Satellite retrieval AK value for level *i* (points to AK_i , **xco2_averaging_kernel**, **xch4_averaging_kernel**)
 Pressure weight for level *i* (points to pw_i , **pressure_weight**)

Figure 3: Overview how to compute XCO₂ or XCH₄ (= XGHG) using the „level-based“ AK method. Additional explanations are given in **Figure 4**.

	ESA Climate Change Initiative (CCI)	Page 26
	Product Specification Document Version 3 (PSDv3)	Version 3 Final
	for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	6 June 2014

Parameters for level-based AKs:




Figure 4: Additional explanations related to the parameters needed to use the “level-based AK approach” (see also **Figure 3**).

3.4 Algorithm-Specific Parameters

In addition to the common parameters listed above, a full description of all algorithm-specific parameters is provided in the Product User Guide (PUG) of each product.

	ESA Climate Change Initiative (CCI)	
	Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 27
		Version 3 Final
		6 June 2014

4 References

/Bennett and James, 2013/ Bennett, V. and James, S., Data Standards Requirements for CCI Data Producers, ESA Technical Note, Reference: CCI-PRGM-EOPS-TN-13-0009, Issue 1, Revision 1, Date of Issue 24/05/2013, Link: http://46.137.76.174/?q=webfm_send/76, 2013.

/Bergamaschi et al., 2007/ Bergamaschi, P., C. Frankenberg, J. F. Meirink, M. Krol, F. Dentener, T. Wagner, U. Platt, J. O. Kaplan, S. Körner, M. Heimann, E. J. Dlugokencky, and A. Goede, Satellite cartography of atmospheric methane from SCIAMACHY on board ENVISAT: 2. Evaluation based on inverse model simulations, *J. Geophys. Res.*, Vol. 112, D02304, doi:10.1029/2006JD007268, 2007.

/Boesch et al., 2006/ Bösch, H., Toon, G. C., Sen, B., Washenfelder, R. A., Wennberg, P. O., Buchwitz, M., de Beek, R., Burrows, J. P., Crisp, D., Christi, M., Connor, B. J., Natraj, V., and Yung, Y. L.: Space-based near-infrared CO₂ measurements: Testing the Orbiting Carbon Observatory retrieval algorithm and validation concept using SCIAMACHY observations over Park Falls, Wisconsin, *J. Geophys. Res.*, 111, D23302, doi:10.1029/2006JD007080, 2006.

/Boesch et al., 2010/ Boesch, H., D. Baker, B. Connor, D. Crisp, and C. Miller, Global characterization of CO₂ column retrievals from shortwave-infrared satellite observations of the orbiting carbon observatory-2 mission, *Remote Sensing*, 2010.

/Bovensmann et al., 1999/ Bovensmann, H., J. P. Burrows, M. Buchwitz, J. Frerick, S. Noël, V. V. Rozanov, K. V. Chance, and A. H. P. Goede, SCIAMACHY – Mission objectives and measurement modes, *J. Atmos. Sci.*, 56, (2), 127-150, 1999.

/Buchwitz et al., 2011/ Buchwitz, M., F. Chevallier, P. Bergamaschi, I. Aben, H. Bösch, O. Hasekamp, J. Notholt, M. Reuter, et al., User Requirements Document for the GHG-CCI project of ESA's Climate Change Initiative, pp. 45, version 1, 3. February 2011, available on the GHG-CCI website <http://www.esa-ghg-cci.org/> -> Documents or directly via http://www.esa-ghg-cci.org/sites/default/files/documents/public/documents/URDv1_GHG-CCI_final.pdf.

/Butz et al., 2009/ Butz, A. O., P. Hasekamp, C. Frankenberg, and I. Aben, Retrievals of atmospheric CO₂ from simulated space-borne measurements of backscattered near-infrared sunlight: accounting for aerosol effects, *Appl. Opt.* **48**, 3322-3336, doi:10.1364/AO.48.003322, 2009.

/Butz et al., 2010/ Butz, A., O. P. Hasekamp, C. Frankenberg, J. Vidot, and I. Aben, CH₄ retrievals from space-based solar backscatter measurements: Performance evaluation against simulated aerosol and cirrus loaded scenes, *J. Geophys. Res.*, **115**, D24302, doi:10.1029/2010JD014514, 2010.

/CMUG-RBD, 2010/ Climate Modelling User Group Requirements Baseline Document, Deliverable 1.2, Number D1.2, Version 1.3, 2 Nov 2010.

/Connor et al., 2008/ Connor, B. J., H. Boesch, G. Toon, B. Sen, C. Miller, and D. Crisp, Orbiting Carbon Observatory: Inverse method and prospective error analysis, *J. Geophys. Res.*, 113, D05305, doi:10.1029/2006JD008336, 2008.

	ESA Climate Change Initiative (CCI)	Page 28
	Product Specification Document Version 3 (PSDv3)	Version 3 Final
	for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	6 June 2014

/Dils et al., 2013/ Dils, B., M. Buchwitz, M. Reuter, O. Schneising, H. Boesch, R. Parker, S. Guerlet, I. Aben, T. Blumenstock, J. P. Burrows, A. Butz, N. M. Deutscher, C. Frankenberg, F. Hase, O. P. Hasekamp, J. Heymann, M. De Maziere, J. Notholt, R. Sussmann, T. Warneke, D. Griffith, V. Sherlock, and D. Wunch, The Greenhouse Gas Climate Change Initiative (GHG-CCI): comparative validation of GHG-CCI SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT CO₂ and CH₄ retrieval algorithm products with measurements from the TCCON network, Atmos. Meas. Tech. Discuss., 6, 8743-8782, 2013.

/Frankenberg et al., 2011/ Frankenberg, C., I. Aben, P. Bergamaschi, E. J. Dlugokencky, R. van Hees, S. Houweling, P. van der Meer, R. Snel, and P. Tol: "Global column-averaged methane mixing ratios from 2003-2009 as derived from SCIAMACHY: Trends and variability" J. Geophys. Res., doi:10.1029/2010JD014849, 2011.

/Frankenberg et al., 2005/ Frankenberg, C., Meirink, J. F., van Weele, M., Platt, U., and Wagner, T.: Assessing methane emissions from global spaceborne observations, Science, 308, 1010-1014, 2005.

/GCOS 2011/ Global Climate Observing System (GCOS), SYSTEMATIC OBSERVATION REQUIREMENTS FOR SATELLITE-BASED DATA PRODUCTS FOR CLIMATE, 2011 Update, Supplemental details to the satellite-based component of the "Implementation Plan for the Global Observing System for Climate in Support of the UNFCCC (2010 Update)", December 2011, GCOS – 154, 2011.

/GHG-CCI ASR/ Buchwitz, M. et al., Algorithm Selection Report (ASR) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG), Technical Report, 29. Aug. 2012, 2012. Link: http://www.esa-ghg-cci.org/?q=webfm_send/158

/GHG-CCI PVIRv2/ Notholt, J., et al., Product Validation and Intercomparison Report (PVIR) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG), Technical Report, version 2.0, 4 Nov. 2013, 2013. Link: http://www.esa-ghg-cci.org/index.php?q=webfm_send/152

/Guerlet et al., 2013a/ Guerlet, S., S. Basu, A. Butz, M. Krol, P. Hahne, S. Houweling, O. P. Hasekamp and I. Aben, Reduced carbon uptake during the 2010 Northern Hemisphere summer from GOSAT, Geophys. Res. Lett., doi: 10.1002/grl.50402, 2013b.

/Guerlet et al., 2013b/ Guerlet, S., A. Butz, D. Schepers, S. Basu, O. P. Hasekamp, A. Kuze, T. Yokota, J.-F. Blavier, N. M. Deutscher, D. W. T. Griffith, F. Hase, E. Kyro, I. Morino, V. Sherlock, R. Sussmann, A. Galli and I. Aben, Impact of aerosol and thin cirrus on retrieving and validating XCO₂ from GOSAT shortwave infrared measurements, J. Geophys. Res., doi: 10.1002/jgrd.50332, 2013a.

/Kuze et al., 2009/ Kuze, A., H. Suto, M. Nakajima, and T. Hamazaki, Thermal and near infrared sensor for carbon observation fourier-transform spectrometer on the greenhouse gases observing satellite for greenhouse gases monitoring, Applied Optics, 48(35), 6716-6733, 2009.

/Natraj et al., 2008/ Natraj, V., H. Boesch, R. J. D. Spurr, and Y. L. Yung, Retrieval of XCO₂ from simulated orbiting carbon observatory measurements using the fast linearized r-2os radiative transfer model, Journal of Geophysical Research D: Atmospheres, 113(11), 2008.

/O'Dell et al., 2010/ O'Dell, C. W., Acceleration of multiple-scattering, hyper-spectral radiative transfer calculations via low-streams interpolation, Journal of Geophysical Research D: Atmospheres, 115(10), 2010.

	ESA Climate Change Initiative (CCI)	
	Product Specification Document Version 3 (PSDv3)	
	for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	
		Page 29
		Version 3 Final
		6 June 2014

/Parker et al., 2011/ R. Parker, H. Boesch, A. Cogan, A. Fraser, L. Feng, P. I. Palmer, J. Messerschmidt, N. Deutscher, D. W. T. Griffith, J. Notholt, P. Wennberg and D. Wunch, Methane observations from the Greenhouse Gases Observing SATellite: Comparison to ground-based TCCON data and model calculations, *Geophys. Res. Lett.*, 38, L15807, doi:10.1029/2011GL047871, 2011.

/Reuter et al., 2010/ M. Reuter, M. Buchwitz, O. Schneising, J. Heymann, H. Bovensmann, J. P. Burrows: A method for improved SCIAMACHY CO₂ retrieval in the presence of optically thin clouds. *Atmospheric Measurement Techniques*, 3, 209-232, 2010.

/Reuter et al., 2011/ M. Reuter, H. Bovensmann, M. Buchwitz, J. P. Burrows, B. J. Connor, N. M. Deutscher, D. W. T. Griffith, J. Heymann, G. Keppel-Aleks, J. Messerschmidt, J. Notholt, C. Petri, J. Robinson, O. Schneising, V. Sherlock, V. Velazco, T. Warneke, P. O. Wennberg, D. Wunch: Retrieval of atmospheric CO₂ with enhanced accuracy and precision from SCIAMACHY: Validation with FTS measurements and comparison with model results. *Journal of Geophysical Research - Atmospheres*, *J. Geophys. Res.*, 116, D04301, doi:10.1029/2010JD015047, 2011.

/Reuter et al., 2013/ Reuter, M., H. Boesch, H. Bovensmann, A. Bril, M. Buchwitz, A. Butz, J. P. Burrows, C. W. O'Dell, S. Guerlet, O. Hasekamp, J. Heymann, N. Kikuchi, S. Oshchepkov, R. Parker, S. Pfeifer, O. Schneising, T. Yokota, and Y. Yoshida, A joint effort to deliver satellite retrieved atmospheric CO₂ concentrations for surface flux inversions: the ensemble median algorithm EMMA, *Atmos. Chem. Phys.*, 13, 1771-1780, 2013.

/Saitoh et al., 2009/ Saitoh, N., R. Imasu, Y. Ota, and Y. Niwa, CO₂ retrieval algorithm for the thermal infrared spectra of the greenhouse gases observing satellite: Potential of retrieving CO₂ vertical profile from high-resolution FTS sensor, *Journal of Geophysical Research D: Atmospheres*, 114(17), 2009.

/Schneising et al., 2011/ Schneising, O., Buchwitz, M., Reuter, M., Heymann, J., Bovensmann, H., and Burrows, J. P.: Long-term analysis of carbon dioxide and methane column-averaged mole fractions retrieved from SCIAMACHY, *Atmos. Chem. Phys.*, 11, 2863-2880, doi:10.5194/acp-11-2863-2011, 2011.

/Schneising et al., 2012/ Schneising, O., Bergamaschi, P., Bovensmann, H., Buchwitz, M., Burrows, J. P., Deutscher, N. M., Griffith, D. W. T., Heymann, J., Macatangay, R., Messerschmidt, J., Notholt, J., Rettinger, M., Reuter, M., Sussmann, R., Velazco, V. A., Warneke, T., Wennberg, P. O., and Wunch, D.: Atmospheric greenhouse gases retrieved from SCIAMACHY: comparison to ground-based FTS measurements and model results, *Atmos. Chem. Phys.*, 12, 1527-1540, doi:10.5194/acp-12-1527-2012, 2012.

/Wunch et al., 2010/ Wunch, D., Toon, G. C., Wennberg, P. O., Wofsy, S. C., Stephens, B. B., Fischer, M. L., Uchino, O., Abshire, J. B., Bernath, P., Biraud, S. C., Blavier, J.-F. L., Boone, C., Bowman, K. P., Browell, E. V., Campos, T., Connor, B. J., Daube, B. C., Deutscher, N. M., Diao, M., Elkins, J. W., Gerbig, C., Gottlieb, E., Griffith, D. W. T., Hurst, D. F., Jimenez, R., Keppel-Aleks, G., Kort, E. A., Macatangay, R., Machida, T., Matsueda, H., Moore, F., Morino, I., Park, S., Robinson, J., Roehl, C. M., Sawa, Y., Sherlock, V., Sweeney, C., Tanaka, T., and Zondlo, M. A.: Calibration of the Total Carbon Column Observing Network using aircraft profile data, *Atmos. Meas. Tech.*, 3, 1351-1362, doi:10.5194/amt-3-1351-2010, 2010.

/Wunch et al., 2011/ Wunch, D., Wennberg, P. O., Toon, G. C., Connor, B. J., Fisher, B., Osterman, G. B., Frankenberg, C., Mandrake, L., O'Dell, C., Ahonen, P., Biraud, S. C.,

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 30
		Version 3 Final
		6 June 2014

Castano, R., Cressie, N., Crisp, D., Deutscher, N. M., Eldering, A., Fisher, M. L., Griffith, D. W. T., Gunson, M., Heikkinen, P., Keppel-Aleks, G., Kyro, E., Lindenmaier, R., Macatangay, R., Mendonca, J., Messerschmidt, J., Miller, C. E., Morino, I., Notholt, J., Oyafuso, F. A., Rettinger, M., Robinson, J., Roehl, C. M., Salawitch, R. J., Sherlock, V., Strong, K., Sussmann, R., Tanaka, T., Thompson, D. R., Uchino, O., Warneke, T., and Wofsy, S. C.: A method for evaluating bias in global measurements of CO₂ total columns from space, *Atmos. Chem. Phys.*, 11, 20899–20946, doi:10.5194/acpd-11-20899-2011, 2011.

/Yoshida et al., 2013/ Yoshida, Y. Kikuchi, N., Morino, I., et al., Improvement of the retrieval algorithm for GOSAT SWIR XCO₂ and XCH₄ and their validation using TCCON data, *Atmos. Meas. Tech.*, 6, 1533–1547, www.atmos-meas-tech.net/6/1533/2013/, doi:10.5194/amt-6-1533-2013, 2013.

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 31
		Version 3 Final
		6 June 2014

5 Acronyms

Abbreviation	Meaning
ACA	Additional Constraints Algorithm (of GHG-CCI)
AK	Averaging Kernel
ASR	Algorithm Selection Report
ATBD	Algorithm Theoretical Basis Document
BESD	Bremen optimal ESTimation DOAS
CCI	Climate Change Initiative
CRDP	Climate Research Data Package
DMF	Dry Mole Fraction
DOAS	Differential Optical Absorption Spectroscopy
ECA	ECV Core Algorithm (of GHG-CCI)
ECV	Essential Climate Variable
EMMA	Ensemble Median Algorithm
ESA	European Space Agency
FCDR	Fundamental Climate Data Record
FP	Full Physics
FTIR	Fourier Transform InfraRed
FTS	Fourier Transform Spectrometer
GCOS	Global Climate Observing System
GEO	Group on Earth Observation
GEOSS	Global Earth Observation System of Systems
GHG	GreenHouse Gas
GMES	Global Monitoring for Environment and Security
GOSAT	Greenhouse Gas Observing Satellite
IMAP-DOAS	Iterative Maximum A posteriori (IMAP) DOAS

	ESA Climate Change Initiative (CCI) Product Specification Document Version 3 (PSDv3) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 32
		Version 3 Final
		6 June 2014

IPCC	International Panel in Climate Change
IUP	Institute of Environmental Physics (IUP) of the University of Bremen, Germany
MACC	Monitoring Atmospheric Composition and Climate, EU GMES project
MIPAS	Michelson Interferometer for Passive Atmospheric Sounding
NASA	National Aeronautics and Space Administration
NIES	National Institute for Environmental Studies
NOAA	National Oceanic and Atmospheric Administration
OCO	Orbiting Carbon Observatory
OE	Optimal Estimation
PBL	Planetary Boundary Layer
PR	PRoxy algorithm
PUG	Product User Guide
PVIR	Product Validation and Intercomparison Report
RTM	Radiative transfer model
SCIATRAN	Radiative transfer model for SCIAMACHY developed and continuously improved at IUP
SCIAMACHY	SCanning Imaging Absorption spectroMeter for Atmospheric CHartographY
TANSO	Thermal And Near infrared Sensor for carbon Observation
TBC	To be confirmed
TCCON	Total Carbon Column Observing Network
TBD	To be defined / to be determined
VMR	Volume Mixing Ratio
WFM-DOAS (or WFMD)	Weighting Function Modified DOAS

End of document